tory nature. Nussbaum's assumptions as to the nature of the renal secretion are based, as stated above, upon the premise that after ligation of the arteries the Malpighian vascular knots are cut off from the circulation. Adami having found that in spite of breaking off the arterial supply, blood would still seem to pass through at least some of the glomeruli, the question remains whether the conclusions that Nussbaum has drawn from the urea researches are correct? According to Heidenhain the capsule secretes water owing to its own iridescent secretory activity. This activity is excited by urea, which must pass with the blood to some of the glomeruli; hence the increased secretion of water by Nussbaum. Adami also found that diuretics, as sodium nitrate, owe their diuretic effect in part, if not wholly, to direct action on the glomerular epithelium, and not, as Nussbaum has thought to have shown and Heidenhain has considered possible, to its influence upon the cells of the tubuli.

Plethysmographic and Vaso-Motor Experiments with Frogs. By F. H. ELLIS, *Journal of Phys.*, *Foster*, vol. vi., No. 6.—Drs. Bowditch and Warren have shown in the cat, by means of the plethysmograph, that slow, rhythmical, electrical stimulation of the peripheral end of the cut sciatic nerve of the curarized cat causes an immediate dilatation of the vessels of the leg evidenced by a rise in the plethysmographic tracing, while shocks of a certain greater degree of frequency cause an immediate contraction. Dr. Ellis has studied similar effects upon frogs. The results of Dr. Ellis' experiments point to the existence of a vaso-dilator as well as a vaso-constrictor mechanism in the frog. The vaso-motor mechanism is not so active in cold-blooded animals as in warm, and therefore he is not surprised to obtain dilatation in the vessels of the leg of a frog with greater difficulty than was experienced by Bowditch and Warren in producing the same result in the leg of the cat.

Spinal Vaso-Motor Nerves. By Dr. WATERS, *Foster's Journal*, vol. vi., No. 6.

Dr. Waters has made a series of experiments upon the vaso-motor functions of the spinal nerves of the frog. He wished to ascertain whether the distribution of vaso-motor nerves in the various regions of the frog's alimentary canal bore any distinct relation to the different spinal nerves as they are given off from the spinal cord.

From the results thus obtained, I feel justified in concluding that the blood-vessels in separate areas of the alimentary canal are governed by vaso-constrictor fibres, which travel along distinct and determinable spinal nerves, each spinal nerve conveying fibres for a certain area.

ISAAC OTT.