say that, as the result of intubation, they recovered more comfortably.

Extreme restlessness.—This is an important symptom. When a child is continually changing its position and tossing about from side to side, we may feel certain that it is in a dangerous condition. While this symptom is usually associated with the other symptoms, especially that of epigastric recession, I have seen cases where this was by far the most prominent feature. Indeed, in two cases, but for the restlessness, the patient would not have been thought in immediate danger. I am satisfied this restlessness is due to a lack of oxygen, and is dangerous not only for this reason, but also because the patient rapidly wears itself out by its exertions. When, therefore, extreme restlessness is a marked symptom, even if the other symptoms do not appear urgent, it demands intubation for its relief. The relief given in these cases is usually very prompt.

Cyanosis.—When not due to a spasmodic condition or loosened membrane cyanosis is usually a late symptom, and a very serious one. We sometimes notice this symptom where the others are not prominent, but as far as I have observed, it has been preceded by the other symptoms mentioned. I am satisfied that many cases of this type are not cyanotic so much on account of the laryngeal stenosis as because of obstruction in the deeper portions of the respiratory tract, and I think this explains why it is that intubation often fails to give complete relief. The operation is indicated, however, to relieve the stenosis and to enable the child to cough and more effectually clear out the deeper portions.

From this brief study I will draw the following conclusions: The majority of cases recognized early are amenable to treatment by antitoxin. A certain number, however, fail to respond to the remedy. In these cases, if at the end of twenty-four hours a study of the symptoms leads us to the conclusion that the patient is not better, then intubation should be done. In all cases presenting any one of the following symptoms prominently, namely: deep epigastric recession with each inspiration, labored and prolonged expiration, extreme restlessness, spasmodic attacks coming on at intervals, or persistent cyanosis, then intubation should be performed. I am satisfied that in our enthusiastic confidence in antitoxin, we often expect too much from the remedy, and that in cases seen late it would be wise to intubate and administer this remedy rather than give the remedy and wait for its effect. Antitoxin and intubation are in no way antagonistic. Intubation is the complement of antitoxin, to be used only where indicated, and when indicated it is generally a life-saving operation.

INTUBATIONS FOR DIPHTHERITIC CROUP.

A REPORT OF THIRTY-EIGHT CONSECUTIVE CASES WITH THIRTY-SIX RECOVERIES. ALL TREATED WITH ANTITOXIN.

Presented to the Section on Diseases of Children at the Forty-ninth Annual Meeting of the American Medical Association, held at Denver, Colo., June 7-10, 1898.

BY F. E. WAXHAM, M.D.

Professor of Clinical Laryngology and Rhinology, College of Physicians and Surgeons; Professor of Diseases of the Chest, and Laryngology, Clinical School.
CHICAGO, ILL.

Desiring to add my testimony to that of many others regarding the value of one of the greatest remedies of the century, I beg leave to place the following

cases on record. These cases were no less severe nor less dangerous than those operated upon before the days of antitoxin, and the great reduction in mortality can not be attributed to a milder type of the disease.

June 30, 1895, through the courtesy of Dr. L. T. Durbin, I was called to operate upon a boy of 3 years. The tube was worn six days, being coughed out once, but returning dyspnea required its reintroduction. Antitoxin used; bacteriologic examination positive; recovery slow, but complete.

Sept. 2, 1895. Dr. Wm. P. Munn called upon me to intubate a child $2\frac{1}{2}$ years old. The patient was nearly dead from asphyxiation. Antitoxin given; tube ejected several times from exfoliation of membrane below the tube; bacteriologic examination positive; uneventful recovery.

Oct. 6, 1895, I was called by Dr. A. M. Bucknum. The patient was $2\frac{1}{2}$ years old. Antitoxin given; child greatly exhausted from difficult breathing; examination positive; good recovery; tube was worn five days.

Oct. 16, 1895. Dr. J. W. Hawk called me to operate upon a boy 3 years old. Patient greatly exhausted; removed tube on third day, but it became necessary to reintroduce it. The tube was permanently removed on the ninth day; examination positive; antitoxin used; good recovery.

Oct. 23, 1895. Courtesy of Dr. David Thompson. Patient 2 years old; very unfavorable case; nasal cavities, pharynx and larynx involved and the child nearly dead from asphyxia. Antitoxin given and intubation performed; examination positive; tube worn five days; perfect recovery.

Oct. 23, 1895. Courtesy of Dr. C. B. Richmond. Patient 3 years old; pharyngeal and laryngeal diphtheria. Antitoxin given; intubation performed; examination positive; tube worn five days; recovery.

Aug. 6, 1896. Courtesy of Dr. Hugo Mager. Examination positive; antitoxin; recovery; tube worn four days; age 6 years.

Sept. 13, 1896. Courtesy of Dr. P. V. Carlin. Examination positive; antitoxin; recovery; tube worn four days; age 16 months.

Nov. 8, 1896. Courtesy of Dr. Charles Jaeger. Examination positive; antitoxin; recovery; tube worn five days; age 4 years.

Dec. 2, 1896. Courtesy of Dr. Catherine Hayden. Examination positive; antitoxin; recovery; tube worn five days; age $2\frac{1}{2}$ years.

Dec. 31, 1896. Courtesy of Dr. George Packard. Antitoxin; recovery; wore tube four days; age 3 years.

Jan. 12, 1897. Courtesy of Dr. L. E. Lemen. The patient had been sick with croup for three days before the doctor was called; membrane had already invaded the bronchi. Intubation done at once and antitoxin given; examination positive. Tracheotomy became necessary twenty-four hours after intubation on account of membrane in lower trachea and larger bronchi. Death occurred within twelve hours, from bronchial obstruction. Age 4 years. In the light of our present knowledge we should have given antitoxin in double the strength and repeated it more frequently.

Jan. 22, 1897. Courtesy of Dr. Herrick. Examination positive; antitoxin; recovery; tube worn two days; age 3 years.

April 7, 1897. Courtesy of Dr. P. V. Carlin. Positive; antitoxin; recovery; age $2\frac{1}{2}$ years; tube worn four days.

Aug. 14, 1897. Courtesy of Dr. F. E. Warren. Positive; antitoxin; recovery; age 4 years; tube worn five days.

Sept. 30, 1897. Courtesy of Dr. H. L. Taylor. Positive; antitoxin; recovery; age 4 years; tube worn five days.

Oct. 16, 1897. Courtesy of Dr. D. R. Lucy. Positive; antitoxin; recovery; age 2 years; wore tube six days.

Oct. 31, 1897. Courtesy of Dr. P. V. Carlin. Positive; antitoxin; death occurred from broncho-pneumonia thirty-six hours after the operation. Age 16 months; surroundings very unfavorable, people living in a shack of two rooms.

Oct. 31, 1897. Courtesy of Dr. Von der Smith. Positive; antitoxin; recovery; expelled tube on the third day; age 4 years.

Nov. 7, 1897. Courtesy of Dr. Leavitt. Positive; antitoxin; recovery; wore tube five days; age 3 years.

Nov. 23, 1897. Courtesy of Drs. H. B. Whitney and H. C. Snitcher. Positive; antitoxin; recovery; wore tube five days; age 5 years.

Nov. 26, 1897. Courtesy of Dr. David Thompson. Positive; antitoxin; recovery; tube worn five and a half days; age 4 years.

Nov. 30, 1897. Courtesy of Drs. Worthington and Munn.

Positive; antitoxin; recovery; wore tube seven days; age 5 years.

Dec. 13, 1897. Courtesy of Dr. A. K. Worthington. Positive; antitoxin; recovery; age 2½ years; wore tube six days.

Dec. 14, 1897. Courtesy of Dr. H. C. Snitcher. Positive; antitoxin; recovery; age 13 months; wore tube nine days. This was the fourth case of diphtheria in the family, and the second one operated upon.

Dec. 21, 1897. Courtesy of Dr. Frank Dulin. Positive; antitoxin; recovery; age 6 years; wore tube five days.

Dec. 30, 1897. Courtesy of Dr. C. P. Perry. Positive; antitoxin; recovery; age 2 years; expelled tube on fifth day.

Jan. 1, 1898. Courtesy of Drs. McLaughlin and Weaver. Positive; antitoxin; recovery; expelled tube on fourth day; age 6 years.

Jan. 15, 1898. Courtesy of Dr. J. S. Neva. Positive; antitoxin; recovery; age 6 years; tube worn five days.

Jan. 31, 1898. Courtesy of Dr. P. V. Carlin. Positive; antitoxin; recovery; age 2½ years. Tube removed on fifth day.

Feb. 3, 1898. Courtesy of Dr. P. V. Carlin. Positive; antitoxin; recovery; age 6 years; removed tube on fourth day.

Feb. 15, 1898. Courtesy of Dr. P. V. Carlin. Positive; antitoxin; recovery; age 8 years; removed tube on fourth day.

Feb. 20, 1898. Courtesy of Dr. D. R. Lucy. Positive; antitoxin; recovery; age 2 years; removed tube on sixth day.

Mar. 3, 1898. Courtesy of Dr. I. B. Perkins. Age 17 months; positive; antitoxin, five doses, each 1000 units; recovery; removed tube on fifth day.

Apr. 27, 1898. Courtesy of Dr. T. J. Carlin. Positive; antitoxin; recovery; removed tube on fifth day; age 2 years.

May 11, 1898. Courtesy of Drs. Ellsner and Gorsuch. Positive; antitoxin; recovery; removed tube on fifth day; age 6 years.

May 12, 1898. Courtesies of Drs. Warren and Worthington. Positive; recovery; removed tube on sixth day; age 2 years.

May 24, 1898. Courtesy of Dr. P. V. Carlin. Positive; antitoxin; recovery; tube ejected two days after introduction; age 20 months.

In this connection I would allude to an adult, 40 years of age, with diphtheritic croup, to whom I was called in consultation with Dr. W. E. Shotwell. The patient expelled a membranous cast of the trachea and recovered under repeated doses of antitoxin without an operation. Recovery was formerly so very rare in the adult, that I believe this case is worth recording. I should also call attention to a case to which I was called by Dr. P. V. Carlin, March 1, 1898, and which has been excluded from this report, as the patient was moribund and died during the operation, and could not be resuscitated, although tracheotomy was also performed. This is the only case that has been excluded.

To recapitulate: of the thirty-eight cases operated upon, five were under 2 years of age with four recoveries, or 80 per cent.; eleven were 2 years old, with eleven recoveries, or 100 per cent.; six were 3 years old, with six recoveries, or 100 per cent.; seven were 4 years old, with six recoveries, or 85.7 per cent.; two were 5 years old, with two recoveries, or 100 per cent.; six were 6 years old, with six recoveries, or 100 per cent.; and one 8 years old, with one recovery, or 100 per cent.; total, thirty-eight cases, with thirty-six recoveries, or 94.7 per cent.—a mortality of only 5.3 per cent. in a class of cases that formerly was dreaded above all others.

A few years since it was my experience, out of several hundreds of operations to have saved 35 per cent. Why this great reduction in mortality from 65 per cent. to 5.3 per cent.? There is but one answer. It has been due to the use of antitoxin. If my experience stood alone the conclusion might well be doubted, but inasmuch as the great reduction in mortality has been noted by all operators, it must be conceded to be correct. In former years patients died, in the great majority of cases, after intubation or tracheotomy, from the extension of the diphtheritic exudate into the bronchi. At present this rarely occurs. I am fully

convinced that when antitoxin is properly and energetically employed in full doses and repeated, the disease is at once cut short and no further progress occurs. I am also convinced that if a patient dies after intubation from bronchial obstruction due to the presence of diphtheritic exudation, the remedy has either been used late, the extension having taken place before its administration, or it has been used with a hesitating hand and in insufficient dosage, or the preparation has been of uncertain strength. When a child is taken ill with diphtheritic or membranous croup—which I consider identical—the danger is so great that hesitation in the heroic use of antitoxin is almost criminal. I certainly should advise a full dose of 2000 units even for a child, and half the strength for infants, repeating the dose in twelve or sixteen hours. In one of the cases reported, four doses of 1000 units were given an infant 13 months old, and in another 17 months old, five doses of 1000 units without any injurious effect.

Even at this late day we occasionally meet with those who still cling to former methods of treatment, hoping to carry their patients through without subjecting them to the dangers (?) of antitoxin, and who are apparently afraid of the remedy and hesitate about repeating the dosage. I would again and again repeat and emphasize the fact that antitoxin *must* be given early. It *must* be given in full doses and repeated once, twice or thrice, if necessary. My confidence in antitoxin is so great that I sincerely believe that with intubation, aided by the proper and free use of this remedy, death from diphtheritic croup should rarely occur.

DIPHTHERIA AS VIEWED BY THE GENERAL PRACTITIONER DURING THE LAST YEAR.

Presented to the Section on Diseases of Children, at the Forty-ninth Annual Meeting of the American Medical Association, held at Denver, Colo., June 7-10, 1898.

BY ALEX. MCALISTER, M.D.

CAMDEN, N. J.

It is not my aim to present a learned dissertation either on the pathology or scientific treatment of diphtheria. These subjects have been before the profession so constantly during the last three years that such a paper would prove only an intrusion upon your valuable time. What I wish to do is simply to make a few practical observations that must be patent to every general practitioner. Until recently diphtheria ranked as the gravest of the anginas, and the most dreaded of the acute febrile affections. This applies with especial emphasis to the laryngeal types, formerly largely regarded as a distinct disease, but now generally recognized as identical with diphtheria of the larynx and properly treated with heroic doses of antitoxin. Yet the nasal and naso-pharyngeal types with their preponderance of constitutional manifestations and former frightful mortality were scarcely less dreaded both by the physician and parent.

Recent advances in diagnosis and treatment, it seems to me, have left scarlet fever at the head of the anginas and typhoid at the head of the acute fevers in point of morbid possibilities. Even the progress which has been made in diagnosis is to be credited to antitoxin, it being due to the new impulses contributed by serum therapy. In the American Pediatric Society's report on the collective investigations of the