the tendon of the triceps above the olecranon, and then drawn out to a corresponding outward point through a very small incision similar to that made on the inner side. The wire was again reinserted and pushed downward under the skin until it was finally brought out through the initial external incision. The circle once completed, traction was exerted on the wire until I was sure that the two fragments were in perfect coaptation, the latter being easily and satisfactorily accomplished. The ends of the wire were twisted several times and then divided by scissors close to the bone. By this procedure the skin was incised at four points (Fig. 2), the largest incision being $\frac{1}{3}$ of an inch in length.

The coaptation of the fragments was perfectly assured by the traction of the wire. The latter could not slip, the lower part of the wire being inserted through a bony canal, while the upper part passed through the tendon of the triceps and pressed against the bony button of the olecranon (Fig. 1, E). The main advantage of this procedure is, first, that the operation is entirely exarticular, an advantage which, of course, needs no further explanation or praise; and second, passive motion can be instituted immediately after the wiring.

The operation finished, the elbow was put in extension and so maintained by an anterior plaster splint which was fastened by a bandage. Following the operation, the patient had slight pain, which in a few days diminished considerably and finally disappeared on the twelfth day, when patient left the hospital entirely free from pain. The splint was temporarily removed on the third day and each of the following days, passive motion being used, and at the end of four weeks entirely abandoned. The patient has since resumed his work and has had perfect use of his arm and no pain.

Postoperative History.—Patient was thoroughly re-examined Oct. 22, 1905. Form and contour of the right elbow similar to that of the left. The natural obtuse angle formed by the axis of the right arm and forearm is equal to that of the left. Flexion and extension of the right elbow is painless and equal in extent to that of the left elbow. Palpation reveals a perfectly smooth surface; in other words, there is no interruption in the continuity of the periosteum of the olecranon. No pain on active or passive motion. The right elbow joint is functionally normal. Where the wires were twisted the continual mechanical irritation produced a small bursa, which does not inconvenience the patient in the least.

I hope the general practitioner will realize that this procedure is both practical and harmless, and does not require great surgical skill for its performance; furthermore, it gives assurance of a perfect result.

A CASE OF INFANTILE SCURVY.

GEORGE DOCK, M.D.

Professor of Medicine in the University of Michigan.

ANN ARBOR, MICH.

Infantile scurvy is a comparatively rare disease everywhere, though becoming more frequent. Chiefly because of its infrequency it is often unrecognized, and so reports of single cases, even if they add nothing to the real knowledge of the subject, may be useful by calling attention to the existence of the disease when it might otherwise be overlooked. For this reason I report the first case that has been treated in the university hospital.

Summary.—Case of infantile scurvy treated for two months with diagnoses of neuritis and probable anterior poliomyelitis, with sore mouth. Easily recognized as infantile scurvy by ocular proptosis with subcutaneous hemorrhages of the eyelids, characteristic swelling and discoloration of gums, depression of anterior thoracic wall, swelling and tenderness of wrists and ankles; slight rickets, sweating of head; rapid recovery on change of food and use of orange juice.

Patient.—E. B., aged 10 months, admitted to university hospital Oct. 23, 1905. The mother, aged 24, was very small as

Fig. 1.—Dr. Dock's case of infantile scurvy. The photograph was taken after improvement began, and the front of the thorax does not look so much like that of a cadaver after necropsy as it did in the beginning.

Fig. 2.—Skiagraph showing swollen arm.

Fig. 3.—Skiagraph showing swollen leg and foot.

a baby, but has always been well since then. She looks well and is of robust frame. The father, a well-to-do farmer, is 26 years old and in good health. The patient, their first child, was born Dec. 26, 1904. The birth was not difficult.

History.—The mother tried to nurse the baby, but after three weeks was obliged to stop. Milk and water, equal parts, sterilized by boiling for one hour, was next used. Barley gruel and milk sugar were added to the milk from the third week. The child vomited curds and was tympanitic, so a change was made, at 5 weeks, to the following preparation: Milk, 10 tablespoonfuls; water, 24 tablespoonfuls; Mellin's food, 4 teaspoonfuls.

As this was not taken well, the Mellin's food was increased and the water lessened, and up to the time the patient came to the hospital she took the following: Milk, 25 tablespoonfuls, water, 7 tablespoonfuls; Mellin's food, 9 teaspoonfuls. The mixture was heated to 165 F. for about ten minutes. The quantity taken at each feeding was rarely more than 3 ounces, and for some time five feedings were given in twenty-four hours.

The weight at birth was 8½ pounds; at 5 months it was 14 pounds; at 8 months, 15½.

Constipation has been the rule, rarely interrupted for a day at a time with as many as three to six stools in a day. In the end of August there was bright red blood in the stools and then for from two to three weeks the stools were almost black, becoming natural only about October 10.

The lower central incisors were cut at six months, without special difficulty. At the end of August the upper central incisors began to appear. At that time it was observed that the baby screamed when her feet were touched. Soon she ceased to use her legs, but there were variations in the degree of this. Then the back became sensitive to touch and she stopped sitting up in the middle of September. Then the arms became quiet, the left almost entirely so. The right arm was only moved to put the thumb in the mouth, where it was kept most of the time. As the upper teeth were slow in coming through the gums were incised. About the last of August the upper gum became swollen, appearing like a blood blister which subsided slightly at times, though the main swelling never disappeared nor became much smaller. The lower gum became swollen after the upper one.

About October 15 the right upper eyelid swelled suddenly. Eye salve was applied. The swelling subsided, but left a bluish red discoloration. The left upper eyelid became swollen October 23, and when admitted the lid was swollen so it did not open, was deep pink and edematous looking. The head has perspired profusely for a long time. There has never been any sign of laryngismus stridulus.

Status Præsens.—October 24: The baby lies with its right index finger in its mouth. Occasionally cries, but cries at once on being disturbed. Is of fair size; weighs 15 pounds. The head is square; anterior fontanelle 38x34 mm.; soft blowing systolic murmur over it. The scalp is sweating slightly.

The right eye is prominent. The upper lid is slightly swollen and shows a pale blue oval discoloration at the upper margin. The left upper eyelid, which was closed yesterday and bright red, is less swollen; pale red; the skin finely wrinkled. The lids open one-eighth inch. The bulbs move naturally; both are very prominent and seem to look down. The conjunctiva is normal.

The rest of the face is slightly puffy, the cheeks flushed, ears pale. There is a scanty eruption of minute red papules in the scalp and on the abdomen, but no petechiæ.

The lower central incisors are about half way through. The gum around them is slightly swollen, reddish purple, the purple most marked close to the teeth. The upper incisors can not be seen. The gum over them is swollen, forming two oblong oval elevations, the right 1 cm., the left 1.5 cm. long. These are dirty red at the bases, blue at the tops, but the extreme tops are broken and appear as ragged dirty-looking ulcers covered with small granulations. The mucosa of the rest of the mouth and pharynx is healthy looking; the tongue is not swollen; has a very thin white coat.

The thorax is of good size. The sternum and costal cartilages are flat and look as if pushed in, forming a ridge running from the nipples to the flanks, parallel to the margin of the ribs, from which the ridge is separated by a shallow groove. The chondrocostal articulations are firm; very slightly swollen. Heart and lungs negative.

The abdomen is below the level of the ribs. Liver and spleen not enlarged. The back is tender, but otherwise negative.

The left arm lies extended; the right thumb is in the mouth and can be moved away without evidences of great tenderness. The wrists and lower fourths of the forearms are swollen; the skin is not tense, but there is a firm swelling seemingly beneath the deep fascia. The hands are emaciated. The thighs and knees are negative. The upper parts of the tibiae are not distinctly thickened. The lower thirds of the legs and the ankles are swollen, the skin glossy, and at the same time pitted, like an orange peel, with slight bluish yellow discoloration. The swelling is very firm. Touching the arms and legs causes severe crying, with many tears. The legs and feet are kept extended; no effort is made to move them.

Blood: Reds, 4,000,000; leucocytes, 11,700; hgn., 80 per cent. Differential count (Dr. H. A. Freund): Small lymphocytes, 14.2 per cent.; large lymphocytes, 9 per cent.; polymorphous, 75.2 per cent.; transitional, 0.8 per cent.; eosinophile, 0.4 per cent.; basophile, 0.4 per cent. Few microcytes; many macrocytes; few poikilocytes.

Treatment and Course.—Orange juice was begun as soon as the baby was admitted. It took one ounce of a 25 per cent. mixture very well, apparently liking it, before each feeding. It also took the following milk mixture: Cream, 8 fluid ounces; milk, 1½ fluid ounces; lime water, 1 fluid ounce; water, 9½ fluid ounces; milk sugar, 6¾ drams. Heated at 99.5 C. for twenty minutes; then cooled; 5 ounces every three hours.

October 25: The baby is distinctly better. It looks more intelligent and does not cry as much as before. The swelling is entirely gone from the left eye. The blue spot on the right lid is more distinct. The gums are less swollen and less blue. The wrists are smaller; ankles about as before. The milk was increased to 2 ounces. Mutton broth cooked with potato and carrots was given.

October 26: The appearance is still better. Moves hands and does not complain as much when touched. Ankles very little smaller than before. Food changed to: Cream, 8; milk, 4; otherwise as before.

October 28: Looks much better. Moves left hand freely, also feet. Wrists and ankles are smaller, but the left internal malleolus is still surrounded by considerable swelling. Milk changed to 5 ounces; other ingredients as before.

October 30: General appearance better. Moves all extremities actively. Gums less swollen; lateral upper incisors show through. Eyes, ankles and thorax about the same. While in the hospital the bowels were slightly constipated. The stools were of natural color and well digested. There was a positive test for occult blood (Dr. Cowie) on October 25, 26 and 27; negative afterward. The urine was not obtained, but the napkins showed no traces of blood. The temperature ranged between 101 C. on admission, and 98.6 C., usually between 99 C. and 100 C. Weight 15½ pounds.

Patient was discharged, with full directions for raw milk, milked with as much cleanliness as possible; aerated at once, filtered through cotton and kept on ice.

November 1: Returned for examination. Looks stronger. Moves more than ever before. Does not cry when handled. Discoloration on right eyelid clearing up. Eyes are further back than before. Upper gum is still spongy and red, but improving. Ankles and wrists about as before.

November 7: Patient sat up yesterday alone. Better in every way.

November 10: Examined in clinic. Moves extremities freely and sat up alone to-day. Color good. Discoloration of right eyelid has disappeared. Left eye normal. The eyes are not prominent. The thorax is deeper. The gums are still slightly

swollen and red. Does not take orange juice as well as at first. Is taking 6 fluid ounces of milk in 20 ounces of the mixture. Weighs 16½ pounds. Food changed to cream 8 and milk 7 fluid ounces; other ingredients as before. Broths and gruels were also prescribed.

November 23: For two or three days has not taken food as well as before. Vomited once on the 19th, probably as the result of cathartic medicine. Looks stronger. Moves actively. Wrists are thick; ankles smaller than before. No discoloration. No tenderness. Gums natural; four upper incisors out; three below and fourth almost through. Costal cartilages still show groove, but body is more rounded than before. Advised five meals a day.

REMARKS.

Diagnosis.—Among the points of interest in the case diagnosis takes the first place. Like many similar cases, the true nature of this one was long overlooked. During this time the patient was in care of two physicians, in turn, of considerable experience and more than average ability. Neither had ever seen scurvy at any age, and they had only the usual knowledge of the less common diseases of childhood. But lack of actual observation of a similar case can not be a valid excuse in practical diagnosis. All progress in medicine would stop if it were. The physician first consulted discussed the case with a surgeon, who suggested neuritis without seeing the patient. The other suspected anterior poliomyelitis, but, not feeling certain, referred the patient to me. The chief cause of failure was the lack of a thorough examination, the cause of most faulty diagnoses. This was assisted by a failure to compare the findings with descriptions in text-books on diagnosis and practice.

A literary knowledge of the subject does not alone prevent error. Heubner tells us that in half the undiagnosed cases of infantile scurvy he saw in consultation the physicians in charge had a literary knowledge of the disease. Until I stripped the baby and examined its extremities, its mother and a very experienced and critical grandmother were not aware of the striking alteration in the shape of the thorax and the swellings and discolorations of the extremities. Had an examination been made, the first diagnosis of neuritis must have been given up on account of the absence of objective evidences of that disease and the presence of other features that could not be explained by neuritis alone. Then it would have been necessary to consider the possibilities that have caused error in other cases. Osteomyelitis, so often diagnosed, even to the point of causing operation, could hardly have been suspected in this case. In some, where the swelling has been greater and less symmetrical, only a complete examination and thoughtful differential diagnosis could prevent error. Against rheumatism, the relative constancy of the local lesions, the fact that they were not in the joints, the absence of superficial redness and of distinct fever would have been useful. On the other hand, the sweating of the head, if not considered in all its bearings, and the slight elevation of temperature, might easily confuse the diagnosis in this respect. Against sarcoma, the multiplicity and symmetry of the lesions, and the failure rapidly to increase should suffice. Against syphilis with pseudo-paralysis the condition of the parents, the age of the patient, the condition of the skin and mucous membranes, the absence of joint involvement and the great degree of tenderness, would be decisive. Anterior poliomyelitis was an insufficient diagnosis in view of the physical conditions. There was also no history of the onset of that disease, and the description of the course of the pseudo-paralysis was clear and agreed more with that of scurvy

than of infantile paralysis. Rickets would have been a better diagnosis, would have been partially correct, and, if made in time, should have led to a change in diet that would, in all probability, have prevented the full development of infantile scurvy. The case illustrates the importance in all cases of artificially fed infants of looking for rickets and the scurvy so often associated with it.

Critical examination of the mouth should have made it clear that the disease in the gums was not due to the incision, as was supposed, for it was also present in the lower gum that had not been incised. Further study of such anomalies must have led to a correct diagnosis of scurvy.

The proptosis and subcutaneous hemorrhage in the eyelids came on late. Diagnosis made in time should have prevented them. Although the orbital changes are relatively rare, the fact that E. Schlesinger has recently reported a case in which orbital hematoma was the first symptom in a case of infantile scurvy shows the importance of a general knowledge of the process. The article of Schlesinger¹ and that of Irving Snow² give a complete résumé of our knowledge of this interesting symptom. In regard to its incidence, the collective investigation of the American Pediatric Society gave only 22 cases out of 379, positively absent in 110. Heubner saw 4 cases out of 65. Barlow, who first described it, has also seen a case in which orbital hematoma was the only symptom. Snow shows the anatomic feature of the condition, a subperiosteal hemorrhage like the process on other bones, and Cohn's report of a case in which the hemorrhage came on suddenly, during a severe crying spell, suggests a traumatic element that is often present in the hemorrhages on the long bones. The successive affection of the two orbits, as in my case, is very frequent.

Etiology and Results of Treatment.—The study of the causal factors in the case reveals nothing new. There is the history, as in most cases, of a monotonous food, composed of a milk mixture that would not seem well adapted to the patient, and yet it seemed to agree fairly well for a longer time than one would have anticipated. The long-continued use of Mellin's food is of interest, but does not give any positive ground for conclusions. It is interesting to observe that severe gastrointestinal disturbance did not occur. The idea that a severe diarrhea or enteritis had been present was entertained by one of the physicians, but was not borne out by the clear and positive descriptions of the relatives nor by the examination of the stools. The bloody stools were certainly the result of the scurvy, without further implication of the intestines.

As in so many cases, we can not say whether the cause here was some qualitative or quantitative alteration of the food elements or some toxic substance, or all combined, in a patient long inappropriately fed.

Some other details of the feeding are interesting. The patient lived on a dairy farm, the herd being made up of Jerseys with a few Holsteins. As I learned later, the milk had undergone official tests and was good in every respect. When I began the treatment I was afraid to use raw milk, fearing the supply on the one hand, the gastrointestinal condition of the patient on the other. I therefore gave orders to Pasteurize the milk at 160 F., but by an error the method then in use in the ward was followed, and the milk was really heated longer than

1. "Zur Symptomatologie der Barlowschen Krankheit," Muench. med. Wochft., Oct. 24, 1905, p. 2073.

2. "Eye Symptoms of Infantile Scurvy," Archives of Pediatrics, August, 1905, p. 576.

before and at a higher temperature. The only differences, then, were in the different source of the milk, an element that can not be analyzed, the withdrawal of the Mellin's food and the use of orange juice. Before anything else was given a distinct improvement was obvious, and recovery occurred while on sterilized milk, orange juice, and a small amount of potato and carrot broth. So rapid an improvement, of course, is not at all uncommon, and the change as soon as the orange juice is given, along with the curious avidity with which my patient took it, like so many others, seems to prove that it contains something much needed by the scorbutic body.

Skiagraphy.—My efforts to get an x-ray picture of the affected bones were not successful. It was impossible to make an exposure the first day, when the legs were spontaneously kept still. By the next day there was some motion when the noise and light of the apparatus were perceived, so that it was not possible to make a slow exposure with a soft tube that would show fine differentiation of the soft parts. The pictures, of which my colleague, Mr. V. H. Willey, made a number, show the swellings, as well as the appearances of the bones, but nothing more.

X-ray pictures of infantile scurvy have been published by Ausset, Rotch³ and Lenndorff.⁴ Those of Ausset and Rotch are not very good. Lenndorff's is given in a rather diagrammatic reproduction. The author says there were shadows along the femur and the tibia and fibula, and a clearing up of the bone shadow in all the bones. He explains the findings by assuming that the periosteum had had time to exert its osteogenetic function, a belief that is borne out by postmortem examinations. Lenndorff, however, very properly warns against too much dependence on the skiagraphic diagnosis of infantile scurvy, though it may be of assistance in clearing up some of the anatomic changes.

AN ANTIGONOCOCCUS SERUM EFFECTIVE IN THE TREATMENT OF GONORRHEAL RHEUMATISM.*

JOHN C. TORREY, Ph.D.

Fellow in the Department of Experimental Pathology, Cornell
University Medical College.

NEW YORK CITY.

The following communication is intended merely as a preliminary announcement of the preparation of an antigonococcus serum of therapeutic value. It will be followed shortly by an article in which the character and properties of this serum are dealt with in greater detail.

There are several reasons why an antigonococcus serum has not been used heretofore in the treatment of gonorrheal rheumatism. They may be tabulated as follows: First, there is not the slightest indication of an immunity in man to gonorrheal infection acquired by a previous attack; nor in chronic gonorrhea is the individual any the less susceptible to an acute superinfection. Second, there is no well-established case of an infection in animals with gonococcus. Wertheim, by injecting agar and gonococci together into the peritoneal cavity of rabbits, was able to produce a local peritonitis, but not a real infection. Heller's statement that he has produced an ophthalmic gonorrhea in young dogs by

injecting gonococci into the anterior chamber of the eye has not been confirmed. Third, it is generally supposed that gonococci can not be induced to grow luxuriantly under artificial conditions. Finally, most authorities agree with Wassermann and Nikolaysen in concluding that animals can not be immunized to the toxin of the gonococcus, as it is embraced within the cells and only diffuses into the surrounding tissues on its death and disintegration. In other words, according to their view, it is an endotoxin, a bacterial protein.

Although there was apparently slight chance of success, at the suggestion of Dr. Rogers I prepared a serum for a patient with gonorrheal rheumatism of some 15 years' standing, who had been treated in every other way without result. We were agreeably surprised to find the serum effective, not only in that case, but also in a high percentage of others in which it has been tried. These cases are fully described in the preceding clinical report by Dr. Rogers.

The culture which was originally employed in producing the serum was isolated from an acute case of gonorrhea in a male, who had not been treated. Gonococci were very numerous in the exudate, principally within the leucocytes. Smears were made from the discharge on petri dishes of ascitic agar. After an incubation of 24 hours, an almost pure culture of very small, transparent colonies appeared. These colonies were composed of small diplococci in every respect identical in morphology, staining reactions and cultural features with the characteristics of gonococci. The micro-organisms were negative to Gram's stain, would not grow on ordinary peptone agar or broth (repeated tests have been made extending over several months), and died out even on ascitic media in a short time. On ascitic agar (equal parts of ascitic fluid and 2 per cent. glycerin agar) the colonies are small and tend to remain separate. At first transparent and with regular edge, after four or five days the center of the colony becomes granular and the edge spreads out thinly and irregularly. All the diplococci have died in from six to eight days. In ascitic broth the length of life is much greater, from thirty to forty days.

For purposes of inoculation, a very satisfactory medium has been found to be a mixture of rich ascitic fluid and slightly acid beef infusion peptone broth in equal parts. Tubes of about 12 c.c. of the mixture are heated to 60 C. for several hours and tested for sterility. To obtain the best results, the temperature of the incubator should be between 36 and 37 C. Gonococci are killed at a relatively low temperature (at from 40 to 41 C., according to Scholtz). After from 18 to 24 hours' incubation, a slight granular growth appears near the surface and along the side of the tube. This slowly increases in amount until, after six days, the medium is well clouded on shaking. Other strains of gonococci which have been used form, after four days, a rather thick and flaky pellicle. In all the strains the growth after a time sinks to the bottom of the culture tube.

Various modifications of the proportion of the ingredients in ascitic broth have been tried, such as the presence and absence of peptone, increase in the amount of albuminous matter and ascitic fluid, but no combination has proved, on the whole, more satisfactory than the one which has been described.

It is interesting to note that the strain of gonococcus which was first used, after being cultivated for over a year on ascitic media, grows fully as well and produces as effective a serum as when first isolated. This is in harmony with the well-known fact that gonococci, iso-

3. Medical News, 1903, vol. II.

4. Archiv. f. Kinderheilkunde, vol. xxxviii, 1904, p. 160.

* Read before the New York Academy of Medicine, 1905.