ADDENDUM.—The writer thought Dr. Howard Kelly would have been present in the surgical Section of the AMERICAN MEDICAL ASSOCIATION and contended in debate for originality in sigmoidoscopy and rectoscopy had he been entitled to it. But Professor Kelly left Atlanta about twelve or twenty-four hours before the writer's paper was read. Here in the surgical Section was the place to settle the priority of claims and originality of sigmoidoscopy in the presence of surgeons and physicians who could see, think and hear for themselves, and must be our judges. We are reminded of the adage, "He who fights and runs away lives to fight another day." We are of one conclusion, that Professor Kelly did not have, nor has he, the facts to sustain him as the author of sigmoidoscopy and rectoscopy, and that rather than suffer defeat he folded his tent and silently went away.

BALTIMORE, Oct. 14, 1896.

Dear Dr. Carpenter:—I am just writing an article including some account of rectal examinations for the *Centralblatt f. Gynaekologie*, which has the widest distribution of any gynecological journal in the world. In it I shall make careful note of your original work published in the *Cincinnati Lancet-Clinic*, Feb. 23, 1895. I am sorry not to have been able to give the matter more prominence at an earlier date, but it is utterly impossible for me to get through the work which is driving me every hour of the day, and important matters sometimes do have to wait.

Remember that I shall always be glad to see you and show you any courtesy I can here in Baltimore, and when I treat the subject of rectal examinations before my classes your name will always occupy a prominent position as a pioneer.

Sincerely yours, HOWARD A. KELLY.

Cincinnati Lancet-Clinic, Feb. 23, 1895, is the second publication of my sigmoidoscopy, the first publication being June, 1886, and August, 1886, before the Kentucky State Medical Society and in the *American Practitioner and News*.

A SYSTEM FOR THE RATIONAL TREATMENT OF INJURIES BY FOMENTATIONS.

BY CHARLES DEWEY CENTER, M.D.

LATE HOUSE SURGEON PRESBYTERIAN HOSPITAL, CHICAGO, ILL.
QUINCY, ILL.

The treatment of injuries by applications of either hot or cold moisture is not a new procedure. There has been, however, a revival of this practice in the past ten years. Listerism is largely responsible for the renewal, and the development of our bacteriologic knowledge combined with our clearer physiologic insight, has given it the impetus toward the position it now holds. These applications of moist heat or cold were formerly known as stupes, but this term is now confined more especially to applications of turpentine, etc., or perhaps a better way to express it, to medicated fomentations applied to strictly medical cases.

There is no doubt that wound treatment by moisture is sometimes overdone, that it is often used with plain empiricism; that hot moisture is used where cold is indicated and *vice versa*, for very little has been written to guide the practitioner, and a doctor does not collect knowledge of fomentations and their uses by intuitional imbibition any more readily than he does other information of a professional character. Hydrotherapy has much of value concealed among its masses of uselessness and harmfulness. All believe that a kidney well flushed each day is of more benefit to the individual than one clogged with urinary salts. We do not doubt that a clean, well bathed skin is

both more useful and more ornamental than a greasy, dirty one, whose pores are filled with disfiguring black-heads. So wound treatment by moisture is rational and has come to stay, since it is based on physiologic and therapeutic facts.

The healing and cleansing properties of water have long been known; so long, in fact, that animal instinct teaches our four-footed and two-footed friends its virtues. We have merely added the knowledge of the difference in action between the hot and the cold. Dogs have been often seen to resort to water in case of injuries. One instance, reported by a reputable doctor, is on record. His dog, belonging to the spotted coach variety, was run over one day by a wagon. The dog was pushed along in front of the wheel for some distance, lacerating the soft tissues of the legs and the hips, but owing to an unevenness of the pavement, had no bones broken. Howling with pain the animal ran to a stream near by, waded into the shallow water, lay down and could not be persuaded to leave it for about seventy hours. The record is made complete by the statement that no inflammation resulted from the wounds and that the dog made an extremely rapid recovery. Water in some form is being used more and more extensively in fevers. Where once the family doctor sternly and solemnly assured the anxious family that cold water, either internally or externally, meant death to the patient, in one case by driving the fever in and the other by driving it out, there is heard now the command to sponge, or cold pack, or ice pack, when the temperature passes a given point. Permission is given to drink as freely as desired. If water, either hot or cold, or both, are useful in systemic inflammations why not in local ones. In one the disease is general, in the other local, and if there is any virtue in the water one would certainly look for its greatest benefit in the local ailment.

There is one feature of this modern hydrotherapy to bring out more clearly: the difference in the action of hot and cold moisture, and a caution against using either indiscriminately. Moist heat acts on inflamed tissue the same way as the removal of a rubber band from the finger acts on the throbbing sensation present before the removal. It dilates the cutaneous vessels, softens the general mass of tissue, allowing the capillaries and arterioles clogged with red and white corpuscles, to free themselves, permitting the process of osmosis to be enacted backward this time, and causing the intercellular stasis of serum or lymph to disappear.

Either heat or cold at first touch, acts as a constrictor or constringer of tissue, causing momentary pain in the congested area. Either the moderate, continued, hot application or the cold one dilates cutaneous vessels. The intermitting, long continued form of either constricts them. Moist heat promotes activity of the leucocytes by increasing their ameboid movements, and hastens suppuration in this way. Then, too, in an inflammation the fixed tissue cells proliferate and accompany the leucocytes in their tissue infiltration. Pathogenic microbes find the artificially warmed atypical area a fruitful spot, consequently suppuration speedily and easily ensues. Of course, in some cases this tissue infiltration reaches such an extreme that a true avascular area is caused by the pressure, and a patch of necrosis results, a frequent condition in the pathologic condition known as furuncle. Here also moist heat is favorable to resolution, since it hastens the separation of the necrotic tissue.

The cold fomentation, while intrinsically perhaps more valuable than the hot one, will in the practice of many find a more restricted usefulness, since cold must be used when inflammations are young or before the inflammatory process has commenced. The matter of cold for local calor is not a serious one, although patients frequently mislead the doctor by insisting that there is so much heat in the inflamed part. Local temperature is rarely any higher and usually not so high as body heat. It is caused entirely by increased blood supply to the part, in all cases where the systemic temperature does not also rise, the latter being caused by the toxicity of ptomaines, or by the not understood physio-chemic product, fibrinogen.

Application of cold is dangerous in cases of extreme congestion, in strangulated hernia and in any instance where the blood supply is already diminished or shut off. The Thompsonian doctrine that, "heat is life," must be believed and practiced in these cases. Cold acts in much the same way as elevation of a part. It reduces the amount of blood in the deep vessels, assisting the return of this diminished amount by dilating the cutaneous ones. Cold decreases vitality and all the vital processes, a fact recognized in our using heat instead of cold in shock. Where heat hastens suppuration cold retards it by constringing tissue; also by delaying or inhibiting the action of pus microbes.

There is no absolute and undeviating rule for the use of hot and cold fomentations. Idiosyncrasy plays a large part; mental impressions or dislikes a smaller one. Some time ago a case of appendicitis came under my notice. He had been complaining of pain for three days; had a temperature of 102.4 degrees F., there was considerable rigidity of the abdominal muscles, dullness on percussion in the region of McBurney's point. An ice bag was ordered for this spot, absolute quiet enforced and to make him think something was being done for him, as much as for any other reason, he was given benzo-naphthol, 5 grains, three times a day. For six days he lay on his back with the ice bag over his affected locality, and he got well. Some time after, another patient with appendicitis came in. It was in the beginning of her attack also. The ice bag was ordered again, but this time the patient declared it increased the pain, increased the vomiting, and it had to be left off. The only explanation that can be given for these cases, is, that there are exceptions to the rules governing the effect of heat and cold on the vaso-constrictors and vaso-dilators, as well as to all other rules.

One proof of the constricting effect of cold upon deep blood vessels is the test of Brown-Sequard. Ice placed over the lumbar region and kept there, first checks and afterward stops the secretion of the kidneys. The laity know of this effect of cold without knowing how it is obtained. They say a warm bath opens the pores and that the bather will catch cold unless he follows with a cold plunge or sponge, which closes the pores. They also know that cold fomentations, or an ice bag, are good for incipient tonsillitis. On the other hand, the backache of many women is relieved by a hot bath along the spine, the ache being the result of cord anemia and the heat producing a physiologic congestion. The opposite condition and result obtain with the use of ice in cerebral congestion.

Concerning the value of medicated fomentations, and by this term is meant those of boric acid, acetate

of aluminum, sulphate of aluminum, bichlorid of mercury, sulphate of zinc, carbolic acid, lead and opium, or opium alone, this may be said: Theoretically they are all right. The boric acid more nearly fills the requirements of all cases than any other. It has the advantage of not being dangerous. Carbolic acid is decidedly untrustworthy under all circumstances as a fomentation. One case where it was used comes to mind. The patient was a man of ordinary intelligence. He had a boil at outer end of right eyebrow. It was opened, the compartments broken up with the end of finger and washed out with 2 per cent. carbolic solution. Then, having nothing else by me, I put on a hot wet dressing of the same solution, telling him to remove this when he reached home, only four blocks, and to apply fomentations of hot water until he went to bed. The carbolic acid made his aching face feel so good he decided no improvement could be made, so left the dressing on all night. Result, considerable superficial necrosis and sloughing. The acetate of aluminum enjoys a good but narrow reputation. Senn advocates it above all other fomentations, and it is at once non-toxic and seemingly has no effect on idiosyncrasy. Sulphate of aluminum had better be left alone. It has an insidious way of removing the body of skin down to the rete, giving no sign of doing so beyond a slight tingling sensation on the part of the patient. Bichlorid of mercury must be employed very cautiously and its use not continued. Sulphate of zinc, unless used in a very dilute solution, is extremely painful to the patient when used over broken skin. Lead and opium are not used so extensively as they were some years ago. The combination is an incompatible on its face. Whether they became unfashionable preparations or whether some untoward results were obtained we do not know. There is no literature on the subject. The employment of the lead and opium wash in hot fomentations gives this result: The lead acetate hinders the conversion of uric acid into urea and favors the deposit of sodium urate about the joints. If long continued about open joints or open bone injuries, there is an increase in the diameter of the bone, or in fractures an unusual amount of callous, which later becomes infiltrated with osseous material. The opium, on the other hand, gives local results that are very desirable, if it is used in small amounts moderately; it must be used early before any degree of inflammation has been reached, for like cold it raises arterial tension, helps to maintain the blood current and hinders the migration of the leucocytes.

Fomentations of one or another kind are frequently preferable to submersions, because of the greater convenience; because submersion, if long continued, tends by great absorption of the fluid to an edematous, spongy condition of the part, and because fomentations allow the patient greater freedom of position.

In conclusion, it may be said that it is just as necessary to make an accurate diagnosis of the existing condition, for the successful use of fomentations, as it is for the successful use of any drug or of an operative procedure.

527 Lind Street.

The number of medical students in France has materially decreased since the more stringent regulations went into effect this year, dropping from 8,996 to 8,485, a difference of 511. The new regulations were described in the JOURNAL August 22, page 450.