(6) that the pulverulent state of the chromatic substance is difficult to admit as being of pathologic nature and due to disease *intra vitam*, as both this and its precedent condition, the cribriform stage, are so readily produced under the influence of putrefactive agencies. The methods of Nissl were used in the author's researches.

JELLIFFE.

PSYCHIATRY.

CONTRIBUTION A L'ÉTUDE DU SUICIDE DANS LA PARALYSIE GÉNÉRALES (Suicide in General Paralysis). A. Monestier. (*Annales médico-psychologiques*, March, 1900, p. 189).

Monestier comes to the following conclusions: Suicide is quite common in general paralysis, both in the course of the disease and in its initial period. There are, however, a number of cases where the suicide is only apparent, and where the patient does not really wish to put an end to himself. He becomes the victim of his illusions, and loses consciousness of the danger he may run on account of the marked disturbance of a clear perception of the reality of things. In most of the cases of deliberate suicide the suicidal ideas have the characteristics not only of dementia, but also of the delusion which the patient may have, a delusion which is almost always irregular and transitory. This irregularity and transitory character are more or less marked; most frequently the ideas of suicide are sudden and of short duration. It is easy to divert the patient from them and he takes no care to hide them. There are, however, other cases where the means employed to commit suicide differ in no degree from those which other insane employ. The patients seem to have a definite idea of committing suicide as shown by the precautions which they take, by the premeditation of the act, and by the length of their preparations for it. In these cases there is less dementia.

JELLIFFE.

PROGNOSTIC ÉLOIGNÉ DES PSYCHOSES DE LA PUBERTÉ. (Prognosis of the Psychoses of Puberty.) Cullerre (*Archives de Neurologie*, x., 1900, September, p. 246.)

Cullerre has studied 120 attacks of the psychoses of puberty in subjects from fifteen to eighteen years of age—53 males and 57 females. Of these 3 died in the first attack. There were 33 cases of dementia præcox which came on in 18 during the first attack, in 9 during the second, and in 2 in the third. In 4 cases the attack degenerated into secondary systematized insanity with mental enfeeblement. There were 20 cases of periodic insanity, all forms being represented. There were 25 cases of recurrence with varying issue. These recurrences were most frequently benign and were at considerable intervals apart. This group was the most favorable in point of view of ultimate prognosis. There were 9 cases of insanity with a consciousness of the condition, obsessions, or impulsive states. These cases were generally permanent. In 30 cases the patient disappeared from observation after the first attack. The clinical aspects of the first attack, however, would permit us to classify them with the subjects of the preceding groups. The prognosis of the attack in the insanity of puberty is favorable in the enormous proportion of 79%, but the ultimate prognosis of the disease is more serious. The individual who is cured of this form of mental disease runs the risk in the future of recurrences, of dementia præcox, of per-