THE
QUARTERLY JOURNAL
OF
THE GEOLOGICAL SOCIETY OF LONDON.
VOL. XL.

1. *On the GEOLOGY of the SOUTH DEVON COAST from TORCROSS to HOPE COVE.* By T. G. BONNEY, D.Sc., F.R.S., Sec. G. S., Professor of Geology in University College, London, and Fellow of St. John's College, Cambridge. (Read November 7, 1883.)

[PLATE I.]

CONTENTS.

1. Introductory.
2. Torcross to the Start Point.
3. Prawle Point and the adjacent coast.
4. Salcombe to the Bolt Head.
5. The Bolt Tail and the neighbouring coast.
6. Remarks on the Microscopic Structure of the principal varieties of Rock.
7. On some peculiarities of structure in the Slaty Series.
8. Conclusions as to the Relations of the various Rocks in the above district.

1. *Introductory.*

THE wild and picturesque coast which fringes the most southern part of Devonshire has already attracted the notice of not a few geologists. As, however, a detailed list of their writings, carried up to the year 1879, will be found in a valuable paper by Mr. Pengelly, published in the 'Report and Transactions of the Devonshire Association for the Advancement of Science, Literature, and Arts' (vol. xi. p. 319); and as no material addition to our knowledge, so far as I am aware, has since been made, it is needless for me to burden the pages of our 'Journal' with a copious history of the literature of the subject. Suffice it to say that the geology of
Q. J. G. S. No. 157.

A. S. Foord, lith.

Mintern Bros. imp.

STRUCTURE IN CLIFF, TORCROSS.

this district is described in the paper of Sedgwick and Murchison "On the Stratified Deposits of Devonshire" (Trans. Geol. Soc. ser. ii. vol. v. p. 633), and in the "Report" of Sir H. De la Beche. In these two works, which after the lapse of more than forty years are still of the greatest value, the stratigraphy of South Devon is excellently sketched, and the main problems relating to it are clearly enunciated. Briefly stated, the latter amount to this: Does this district indicate a progressive metamorphic action, which increases in intensity as we proceed southwards? or do we find here two distinct and independent series, the southern, a group of highly metamorphosed rocks, true schists, the northern, slates, such as are common elsewhere in Devonshire, with no signs of a transition from the one to the other? Sedgwick and Murchison, if I understand them rightly, incline to the latter view; Sir Henry De la Beche inclines, though hesitatingly, to the former, which appears generally to have found favour with later writers, including Mr. Pengelly himself*. Accordingly the schists of the southern part of this district are regarded simply as altered Palæozoic rocks, possibly of Devonian age, whose upheaval is connected with the series of

Fig. 1.—Sketch-map of the South Devon Coast from Torcross to Hope Cove. (Scale about 3 miles to 1 inch.)

post-Carboniferous disturbances which have affected the south-west of England, and whose metamorphism is probably due to some proximate protrusions of igneous rock, analogous to those yet visible

* This opinion is also adopted, with some hesitation, by Dr. H. B. Holl, in his invaluable paper "On the Older Rocks of S. Devon and Cornwall," Quart. Journ. Geol. Soc. vol. xxiv. p. 400.

in the granite bosses of Dartmoor and of sundry districts of Cornwall. All these observers, however, had been obliged to labour without the aid of the microscope; yet it appeared to me from their statements that the question was one in the settlement of which that instrument would be of exceptional value. Accordingly, last Easter I spent a week in examining the coast section from Torcross to the eastern side of the Salcombe estuary, as far as Portlemouth, and from some distance north of the town of Salcombe on the opposite shore round by the Bolt Head and Bolt Tail to Hope Cove (see Map, fig. 1). Favoured by exceptionally fine weather I had time enough to traverse some of the more difficult parts of the coast section three or even four times, and to collect a considerable series of specimens, the more important of which have been subjected to microscopic examination. The results of these studies I now lay before the Society. As will be seen, there are several matters relating to the stratigraphy of the district upon which I have not been able to come to a satisfactory conclusion. Still I hope to have succeeded in clearing up some difficulties, in narrowing the issues involved, and incidentally in throwing light upon one or two difficult general problems*. In excuse for the incompleteness of some parts of this paper, I may plead that the region presents unusual difficulties. The magnificent cliffs sometimes render a close scrutiny impossible, and their bases are unapproachable except in a boat, and in the calmest weather; while in the neighbourhood of Salcombe, for a considerable distance, the gardens and enclosures around houses make the shore difficult of access. Further, the indubitably metamorphic series exhibits but little variety; it consists of a thick mass of a chloritic rock, overlain and perhaps underlain by a dark mica-schist, and these are bent into such sharp folds that when the green rock and the dark schist emerge from the sea, it is difficult to say which is the upper of the two. Nature, in fact, if I may be allowed the expression, seems to have taken a pleasure in puzzling the geologist by placing the rocks at critical sections, in a nearly vertical position, or by bringing together on opposite sides of a fault rocks, which, though really very different in the amount of metamorphism which they have undergone, to the unaided eye bear a very close resemblance one to another. I may say, however, that I had no hope of clearing up my difficulties by any moderate increase of the time allotted to my work in the field, and am therefore compelled by various duties to leave to others the final settlement of some of the questions proposed in this paper.

2. *Torcross to the Start Point.*

The little village of Torcross, sheltered from the southern gales by a rocky headland, stands on the margin of a singular sheet of water called the Lee†. This is one of those freshwater lakelets

* The nature of these has rendered necessary the occasional repetition of details of stratigraphy already given by earlier authors.

† There is another example (completed, however, by the intervention of man

which have been formed by a bar of shingle and sand damming up the streams issuing from one or more valleys. It is, however, on an exceptionally large scale, and is worthy of more attention from physical geologists than it seems to have received. In the headland bounding the northern side of this sheet of water grey satiny slates, not unlike those in the mid-Devonian series of the northern part of the county, are exposed*. The headland immediately south of Torcross consists of dark slaty and fine gritty beds thrown into great folds with many minor contortions. On its southern side is a thick mass of dark glossy slate, which is quarried for roofing-purposes. All these rocks have undergone little alteration other than mechanical. They are distinctly cleaved, the planes dipping, usually at high angles, to the N. or slightly to the W. of N., and the strike both of the cleavage and the bedding being not far away from E. and W. The petrology of this group is interesting in more than one respect; but this I shall reserve for a separate section (No. 7), in order not to interrupt the description of the general stratigraphy of the district. South of the slate-quarry just named is another valley, now occupied by a small 'Lee,' and at the end of the bar and shore called Bossons Sand comes another headland, on the northern side of which nestles a hamlet bearing the same name. Here, according to the map of the Geological Survey, begins the tract of metamorphic rock which occupies the remaining part of S. Devon, the boundary of this and of the unaltered region running nearly due west by South Pool and Marlborough, until it finally reaches the sea at Hope Cove. At first sight there seems good reason for putting the boundary here on the eastern coast, and much evidence in favour of those who maintain that the rocks between Torcross and the Start Point exhibit progressive metamorphism. Indeed, on my first traverse of the ground, I inclined to the same view; but after carefully reexamining the section from Bossons Sands to some distance south of the village of Hall Sands (similarly situated at the next opening in the uplands), I observed that the rocks in the headland between these two valleys, though locally they might assume the macroscopic aspect of schists †, reverted again and

within the last few years) in the next valley to the south. The Loe Pool, near Helston, is another; and it is possible that the Chesil Bank at Portland may be the remains of a like phenomenon. I think it probable that these shingle beaches are memorials of a period when the coast was depressed slightly below its present level, and thus are bars once formed at the mouths of estuaries and since elevated.

* Dr. Holl (Quart. Journ. Geol. Soc. vol. xxiv. p. 400) considers the whole series of this Torcross district as higher in position than the Plymouth Limestone, and thus equivalent to the upper part of the middle and lower part of the Upper Devonian of the north.

† Throughout this paper I apply the term "schist" (as is my constant practice) only to *foliated* rocks, in accordance with the limitation insisted on by the late Professor Jukes. The lax use of the term by many continental and some English petrologists (by whom it is made to include not only highly metamorphic rocks, but also some where the chemical and mineral changes are but slight) causes much confusion, both in description and (as I have observed) in reasoning.

again to the slaty type, while south of the valley the characteristics of a schist are seen on closer examination to be not only more distinctly marked, but also permanent. Thus in the very first craggy boss N. of the valley at Hall Sands, after a few yards of rock so much decomposed and cut up by quartz veins as to be incapable of giving testimony, we come to a true slate. This can be split up into thin quadrangular or polygonal plates, whose upper and under surfaces have a glimmering satiny lustre, crumbling, or rather cracking and splitting up, when manipulated, after the manner of the fine-grained less flinty slates; while in all the rocks south of the valley, beginning with the very first crag, the surfaces are more waved, and have a more silky mica-like lustre; the thinner edges of fragments can be split up by the knife into rather translucent wavy folia; in short the demeanour of the rock is in all respects that of a micaceous schist.

I examined the crags beneath the hamlet of Hall Sands, until a reef, which projected into the sea, compelled me to mount above them. The rock is fairly uniform in character, being commonly a dull greenish schist, often marvellously corrugated, with "eyes" of quartz and with innumerable quartz-veins, the latter often being best developed in the loops of the corrugations, and seemingly produced by mineral segregation from the darker bands. Sometimes, however, these veins are on a large scale, and traverse irregularly the mass of the rock. They also are frequently crumpled and corrugated to an extraordinary extent, the dip of the schist, if we may trust the alternation of the finer quartzose and micaceous laminae (*i. e.* the foliation planes) as indicative of bedding, is to N.N.W. and is high, being perhaps on the average from 66° to 70° ; but the crumpling and zigzagging of the layers is such that it is most difficult to estimate the dip, though there is a fairly clear intimation of a general E.N.E. strike. Mica-schists (often greatly crumpled and corrugated), varying from a dull greyish green to a leaden or brownish colour, are traversed by the path leading to the Start lighthouse and the rock. About the buildings is a mica-schist, which reminded me of those which so commonly occur on the flanks of the higher ranges in the Alps. Everywhere the schist, in addition to the ordinary aspects of a foliated rock, has a peculiar slickensided look, as though it had been subjected to great compression long after it had become a normal mica-schist.

3. Prawle Point and the adjacent Coast.

Instead of continuing the section westward from the Start lighthouse, it will, I think, save time to describe next the rock which forms the fine headland of Prawle Point, and then to follow the coast back to the Start, on the eastern, and to Portsmouth on the western side. The first rock seen after quitting the village of East Prawle for the Point, is a mica-schist presenting a general resemblance to that at the Start. It forms a sort of scarp running westward, on the southern slope of the hilly upland. Apparently emerging from

beneath this mica-schist comes the mass of rock which forms the headland of Prawle Point, and is exposed in the grand cliffs for a considerable distance on either side. This latter rock varies in colour from a dull green to a light yellowish (epidote) green. Sometimes rapidly alternating bands of different tint appear to indicate a very distinct bedding, at others it is more massive and homogeneous. Generally it has a somewhat decomposed aspect; in the latter case it is often very rotten, passing into a crumbling earthy or sandy material, sometimes of a rusty colour, and forming a pitted carious-looking surface, the rounded hollows being occasionally several inches in diameter and three or four inches deep. It frequently reminded me of a rotten "greenstone tuff;" and now and then I could have supposed myself to be looking at a decomposed diabase. In two or three places I fancied I detected a slightly fragmental aspect, but could never satisfy myself that this was more than an accidental resemblance. In other places, however, the banding of the rock is very distinct, and it has more the aspect of an ordinary schist, though the foliation is seldom or never of a kind that renders it conspicuously fissile. Still, though rightly classed among the metamorphic rocks, it may be well to keep in view the possibility of this having once been a fine basic tuff. As will presently be seen from examination of the microscopic structure, it is rather difficult to give this rock a name. We may continue to call it "chlorite rock," with De la Beche, or adopt "chloritic schist" as being slightly less definite as to its mineral character, remembering that it occasionally becomes rather massive. The dip indicated by the banding, which I think can only result from bedding, at one place rather to the north-west of the Point, was about 45° towards a point between N.N.W. and N.W. At the Signal House it is about 45° or 50° to N.W., or even slightly to the west of this.

The steep cliffs on the eastern side of the Point rendered examination difficult; but, so far as I could see, the general character of the rock for about three quarters of a mile is similar to that already described. I then descended to the shore, the chloritic schist con-

Fig. 2.—Section on shore E. of Prawle Point.

tinuing to form an inland line of crags; the reefs by the water afford the section given in the diagram (fig. 2). Three bands of mica-

schist are intercalated in the chloritic schist, the lowest forming a skerry insulated at high water, the other two exposed on the face of a rocky mass on the shore. The lower of these is about four feet at its thickest part; but both are rather lenticular in form, and in one place the chloritic schist appears to be nipped between two beds of mica-schist. The latter is banded with thin quartzose laminae, and has a more gneissic aspect than the rock at the Start*. I also observed that it was more corrugated than the interbedded chloritic schist. The dip varies from about 20° to 30° to N.N.W., or a few degrees more to the north. These interstratified schists, so far as I can ascertain, are the lowest rocks exposed in this district. I followed these associated schists for some distance, the rock at one place being much contorted and cut by quartz veins several inches thick. Then I ascended the inland cliff, which consists of chloritic schists with one or two bands of mica-schist. After this, keeping along the coast, which now trends nearly N.E., I continued to traverse interbedded chloritic and mica-schists, sometimes much rolled about, but generally striking not far from E. and W., and with the prevalent dip on the northern side at a rather low angle, until (I omit details of minor importance) I lost the chloritic schist on reaching a headland projecting from the Start promontory towards the south. The last band of the chloritic schist appears to strike out to sea, so as to pass beneath the lowest rocks of the Start. My examination of this part of the coast, especially about Lannacombe Mill, and to the east of it, led me to the conclusion that the mass of chloritic schist probably thins, and is certainly split up by bands of mica-schist as it is traced eastward.

Returning to Prawle Point, we can follow the chloritic schist already described along the limit of free cliffs for a distance of nearly half a mile to a projecting headland†. The evidence as to the dip of the strata is sometimes obscure, sometimes variable; but on the whole the dip seems to be high, perhaps about 60° , and the inclination slightly to the W. of N. Beyond the headland the readings are discordant, the beds in one place appearing to plunge to the S., in another to the N.E. The strike, however, seems generally to be about E. and W., and folds are very probable. A short distance to north of the headland the chloritic schist is replaced by mica-schist (the prolongation of that already mentioned as forming the ridge just below Prawle village), the line of junction being almost vertical (fig. 3). The latter is rather greener in colour than usual. In the cove beyond we again find chloritic schist, but on the further side mica-schist occurs again. The latter is wonderfully gnarled and quartz-veined, the zigzagging bands appearing generally to be almost vertical. The former also indicates great disturbance, though

* Probably this is the rock spoken of by De la Beche as "gneiss;" but I should agree with Sedgwick and Murchison in preferring to call it a mica-schist.

† Here I may venture to call attention to the coast scenery. Rarely, except in the recesses of the Alps, have I found a spot so perfect in its solitude or so impressive in its grandeur as on the crags of this headland.

at one place on the southern side the beds dipped pretty clearly to the south. By following these along the shore I could come to no satisfactory conclusion as to the relative position of the two schists, but afterwards, by traversing the little glen which leads down to

Fig. 4.—Chloritic and Mica Schist on shore south of Portsmouth.

the cove, I found that I only lost sight of the mica-schist beneath the turf for a very short distance, and have no doubt that the rock exposed in the headlands on either side of this cove belongs to the same bed, and that the sea has cut across the crown of a sharp arch which brings up the underlying chloritic schist (fig. 3).

This second mass of mica-schist forms a considerable headland, which descends in steep crags to the sea. The rock is much "gnarled," and is slightly greenish in colour, the strike being very nearly E. and W., but the dominant direction of the dip is very uncertain. In the next cove, on the S. side of a small stream which here descends to the sea, we again meet with the chloritic schist. Here also the junction appears to be nearly vertical, but in the cove itself a sharp N.N.W. dip may be noticed in the last-named rock, which is, I think, the lower of the two. Hence, for some distance along the shore, bands of mica-schist occur in the chloritic rock, the former being always the more corrugated and having more quartz-veins. The dips which I observed were now less steep, first to W.S.W., then to S.W. at about 45° , the latter being near the westernmost angle of the coast. The trend of this is now a little E. of N., and we pass over chloritic schist till in a little sandy cove we find a darkish mica-schist pretty clearly underlying it. The annexed diagram (fig. 4) roughly illustrates the relation of the two rocks, and indicates that here, at any rate, the foliation-planes of the mica-schist cannot be trusted as evidence of bedding: for they make a considerable angle with that (indicated by marked mineral changes) in the chloritic schist, and correspond with a faint imperfect cleavage in the latter. Hereabouts the rocks evidently are greatly folded, but the dark mica-schist is pretty clearly shown to be the lower of the two.

Mica-schist, with its foliation-planes generally dipping northward at a high angle, continues as far as the ferry at Portlemouth; and the same rock can be traced for a considerable distance inland up the hill, at least as far as the church; but a little beyond the ferry the chloritic schist seems to set in again and to continue along the shore for some distance. The mica-schist at Portlemouth is evidently thick, and the evidence just mentioned would naturally lead us to place it below the chloritic schist: still after consideration of the whole question, I doubt whether this is not another repetition of the upper schist, the apparent infraposition of that in the cove being due to the folding over of a banded part at the top of the chloritic schist. I was unable, for want of time, to examine the coast north of Portlemouth.

4. *Salcombe to the Bolt Head.*

The ramifications of the coast and the numerous houses and enclosures cause this district to present even greater difficulties than the former. I visited the junction between the true schist and the slaty rock laid down on the geological map to the north of Salcombe, and by making use of a boat, formed some idea of the stratigraphy of the very irregular coast-line between this locality and the town. Landing a little north of the spot where the junction is indicated, I found the rock very imperfectly exposed; but satisfied myself that though, as at Bossons Sands, it occasionally assumed a schistose character, all is indubitably slaty, the cleavage-planes

dipping apparently to points between N. and N.N.W. I place the actual junction, which I have no doubt is a faulted one, about 200 yards S. of where it is drawn on the Geological map; for I succeeded in tracing the slaty rock almost to this point *, and then after a few yards' interval (the rock hereabouts being very imperfectly exposed) found an indubitable mica-schist of rather chloritic aspect. Beyond this, at a slight headland, is a very "gnarled" greenish schist, and at the headland bounding the little bay chloritic schist appears to rise from below the mica-schist.

Along the shore of the inlet leading to Batson, which cuts obliquely across the above-named beds, we have chloritic schist more or less banded with mica-schist, and at last, near the inlet leading to Snapes Farm, these appear to be overlain by a considerable thickness of darkish mica-schist †.

The town of Salcombe is scattered over a headland formed of chloritic schist (fig. 5), of which rock many exposures will be found by searching; but a band of mica-schist occurs near the base, as may be seen in an alley not far from the ferry; and at the upper part of the hill is found a considerable mass of mica-schist, resembling that already described, and clearly overlying the chloritic schist. This is, in all probability, continuous with a mica-schist exposed in a quarry to the west of Shabicombe, on the road to Marlborough.

From Salcombe to North Sands the coast, for the reasons given above, is generally difficult of access. So far as I could ascertain, the rocks are chloritic schist, but in one or two places I observed bands of mica-schist underlying the basal part of the former rock. In the southern flank of the hill, forming the northern boundary of North Sands, the chloritic schist, here pretty well banded, is quarried. The dip is steep, approximately to S. The same rock forms the headland dividing North Sands from South Sands; but by the roadside, low down on its southern face, we find a darkish mica-schist and on the shore at its base chloritic and mica-schists are interbedded and, I think, repeated by minor flexures. The general strike is about E. and W., but it is difficult to say whether the dominant dip is northwards or southwards. On the south side of the little bay called South Sands is a darkish mica-schist, the commencement of the great mass which forms the Bolt Head. The rock is greatly disturbed and "gnarled." At first I inclined to the view that the whole rose up steeply from beneath the chloritic schist, but on repeated examination I fancied I could detect some indication of bedding which gave a southerly dip, and after studying specimens with the microscope, and a review of the whole evidence, I think it more probable that the Bolt-Head mica-schist is only a repetition of the upper schist of the Salcombe hill and is thus a continuation of that already noticed on the eastern side of the estuary. The Bolt Head, so far as I could see, is a mass of darkish mica-schist, the general strike of which appears to be a few degrees

* About 80 yards from a gate, opening from the shore into a field.

† The tide was falling, and mud prevented me from landing.

N. of E. The foliation-planes seem generally to incline rather to the southern side, but the thinner quartz bands among the micaceous laminae are intensely "gnarled," and the quartz veins, by which the rock is often traversed, are commonly completely smashed. I do not think that we can rely much upon the evidence of the minor features of this mass: but the general aspect of the crags—of marvellous wildness and beauty—produced upon me the impression that the headland consisted of a series of grand folds, the loops of which had their axes in a nearly vertical position.

5. *The Bolt Tail and the neighbouring Coast* (fig. 6).

At the Bolt Tail we have, as at Prawle Point, a great mass of chloritic schist. The coast between the Head and Tail may be dismissed in a few words, though the walk from the one to the other is rather a long one (more than four miles as the crow flies, and over very broken ground). As a rule, it is not possible to descend the cliffs to the water's edge; but we can see clearly that the rock is very uniform in character, a dark mica-schist, the interlamination of quartz being very thin, often barely perceptible, while the mass is frequently traversed by quartz veins. There can be little doubt that the general strike is to the east, and the bedding (if we can trust the indications) dips at a high angle to the south. We do not touch the chloritic schist until we reach a spot almost due south of Hope village, where it is seen to rise abruptly from the

Fig. 5.—*Diagrammatic Section near West Side of Estuary, Salcombe.* (Horizontal scale 2 inches to 1 mile.)

sea in the face of a precipitous headland, difficult of approach, and impossible to descend. The two rocks strike a little S. of E., and the mica-schist, so far as can be seen from above, appears to plunge at a very high angle beneath the chloritic rock, but not more than would be perfectly consistent with a slight overthrow in the folding. As we proceed towards the "Tail," the strike becomes E. and W., and about a furlong from the junction the dip is distinctly about 60° N. We see no more of the mica-schist until, as we proceed from the Tail to Hope village by the path along the northern margin of the headland, we note some outcropping beds of it in a cove below; and then, beyond another little cove, where the coast-line bends sharply to the north, we find in the crags of a headland first mica-schist, then chloritic schist, and lastly mica-schist, all seemingly dipping at a high angle to the south. The last-named mica-schist also forms the little "look-out" headland, which divides the two coves, between which extends the village of Hope. It is rather dark in colour and marvellously gnarled and broken, but the foliation-planes are often vertical.

We have only to traverse the rocky shore in the northern cove at Hope for a very short distance to be convinced that, as is indicated in the map, we have returned to the slaty series. Here again the mimicry of foliation, and the other results of the tremendous pressure which the beds have undergone, make it difficult to fix the exact position of the junction, but after repeated examination on more than one occasion, I came to the conclusion, fully confirmed by subsequent microscopic examination, that the beds outcropping on the shore were never more than schistose, while those in the headland itself, wherever I could get at the crag (rendered difficult each time by the state of tide), were true schists, so that the fault has probably determined the northern face of the headland.

Fig. 6.—*Diagrammatic Section near Bolt Tail.* (Horizontal scale 3 inches to 1 mile.)

In the floor of the cove the newer series exhibits considerable variety of mineral character; but in the crags beyond, dark slaty rocks predominate, bearing a general resemblance to the series in the headland between Hall Sands and Bossons Sands. In the distance, beyond Avonmouth, the peculiar sheen of the cliffs recalls to mind the satiny slates seen at Slapton on the east coast, and at Kingsbridge at the head of the estuary.

I postpone the full consideration of the facts recorded in the above descriptions until I have given the results of the microscopic examination of the rocks, merely remarking that the evidence obtained in the field leads us to the following conclusions:—(1) that the foliated and slaty series are quite distinct rocks, the one much older than the other, so that very possibly the foliation of the former was anterior to the deposition of the latter, and that their junction is a faulted one; (2) that both have subsequently been subjected to tremendous lateral pressure, which has bent them into huge folds, cleaving the newer and producing marked effects on the older rock.

6. *Microscopic structure of (A) the Foliated Series, (B) the Slaty Series.*

(A.) Regarded lithologically, the foliated series may be divided into two groups, the mica-schists and the chloritic schists, of which the former are decidedly the more fissile, the latter being occasionally almost massive. Macroscopically the mica-schists do not materially differ; they are of a dull lead-colour, sometimes rather dark, but often relieved by minute spangles of silvery mica, sometimes distinctly laminated with quartz; but not seldom the latter mineral is inconspicuous. The surface often weathers to a rusty brown. The rock, almost everywhere, appears to have been much squeezed after it had assumed its present mineral character. Sometimes it has a peculiar slickensided look, as near the Start; sometimes the surface of the foliation-planes is traversed by minute wavelike wrinklins; sometimes, as especially about the Bolt Head, it is so crumpled as to break up under the hammer into rude prisms with irregularly fluted sides.

Microscopically examined, the differences are merely varietal, and so slight that one description will suffice for the predominant characters of the specimens which I selected for cutting. The chief constituents are quartz, mica, and a variable amount of a minute black mineral.

Quartz.—This occurs in granules variable in size. These are sometimes at least as old as the time when the foliation was produced; at others they have a chalcedonic aspect, as if a secondary product, in interstices due to subsequent disturbance. Cavities, especially in the former, are rather numerous, but extremely minute, so as to require for examination a magnification of at least from 300 to 400 diameters. Some are rounded in outline, some vermiform, a few are irregular, and may possibly be films of some micaceous minerals. Bubbles are extremely rare. Sometimes the cavity is slightly darkened as if by the internal deposit of a ferruginous coating.

Mica.—Of this mineral there appear to be three varieties—(a) a colourless mineral in flakes rarely exceeding .01" in length, exhibiting with the polarizing apparatus a satiny chromatic polarization,

the tints of moderate brilliancy, varying from pale green to pink, and extinguishing when the cleavage-planes are parallel with the vibration-planes of the crossed nicols; probably a hydrous potash or soda-mica; (b) a dark brown mica, associated with, and often rendered almost opaque by, the black dust named above, probably akin to biotite, or a hydrous variety of it; and (c) a pale olive-green slightly dichroic mineral in rather smaller flakes which seems not to extinguish in quite the same positions as the first mineral, and so may be rather a chlorite than a mica.

The white and the dark mica are closely interbanded, sometimes the former, sometimes the latter predominating, in which case the slides become locally only very feebly translucent. The crumpling of these interlaminated minerals is often singularly beautiful, equalling the most complicated arrangements of banded agates (see fig. 7). In this case I have observed that the white mica, when seen with the polarizing apparatus, assumed a peculiar granular aspect, as if the mineral had been somewhat "crumbled" *in situ*. The microscopic structure fully confirms the conclusion, formed in the field, that after the rock had assumed its present mineral character, it was subjected to great lateral pressure, often crushing the minuter laminae of quartz as well as the larger veins of the same mineral, and abruptly bending the more yielding laminae of mica, which were now and then somewhat broken.

The Black Mineral.—This is generally associated with the darker mica, occurring in minute specks, vermicules, and rods, forming fine black bands and streaks, which under a low power are often almost wholly opaque. Among the quartz occasional specks and rare larger grains occur; but even in the latter anything like a definite crystalline outline is wanting. Mr. J. J. H. Teall has kindly undertaken some investigations, experimenting with much care both on prepared slides and on a specimen of the rock powdered, with the results (i) that the colouring matter is principally iron-peroxide, but that there is probably a small amount of carbonaceous matter, and (ii) that the rock contains about 6 per cent. of combined water. Besides the above there is often a little iron-glance, limonite, and a grain or two which we may regard as impure epidote.

Fifteen specimens have been examined, on which I will make a few remarks. One, from the first rock visible on the north of Hall-Sands village (p. 5), *i. e.* from the first outcrop of true schist on the east coast, though much affected by crushing and decomposition (as its macroscopic aspect led me to expect), appears to me to be a true foliated rock. Another specimen, from near a fissure about 100 yards further south, is a mica-schist, like several of those following, but extraordinarily crumpled. A specimen taken from a greenish band in this neighbourhood has, as we might expect, more the aspect of the slides of the chloritic schist presently to be described, but it contains some mica and a fair amount of calcite. It has, as is common with the chloritic schists, a rather "fragmental" look; but this I believe to be the result of subsequent crushing.

Two specimens from different parts of the Start headland call for no special remark. Three specimens, one from the schist to the west of Prawle village, the others from the two principal masses further west, selected to see if they were microscopically as well as macroscopically identical, agree in all essential particulars with one another and with the last two; so also do three from the Bolt Head, one from a quarry just inside the grounds, another from the eastern face of the Head, and a third from near the Mewstone, except that in these, perhaps, there may be a little more of the black mineral. There is therefore nothing to be discovered under the microscope which forbids us to consider all the thick masses of mica-schist one and the same rock. I figure a portion of one from the Bolt Head

Fig. 7.—*Section of Mica-schist from the Bolt Head ($\times 50$).*

(fig. 7), to give an idea of the general structure and the extraordinary corrugations. I examined slices from the bands (A_1 and A_2 of fig. 2) which are undoubtedly below the main mass of the chloritic schist, in the hope of finding something that would characterize a lower mica-schist, but without success. They contain a few grains of a mineral which also occurs in the chloritic schist, and rarely in the ordinary mica-schist; as I shall presently explain, this may be a felspar, but I have doubts as to its nature. I have also examined one of the lower schists from the northern side of South Sands, but it does not help in our attempts at differentiation. Hence, as might be expected, the microscopic evidence as to the position of the mica-schist is inconclusive.

The macroscopic aspect of the chloritic schist has already been described (p. 6). Seven specimens have been cut for microscopic examination. In these three minerals predominate:—(1) the most

abundant, a laminated yellowish green mineral occurring in isolated folia, in grains exhibiting a marked scaly cleavage, and in felted aggregates. Viewed with the polarizer alone, it is seen to be moderately dichroic, a dull green when the lines formed by the cleavage-planes are parallel with the vibration-plane of the transmitted beam, and yellow-green when they are perpendicular to it. We may safely regard this mineral as a species of the chlorite group. (2) A granular mineral, often dusty-looking, giving, when clear, fairly bright tints with the two nicols. This I have no doubt is epidote. (3) Quartz in rather irregular granules, frequently containing microliths of a pale green mineral; some of these, I have little doubt, are flakes of a chlorite, but others appear to be belonites, minute elongated prisms. The comparative thickness of the including quartz grains renders it most difficult to examine their optical characters; but I think that the greatest extinction is when the belonite forms a small angle with the vibration-plane of one of the crossed nicols, so that they are probably hornblende. The quartz also contains numerous extremely minute cavities; these generally seem to be empty, but in a very few I have detected little moving bubbles. (4) A mineral in grains of moderate size, present, but in variable amount, in most of the slides. It has one well-marked cleavage, and another making with it an angle not far from 90° . It may be orthoclase, but I should have expected this under the circumstances to show more indication of decomposition, and I should consider that a few earthy-looking grains occasionally present are more likely to represent a felspathic constituent. The present mineral has considerable resemblance to my specimens of kyanite, which would be a mineral very likely to occur in a rock of this nature. Calcite, grains of iron

Fig. 8.—Section of *Chloritic Schist* from the west side of *Prawle Point* ($\times 50$).

peroxide, and ferrite stains are occasionally and variably present. The quartz granules often have a very elastic aspect, but as they contain frequent microliths of the green constituent, I consider this illusory, the result of subsequent pressure. It will be remembered

that the chloritic schist exhibits less marked crumpling than the mica-schist. The more granular and uniform character of its constituents probably allowed of their being compressed into a smaller space, while the platy laminæ of the mica were necessarily folded by a transverse pressure. It is rather difficult to obtain thin slices of this chloritic schist. Seven slides have been prepared, one from the rock by the signal-house on Prawle Point, another from the cliffs on the west side, and a third from the beds associated with mica-schist on the east (p. 6, in fig. 2). There is nothing in these calling for special remark, except that the belonitic character of some of the microliths is very conspicuous; they pierce the grains of quartz (and of felspar?) like pins in a cushion; there are also bands of secondary quartz, probably filling interstices produced by crumpling. A slide from the quarry near North Sands (p. 10) shows, more than any other, quartz grains like original clastic constituents, and contains some calcite; but, as already stated, I believe the structure to be due to unequal pressure. One from the other side of North Sands, and one from near the Signal on the Bolt Tail, call for no special remark; that from the headland south of Hope, has the felspar-like mineral in greater quantity and in larger grains than is usual.

B. *The Slaty Series.*

Of this group I have examined, microscopically, eight specimens, the majority being selected from critical positions. As an example of one of the normal rocks of the district, I take the silky slate from the quarry on the south side of the Torcross headland. This, cut perpendicular to the cleavage-planes, exhibits a moderately clear, finely granulated ground-mass, divided into minute lenticular streaks by dark lines of variable thickness—the usual aspect, in short, of a fine-grained blackish slate; the dark lines which are parallel with the cleavage-planes are due to the presence of carbonaceous matter, perhaps graphite, with probably some iron oxide. The clearer interspaces (not seldom minutely granulated with similar minerals), when magnified 300 or 400 diameters, appear to be composed largely of very minute scales, rather irregular in form, of a filmy mineral, which on applying the nicols gives pale colour tints; these are probably one of the hydrous micas which seem to be common to many slates, especially the more ‘satiny’ varieties, and to the phyllites; with them are associated extremely minute granules of (apparently) quartz. The slide, as a whole, is darkest when the cleavage-lines are parallel with the vibration-planes of one of the crossed nicols.

A specimen of a slaty rock collected a little to the north of the marshy valley at Hall Sands (just to the north of a stone building), and so one of the nearest to the metamorphic rocks south of the valley, exhibits a structure generally similar to the above, except that the rock has evidently been much crumpled. A sharp zigzag fold traverses the slide, and some of the bands have been broken across, and one part of the fold thrust over the broken ends of the other,

so as to exhibit a structure in one part of the slide which might readily be mistaken for false-bedding*.

A specimen from the curiously corrugated rock at the headland north of the last locality (p. 4) shows a similar mineral structure, but one more conspicuously banded, the slide having been very skilfully cut from the apex of a fold. In this, however, the micaceous constituent occurs in slightly larger flakes. Still, notwithstanding the presence of a hydrous mica, which perhaps we may venture to designate sericite, these rocks differ widely from the normal foliated rocks and, if we restrict the term schist to the latter, they have no claim to that title. These filmy minerals appear to be very readily formed under pressure from damp argillaceous material in a fine state of division, and rocks in which they occur alone differ widely under the microscope, and usually to the naked eye, from the true mica-schists. I call special attention to this because I have noticed in some authors a tendency to confuse these two generally well-defined and widely separated groups of rocks.

A specimen from a rather gritty-looking band in the cliffs south of Torcross shows a minutely granular, somewhat streaky structure, and appears to consist of quartz, white mica, a dusty mineral, probably kaolin, perhaps a little calcite, with ferrite, opacite, and a filmy mineral belonging to the chloritic group. Bands of secondary quartz associated with some of the last-named mineral traverse the slide. Another somewhat similar rock, in which scales of silvery mica are distinctly visible to the eye, from south of Bossons Sands, exhibits a coarser texture under the microscope, consisting chiefly of grains of quartz and flakes of white mica with streaky ferrite and opacite; the quartz is rather full of minute cavities, bubbles being very rare, if not absent. The rock has evidently been much compressed, and some slight amount of mineral change has taken place. I have little doubt that the more conspicuous quartz grains and mica flakes are derivative and indicate the destruction of a gneiss or mica-schist.

Specimens of the greyish, greenish, and blackish rocks from near the fault in the Hope Cove have also been examined; but a very minute description does not seem to me necessary. They are clearly not foliated rocks, though they have been much compressed, and consist of fine aluminous sediments of rather varied nature, associated with some quartz, calcite, and numerous microlithic minerals which it is hopeless to identify with precision. For my present purpose it suffices to say that I have no hesitation in classing these with the slaty group, not with the true schists of South Devon.

7. Peculiarities of Structure in the Slaty Series.

I have already called attention to the remarkable corrugations and schistose structure locally exhibited by members of the slaty

* This structure is described by Prof. Lapworth; *Geol. Mag.*, Dec. ii. vol. viii. p. 343.

series; but there is one case which appears to me so interesting and suggestive that I have reserved it for a separate notice. This is exhibited in the rocky headland upon whose northern slope the village of Torcross rests, and the most striking examples are exhibited in the smooth face of a low cliff rising from the shingly beach within a two minutes' walk from the hotel door. The rock appears originally to have consisted of alternating layers of fine dark mud and silt. The former predominate; the latter, though differing so little in mineral composition or texture as to be with difficulty visible on a freshly broken surface, are more distinct on a weathered one, as their colour changes to a lighter grey than the former. The beds are much folded, and so are inclined at various angles to the direction of pressure. Where this has been normal to the bedding, the different layers have evidently been much compressed, the stripe is remarkably clear and sharply defined, cleavage corresponding with bedding; but in other parts, where the angle between the direction of pressure and the normal to the surface of the bed has been a fairly large one, the results are very remarkable. The stripe is obscured or obliterated, and a new and often more conspicuous structure produced, which is parallel to the planes of cleavage.

Fig. 9.—*Diagrams of Pressure Structure in Banded Rock.*

1. Banded gritty (x, x') and slaty (y, y') rock, cleavage parallel with x, x' . Gritty band about 4 inches thick.
2. Enlarged drawing of part of gritty band between x, x' .
3. Study of a portion of rock, about $1\frac{1}{2}''$ by $1''$, where these gritty bands, which have lain diagonally, have been squeezed out by a nearly horizontal pressure.
4. A small diagonal band of gritty rock squeezed out on one side only.

An examination of Plate I*, which represents a small portion of the cliff face, aided by the smaller diagrammatic figures (fig. 9), may, I hope, render intelligible the following extract from my notebook, and thus render a lengthy description unnecessary. "The whole of this bedded mass exhibits a cleavage-structure, and the more gritty bands are squeezed out into long lenticular streaks, as if the bedding coincided with the cleavage. Where the gritty bands are broadest, there the ends are all 'frayed out' (fig. 9, no. 2); where the banding has been fine and rather close, there it has been entirely obliterated and replaced by this new structure of parallel lenticular streaks or elongated 'eyes' (fig. 9, no. 3). Round these the darker part of the rock bends, just as the folds of mica in a gneiss pass round the lenticular patches of quartz and felspar. Occasionally, perhaps where the materials of the gritty band have been coarser than usual, rather larger lumps have been formed, like the 'eyes' in an augen-gneiss. These are 'drawn out,' as it were, into long flame-like ends, with more distinctly indicated central eyes. They vary in length from about 1 to 3 inches, and in breadth from $\frac{1}{8}$ to $\frac{1}{4}$ inch. In some cases I noticed that the thinner bands were much more squeezed out on one side than on the other, like the feather of a pen, when the pinnules are a little separated (fig. 9, no. 4). Commonly, during the latest stage of the squeezing-out, there appears to have been a slight slipping of one part over another, and a thin film of the darker material has been either infiltrated or dragged in between; this aids greatly in obliterating the traces of the original bedding and in simulating the aspect of a schist." I may also note that in one or two of the coarser gritty bands in this neighbourhood (not "teazed" out as above described) I noticed indications of stratulæ, which might readily be taken as marks of ripple-drifts. I came, however, to the conclusion that these were more probably indications of a rude cleavage—minute splits along which perhaps a slight infiltration had taken place; they made an angle of about 15° or a little more with the well-defined and level surface of the bed. It may be well to bear this instance in mind, for the structure might easily have been mistaken for one of the cases of ripple-drift noticed by Dr. Sorby †.

The peculiar structures described above are, I think, interesting as illustrating some of those seen in foliated rocks. Though the Torcross rock is in no true sense of the word foliated, yet it is often an excellent model of one that is foliated. In regard to rocks of the latter class it is often difficult to say whether foliation corresponds with bedding or is transverse in direction to it, whether it is independent, or only a further development of cleavage; further, an arrangement of the minerals in lenticular streaks with their longer axes parallel is common, especially in gneisses. If the action

* Copied from a photograph. For this I am indebted to my brother, Mr. F. Bonney, who kindly undertook a journey to Torcross to make it. Shingle thrown up by a recent storm had buried the best example, but the one which he secured is nearly as good.

† Quart. Journ. Geol. Soc. vol. xix. p. 401.

of heat, pressure, and alkaline water on these slaty rocks of Torcross had been sufficiently prolonged, we can hardly doubt that the darker laminae would have been converted into bands of mica, the greyer streaks into lenticular aggregates of quartz and felspar. All trace of the original bedding might thus have been lost, and the presumption from the general aspect of the foliation would have been that the original stratification had been parallel to it. Certain portions also of the rock would have closely resembled an augen-gneiss.

In offering these suggestions, I wish to guard myself against being supposed to put forward a general theory of foliated rocks. I merely mention these as a possible explanation of certain cases of foliation, and certain cases of "eyed" structure. I am only prepared to make one general induction, namely, that a lenticular structure in a foliated rock is probably an indication that it has been subjected to pressure in a definite direction, which, though making a fairly large angle with, is not at right angles to, the dominant mineral banding. From what I have seen of foliated rocks in a rather wide experience, I am convinced that very commonly their characteristic structure, speaking in general terms, is parallel with the original stratification, and that there are many augen-gneisses to which the above explanation does not seem to be applicable*.

8. *Conclusions as to the Relations of the various Rocks in the above District.*

From the foregoing remarks it is evident that in this district of South Devon two questions especially present themselves for consideration:—one, the relation of the foliated to the slaty series, the other, the mutual relation of the members of the former series. As regards the former question, we have to decide whether we find in the sections above described—a case of progressive metamorphism, so that we pass in descending order, from true slates to true schists—or two distinct groups, of which the lower was probably metamorphosed before the upper was deposited.

The facts mentioned above in detail prove, I think, that there is no valid evidence of a passage from schist to slate. While I cannot assert that there is a conspicuous fault, as in the Lizard district†, yet the transition from the one class of rock to the other is so abrupt as to be incompatible with the idea of progressive metamorphism, while at Hope Cove and to the north of Salcombe the chloritic schist appears to approach much nearer to the slaty series than at the Start; but apart from the former of these considerations—that we pass from slates and phyllites (to use the strongest term that can be applied to them), from rocks in which the metamorphism can at most be called incipient, to true schists, too rapidly for us to regard

* I may refer in connexion with this subject to a paper read before our Society by the late Mr. G. Poulett Scrope on May 12. 1858. and published, with an admirable illustration, in the 'Geologist' for 1858, p. 361; a paper replete with valuable suggestions, which, as it appears to me, has not attracted the notice which it deserves.

† Quart. Journ. Geol. Soc. vol. xxxix. p. 1.

the change as resulting from an action which has affected both masses to a different extent—there is another which to my mind is conclusive. All the rocks of this region—slates as well as schists—have been greatly folded and bear unmistakable indications of lateral compression. In those which some observers might call phyllites, those which, as described, at first sight resemble schists, the aspect of the rock leads us irresistibly to the conclusion that to this compression is due such modification in mineral structure as they have undergone; but the contrary is the case with the schists, and especially the mica-schists. Folded and corrugated schists are far from uncommon in nature. In some cases the structure of the crystalline constituents indicates that the folding has been the first stage in the process of metamorphism, and that the chemical or mineral change has been subsequent to the mechanical one. In others the former appears in immediate connexion with the latter, subsequent to it, indeed, but in such connexion that the mineral constituents, owing to the plasticity of the mass, were capable of change of position mutually without visible rupture of continuity or sign of crushing; but in a third case it is obvious that the rock, when compressed, was substantially the same that it is now. In the first and second cases we find that the rock does not separate along the different mineral layers, where corrugated, more readily than where these are level; pieces may be struck off from it upon whose broader faces sections of many corrugations of the mineral layers are exhibited. Not so, however, when the lateral compression has been long subsequent to the metamorphism. Then not only do we find evidence of brecciation in the laminae and veins of quartz &c., but also the arched bands of different minerals fall readily apart, and the rock, when hammered, demeans itself in a wholly different manner.

We also meet with abundant confirmatory evidence of the above view in examining the microscopic structure of the rock. The mica-layers have evidently been bent into zigzags subsequently to their crystallization. Sometimes they have gaped, and the aperture has been filled by infiltrated quartz or other minerals; sometimes there is distinct brecciation of the mineral constituents and the like*. Now in the case of these metamorphic rocks of South Devon—the chloritic rock, with its frequent elastic aspect, especially as regards its quartz, the mica-schist with its wrinkled surface, and, especially at the Bolt Head, its peculiar fracture into prisms with corrugated sides, and its wonderful microscopic structure—there is, to my mind, every sign that the metamorphism and the great lateral compression of the rocks were wholly distinct, and the latter long subsequent to the former; and I believe that any petrologist, accustomed both to handle rocks in the field and to examine them with the microscope, will be led irresistibly to the same conclusion.

* I only mention two or three of the most obvious signs. There are others familiar to me which I do not mention, because they would require too long a digression, in order to convince any but the very few who have made a special study of these questions.

I incline further to the opinion that the fault at the junction of the two groups is a considerable one, and that the slaty series of Torcross and Hope Cove was not deposited in immediate succession to the southern schists. The former are generally composed of extremely comminuted materials; even fine grits are not common, conglomerates I never saw. Now the mica-schists especially abound in quartz-veins, and contain layers with fair-sized plates of mica. The former would furnish pebbles of white quartz, the latter mica flakes, not comminuted, because they would be carried for considerable distances without much attrition. Hence, if the slaty series derived much of its materials from the neighbouring schists, I should expect occasional conglomerates of rolled quartz, not unfrequent micaceous grits, and abundant scales of the latter mineral in the muddy beds. Yet in only one of the specimens examined (p. 18) have I found distinctly fragmental quartz and mica, and the micaceous constituents of the most satiny slates or phyllites are so minute as to render it very possible that they are endogenous rather than exogenous*.

The absence of fossils makes it difficult to fix precisely the age of the slaty series of South Devon. Dr. Holl regards the Torcross group as newer than the Plymouth Limestone, and so fairly well up in the Devonian series; but, at any rate in Cornwall, we can trace unmetamorphosed rocks at least as far back as the Ordovician series; so that we may fairly assume any metamorphic rock in a district so near to be considerably earlier than this geological period. I think therefore that we may safely regard the South-Devon schists as Archæan, and as a prolongation of the *massif* so distinctly indicated in the Lizard, to which also belonged the gneiss of the Ed-dystone † and those granitoid rocks which furnished the pebbles in the conglomerate at the Nare Point.

The physical history of the district of which the south-west of England is a fragment seems then to be identical with that of many other regions which have undergone great disturbance from lateral pressure. The first stage—a very remote and doubtless very long one—was the formation of a great mass of more or less crystalline rock, in which perhaps we may discover the usual progression from a basal group of granitoid gneisses through more distinctly foliated gneisses and strong schists, to a well-bedded and more variable series, in which felspar is comparatively rare. The next stage, an uplifting and denudation of this mass; the third, a long period of depression and deposition of sediment; and the fourth, a period of continued lateral pressure, folding, cleaving, and otherwise modifying the rocks affected by it. To this has of course succeeded, as usual, another period, characterized chiefly by denudation, which has lasted down to the present time. A physical history this, which, with

* It will be remembered that in the slaty series of the district N.E. of the Lizard, fragments of metamorphic rocks are found in the associated conglomerate. Quart. Journ. Geol. Soc. vol. xxxix. pp. 9, 10.

† See for indications of the occurrence of crystalline rocks in the bed of the Channel, papers by Mr. A. R. Hunt, in Trans. Devon Assoc. vols. xi., xii., xiii.; also by Mr. Pengelly, in vol. xi. p. 319.

more or less complication, we find in the Scotch Highlands, in North and in South Wales, in the Alps, the Pyrenees, and other mountain chains.

The question as to the mutual relation of the members of the presumed Archæan rocks of South Devon may be quickly dismissed. We have simply to decide whether the chloritic schist is interbedded between two great groups of mica-schist, or whether there is only one thick mass of the latter, with some thin bands of the same rock in the lowest visible part of the chloritic schist. For the reasons given above I consider the latter view to be the more probable, and regard even the mass of the Bolt Head as simply the upper mica-schist repeated by an extremely sharp fold, or rather series of folds.

If I were to attempt to correlate the South-Devon rocks with others of Archæan age, I could only say that they cannot, in my opinion, be referred to the earlier portion of it. In all Archæan rocks which I have examined, where there is any thing like a full development, the rocks become, broadly speaking, more distinctly bedded, more variable in their mineral character, more markedly foliated as we ascend. The threefold division mentioned above (I do not say that further subdivision is not possible) is very recognizable in the Alps; and this—into the group of granitoid gneisses, the group of more fissile gneisses and strong schists, and the group of variable well-bedded schists—suffices for my present purpose. The rocks of South Devon remind me most of members of the last group. I cannot find an exact parallel to them elsewhere in England; but so far as lithological evidence (and we have no other obtainable) avails us, I should consider them to belong to this third group, to be not very far away in geological position from the schists of the Lizard, but, if any thing, a little more recent.

The above conclusions, if established, suggest one or two inferences of a more general nature. One as regards the physical history of a large adjacent area of country. I have indicated the successive changes which the rocks of South Devon have probably undergone. The great period of disturbance—the formation of the boundary fault, followed by the folding and the various results of lateral compression, including, in all, a long series of events—was doubtless subsequent to the deposition of the Carboniferous series as a whole. The flexures &c. of South Devon form parts of the great series of flexures, affecting all rocks, belonging and prior to that age, which can be traced from Belgium to the South of Ireland. It would seem to have been a wrinkling of the earth's crust, scarcely less important than that to which the existing Alpine chain is due. We can hardly help associating with its phenomena the great granitic intrusions and the other igneous masses of Cornwall and Devon, which fringe the north of the most highly compressed area, much as granitic and felsitic bosses now stud the southern margin of the Alps*. In short,

* It may be worth noting the apparent absence of intrusive igneous rocks in the district described in this paper. There is a greenstone dyke near Torcross; but I found no igneous rock in any other place to the south of it. Yet in the Lizard district, where the beds are less sharply folded and there is less evidence

as perhaps will have been inferred from some of my preceding remarks, the phenomena of the district described in this paper are eminently those indicative of a mountain chain*. They record more than one chapter in the history of mountain-making. I can hardly understand the production of those great parallel flexures which affect all south-western England, and especially this district—flexures whose undulations become broader and flatter as we trace them northwards, without assuming the existence to the south of them of a great axial mass of much harder rock, to the pressure of which these sharper crumplings are due. On comparing a section of South Devon with a series of sections across the Alps we cannot fail to be struck with the resemblance between them, except that in the former we do not find the crystalline nucleus rising up as the central mass of the chain, often overtopping and apparently overlying the later deposits. But when we look to the south, beyond the South-Devon and South-Cornish schists, we find at the Eddy-stone—we find probably in the shallower parts of the neighbouring sea—we certainly find yet further south in the Channel Islands and on the adjoining coast of France—large masses, projecting from the ocean, of much more coarsely crystalline rock. Is it possible that these may be the foundation-stones of a great mountain region, of whose outer zone South Devon and South Cornwall are the last remnants—the relics of a long-vanished highland region of Archæan rock? If so, the mountain-making—at any rate that with which we are now concerned—occurred after the Carboniferous, and prior to the Triassic epoch,—was, in short, one of the closing incidents of the Palæozoic volume of geological history. Yet the Channel sea now occupies the place where this range was upraised, and of its ruins some fragments only remain *in situ* †. These remarks may seem to belong rather to the poetry than to the prose of science; but I think that any one who will spend a few years in studying the structure of rocks and of mountain-masses will see that the idea conveyed by them is something more than a baseless flight of imagination‡.

of compression, intrusive igneous rocks abound. This is, of course, what was to be expected.

* See also in illustration of this the sections in Dr. Holl's paper, especially fig. 1, p. 409, and fig. 5, p. 438.

† A thorough study of the rocks composing the Triassic conglomerates of Devonshire would give, I am sure, most interesting results. Whoever undertakes it must be a resident in Devonshire; for the field-work will take no little time; he must be a skilled petrologist, or his results will be worse than useless; and he must have a special knowledge of the rock masses of Devonshire and Cornwall. Among the many excellent geologists and enthusiastic students of the West of England is there no one to undertake this task, and endeavour to replace the covering which has been stripped from the granitic bosses, and rebuild the hills which the sea has destroyed?

‡ There is another more general conclusion to which I may call attention, namely, the importance of pressure as modifying rock structure. Rocks subjected to much pressure, folded, and squeezed out of shape, appear to undergo more micro-mineralogical change than similar rocks when not so maltreated. Perhaps it is hardly too much to say that the difference between a satiny slate or phyllite and an ordinary shale is due even more to the action of

EXPLANATION OF PLATE I.

Structure imitative of foliation produced by pressure on a stratified rock without important mineral change.

The mass of rock figured is about four feet high. The cleavage-structure is vertical; the beds (of alternating dark mud and paler silt) are bent into rather irregular curves pointing downwards, and are separated as described in the text. The paler parts of the figure represent the 'streaks' and 'eyes' of the silty rock, made sometimes more distinct by aggregations of quartz.

DISCUSSION.

Mr. J. STARKIE GARDNER stated that the Eocenes of freshwater origin of the south of England were formed from granitic materials, and not from flints, the Chalk not being then upheaved; and these facts seemed to lend support to the author's conclusions.

Rev. G. F. WHIDBORNE said that somewhat similar beds at Hope Nose contained the fossils of the Middle Devonian. He asked if the rocks dredged in mid channel by Mr. Hunt at all resembled those of the coast.

Rev. E. HILL asked if the gneiss mentioned by Sir H. de la Beche resembled any rock of the Lizard. He agreed with the author that the lenticular structure described is such as would be produced by the oblique crushing of a mass. He stated that in the Channel Islands none but the most highly metamorphic rocks occur.

Prof. T. M^cK. HUGHES pointed out the close resemblance of the gnarled beds of Prof. Bonney's post-Archæan slates to the gnarled beds of Amlwch, which he (Professor Hughes) had described as not truly metamorphic, but only mechanically altered rock. He thought that the lowest Archæan of Britain and France was sometimes granitoid rock, sometimes schist, and that they alternated and passed into one another, so that no general order could be established between granitoid rock and the lowest schists. In Brittany he had observed such alternations, especially in the remarkable sections along the river Rance, near Dinan.

He called attention to the thickening of sediment from north to south, *i. e.* towards the English Channel, in the two groups which next succeeded the Archæan, viz. the Lower Cambrian and the Devonian-Carboniferous, and thought that these facts must be taken into consideration when speculating upon the probability of the

pressure than to mineral composition or geological age. Such rocks as the phyllites of the Ardennes and of the Alps, not to name others, have, as a rule, little claim to be classed with the true schists—the foliated rocks. They have, indeed, taken a step in that direction, but it is only a small one, and their alteration should be classed with the phenomena of mechanical rather than of chemical metamorphism, although the latter has operated to a slight extent. Still, though pressure is an important agent, it is not, as these words imply, the only agent; and when marked regional metamorphosis prevails, when we are considering great masses of truly foliated rock (gneisses and schists), we seem to require something more. Are we justified in asserting that the other necessary agencies operated more freely in Archæan times than they have subsequently done?

former existence of Archæan mountain land over the area now occupied by the Channel.

Dr. HICKS objected to Prof. Hughes's view that schistose rocks such as those exhibited alternate with the granitoid series. He agreed with the author in referring the Devonshire rocks to the newer of the Archæan series. The same succession had been described by Barrois in the North of Spain.

THE AUTHOR agreed with Mr. Starkie Gardner that many of the materials of the Eocene and also of the Mesozoic rocks were derived from granitoid materials. In reply to Mr. Whidborne he stated that the rocks dredged by Mr. Hunt in the Channel were generally of a much older type than those of South Devon. The so-called gneiss near Prawle Point did not resemble any rock at the Lizard. In reply to Prof. Hughes he stated that he had never seen schists of the type now described by him alternating with granitoid rock, and that the two varieties of rock in South Devon, so much alike to the eye, yet really so different, formed a good parallel to the cases in Anglesey which had so misled Prof. Hughes.