be in proportion to his practical experience and the careful study of his cases; we must not lose sight of the fact that in the great majority of instances early diagnosis is dependent on the family physician rather than on the skilled specialist.

As has been said, I have been personally impressed with the value of diagnostic excision. Whether or not this is wise there can be no question as to the absolute value of the most painstaking and far-reaching microscopic study of all tumor tissues after removal. This study should be made in direct connection with the history of the case and the clinical examination of the patient, and the findings should be verified by the operator himself. In the one column of his case-book may be placed his provisional clinical diagnosis; in another the pathologist's report. A comparative study of these will be productive of the greatest good. On the one hand we have the family and life history of the patient—the onset, duration and course of the growth; its general characteristics, resistance, "feel," tenderness, mobility or fixation; the condition of the skin or mucous membrane, of the deeper parts and glands; the blood examination if you like; in a word, all possible factors which can lead to provisional diagnosis. On the other hand lies the pathologist's report. Constant, comparative study of the two will lead to an increasing probability of accuracy in diagnosis.

I do not know that we can at this time go beyond this. In the smaller cities there is a great and increasing need of careful and capable pathologists. The accuracy of our work must be determined by them. Not long ago a well-known surgeon of Boston said to me as we were going through the superb Municipal Hospital there: "When Councilman came to this hospital he revolutionized our methods; we surgeons found that we had to live up to him and his work." We, whose lot is cast in the smaller places, must strive without ceasing for better laboratory work. Our laboratories must be consolidated, enlarged and endowed. It is not enough that the clinicians in the great centers should be in close touch with the pathologists; those in the minor cities and towns must enjoy a like privilege. Happily, facilities for the better training of pathologists are increasing rapidly.

In a recent presidential address before the Association of American Physicians, Professor Welch, the distinguished scientist, compared the opportunities enjoyed in this country by the men who desire to study the branches of scientific medicine and the opportunities open to those who wish to become proficient in clinical medicine and surgery. There are laboratories in this country, equal to any in the world, that are open to young men who have graduated and who wish to become teachers of anatomy, physiology or pathology. In these laboratories a young man must serve his apprenticeship, but Professor Welch says that his promotion is sure. May I not ask you whether (and this especially in the smaller places) we as clinicians recognize as we should the value and the importance of the work which these scientists do, and whether we sufficiently aid in making it possible for them to devote their lives to the purely scientific side of our calling? It is in this direction that advance is to be made, and it is in this direction that our duty lies; a duty seen by such wise and earnest men as Roswell Park, whose Buffalo Laboratory for the study of cancer attracts the attention and admiration of the medical world. Not all of us can create such laboratories, but we can all do some-

thing toward a better appreciation of the importance and value of the most painstaking, thorough and far-reaching laboratory investigation.

BIBLIOGRAPHY.

1. *La Presse Médicale*, Aug. 27, 1898.
2. *Arch. f. Verdauungs-Krankheiten*, 1900, vi, p. 25.
3. *La Semaine Médicale*, March 6, 1901.
4. *Medical Review of Reviews*, April, 1900.
5. *Memorabilien*, 1895, vol. xv, new series, p. 203.
6. *Boston Med. and Surg. Jour.*, May 16, 1901.
7. *The Microscope and the Early Stages of Cancer*. *Archives of Surgery*, 1897, viii, 238.
8. *Transactions Am. Surg. Association*, 1898.

SOME PHASES OF MALARIA.*

J. B. McFELROY, M.D.,

STOVALL, MISS.

On February 12, of this year, I was called to come quickly to see N. M., who the messenger said was speechless. He was a well-developed, strong negro man, 60 years of age, his family history good. He had smallpox about thirty years ago, since which time he has not been sick enough to require the services of a physician. He has never had syphilis, gonorrhea, nor rheumatism. He has lived in the Mississippi Valley eleven years, during which time he has had frequent attacks of "dumb chills." For four days prior to this illness, he had daily paroxysms of fever, which came on about noon. At 11 p. m., February 11, he had a hard chill which lasted for two hours. The fever rose quite high and was attended with vomiting. The patient became delirious, and from 2 a. m. to 9 a. m. he was comatose, his wife being unable to arouse him by calling, shaking or pinching. The coma was attended by stertorous breathing. I saw the patient about 10 a. m. He was complaining of headache and a slight burning sensation about his knees. Over each knee were large bullæ, extending from a few centimeters above the upper border of the patella to a few below the tubercle of the tibia. There were also two of much smaller size about the middle of the leg anteriorly. On the very black skin of this patient, these bullæ looked like burns of the second degree. The patient, in his delirium, had run across the room, taking his seat in a chair which sat before an open fire-place, in which were a few coals of fire. His wife states positively that he did not fall into the fire. His clothing was not burned. The temperature was 102 F., pulse 88, respiration about normal. The tongue was large and flabby, covered with a white coat; the mucous membranes were of fairly good color; the bowels were constipated, the spleen was enlarged, being palpable at the costal margin. The blood showed numerous estivo-autumnal parasites, all apparently of the same stage of development—about one-fourth the size of the red blood corpuscle. The urine did not show any albumin nor sugar.

Under vigorous antimalarial treatment for seventy-two hours, the fever subsided and did not return. The bullæ were pricked and a rather dark serum exuded. The places were dressed with gauze greased with vaselin. The patient was told to report if his legs gave him any trouble. I was called to see him again after about four weeks. I found gangrenous patches over areas referred to above, involving the skin and subcutaneous tissue. The ligamentum patella was exposed in the left leg. There was a gangrenous odor and the tissue was sloughing. The temperature was 99 F. The blood showed a few endoglobular, ring-form parasites and several estivo-autumnal gametes; red blood corpuscles, 4,028,000 per cm.; leucocytes, 6400. The differential count showed polymorphonuclears, 68.5 per cent.; small mononuclears, 23 per cent.; large mononuclears, 8 per cent.; eosinophiles, .5 per cent. Under the influence of quinin and antiseptic poultices, the gangrenous tissue sloughed away, leaving quite large ulcers which I show in the accompanying photographs.

The object in thus presenting this case in detail is to discuss some of the phases of malaria suggested by

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Practice of Medicine, and approved for publication by the Executive Committee: Drs. Frank Billings, George Dock and J. M. Anders.

it, which are exceedingly interesting to practitioners in malarial regions. The first phase which I desire to call attention to is

PERNICIOUSNESS IN MALARIA.

The clinical forms of pernicious malaria have been recognized since Torti's classic description of the periodic pernicious fevers, in 1743. These do not differ materially from the graphic description by Dr. Joseph Jones, of New Orleans, of these fevers occurring in the southern portion of the United States.

Marchiafava and Bignami state that "since the time when the parasites of the tertian and quartan were first known, there is no instance on record of a malignant infection caused by them, and no autopsy has ever been made in connection with a spring tertian or quartan." However, in this country, a few cases showing pernicious symptoms in infections with the benign parasites have been reported. They must, therefore, be regarded as exceptions to the general law that "the malignant infections are caused by the summer-autumn varieties of the malarial parasites." My experience has

sites. The first of these is their capacity for rapid propagation. This is seen in the great number of young forms which are seen in the blood a few hours after the beginning of the paroxysm. It is another characteristic of these parasites to disappear from the peripheral circulation to complete their cycle of development in other vascular areas.

The pathologic anatomy of these fevers teaches us that there is a decided tendency for the adult forms and forms of fission to accumulate in the capillary network of certain organs, e. g., the spleen, where the circulation is physiologically slow; in the brain, where the lumens of the vessels is small; and in the capillaries of the gastro-intestinal tract. Indeed, it has been proposed to explain the cerebral and choleriform types of these fevers by the mechanical obstruction thus produced in the respective areas. Frerichs thought this obstruction in the cerebral capillaries was produced by thromboses of malarial pigment. Laveran recognized the parasitic nature of these thromboses. Marchiafava and Bignami state: "Only exceptionally do we find in the

Estivo-autumnal Infection, with Multiple Spontaneous Gangrene.

Estivo-autumnal Infection, with Multiple Spontaneous Gangrene.

been that the large majority of cases of malarial fevers occurring in the Mississippi Valley are estivo-autumnal infections.

It is a well-known fact that in these infections the estivo-autumnal parasites are found in exceedingly large numbers. If this is not shown in the blood from the peripheral circulation during life, the autopsy usually shows them markedly increased in the viscera. It also appears that there are usually two predominating broods of parasites, sometimes more, in the blood of patients with pernicious fever. But Marchiafava and Bignami report cases in which the viscera, as well as the peripheral circulation, show a very small number of parasites. Dr. Ewing emphasizes this point in his report on the cases of malaria at Montauk Point. The cases of pernicious fever where only one brood of the parasites is present, as seems to be the condition in the case I report, are not infrequent. So it is evident that we must look for the cause of malignancy in another direction than merely in the large quantity of parasites.

These causes may be found in what has been termed the biologic characteristics of the estivo-autumnal para-

cerebral vessels free pigment (thromboses of pigment), or masses of melaniferous leucocytes (thromboses of phagocytes), or rarely also free adult parasitic forms or accumulations of free spores (parasitic thrombi in the proper sense of the term)." But they attribute the cerebral hyperemia to the accumulation, in the cerebral vessels, of red blood corpuscles loaded with amebæ, this stagnation being brought about by the degenerative changes wrought by the parasites in the red blood corpuscles, which impair their capacity for free circulation.

No doubt we should attribute great weight to the mechanical theory in explanation of these symptoms. Another factor, however, must be taken into account in these as in other forms of pernicious malaria, viz., the virulence of these parasites.

Many attempts have been made to demonstrate a malarial toxin, chief among which are the experiments of Professor Celli. The injection of 25 c. c. of serum, from a case of comatose malaria, into a little child, failed to show any toxic effect. The experiment to find the toxin shut up in the red blood corpuscles proved

equally futile. Still, almost every one now speaks of the toxin or toxins set free at the time of sporulation of the parasites. There is good evidence for believing that a malarial toxin exists, although it can not be demonstrated, so it is probably this property of the estivo-autumnal parasites more than any other which accounts for malignancy in malaria.

It is known that many uninfected corpuscles give up their hemoglobin. The presence of brassy corpuscles, the *ottonati*, shows the toxic effect of the parasite in its endoglobular phase.

Another proof of the great virulence of these parasites is the occurrence of hemoglobinuria in these infections. It has been stated by a distinguished authority that hemoglobinuria is not malaria, because he has been unable to observe the correspondence between the amount of hemoglobinuria and the number of parasites in malarial fever; and in his series there was no case in which hemoglobinuria was present with a large number of parasites, whereas in many cases with a large number of parasites no hemoglobinuria was present. As against this view it is interesting to note the findings of the Malaria Commission to the Royal Society. It tabulated a series of 28 cases, including 17 reported by Koch, 3 by Ollwig, and 5 studied by the members themselves, in British Central Africa; and show that of these, 75 per cent. showed malarial infection. It also points out that of 21 postmortems in cases of blackwater fever, 90 per cent. showed malarial infection. In these cases, we most infrequently are unable to find the parasite in the peripheral circulation during the attack. I have seen them in only two of nine cases in which the blood was studied. But the blood affords other evidence of malarial infection than the presence of parasites in it, viz., the presence of pigmented leucocytes, and the characteristic leucocyte variation of malarial infection. There were pigmented leucocytes in all the cases studied by me. This pigmentation was seen especially in the large mononuclears. It has been stated that there is a true leucocytosis in these cases, i. e., an absolute increase in the white cells and a relative increase of the polymorphonuclears. I do not believe this statement is correct. Numerous observers have noted that there is often a large increase of white cells in pernicious fevers. Dr. Thayer reports one in which there was an enormous increase, but the differential count showed the characteristic variation of malaria—an increase of the large mononuclears at the expense of the polymorphonuclears. The malaria commission appointed by the Royal Society has made more extensive investigation of the leucocytes in the estivo-autumnal infections than I have been able to find elsewhere. As a rule, a slight leucopenia was found. During the intermission or the remission, when marked, there was a relative increase of the large mononuclears; but during the paroxysm of fever, the polymorphonuclears were not only reduced, but sometimes increased. The large mononuclears were increased at the expense of the polymorphonuclears when the large forms of the parasites were in the blood, and the polymorphonuclears were not decreased when the small forms were present. I have been able to confirm these observations in several counts recently made. In all 5 cases studied by them they found an absolute increase in the white cells, and the leucocytic variation characteristic of malaria. I made differential counts in 2 of the 9 cases referred to by me. One showed P. M. N., 64 per cent.; large L. M., 20 per cent.; S. M., 16 per cent. The blood was obtained during the decline of the fever, near the first intermission. The same patient showed

P. M. N., 75 per cent.; L. M., 15 per cent.; S. M., 10 per cent. a few hours before death. Another case showed P. M. N., 64 per cent.; L. M., 19 per cent.; S. M., 16 per cent.; E., 1 per cent. So if pigmentation and leucocytic variations are evidences of malarial infection, where the parasites are few or absent in the peripheral circulation, hemoglobinuria must be malaria, and, being malaria, is a most striking evidence of the extreme virulence of the estivo-autumnal parasites.

Phenomena occurring in these infections, after the disappearance of the parasites, which have been termed post-malarial, are also evidence of the toxin theory. Kelch and Kiener, Mannaberg, and others, have noted the decrease in the percentage of hemoglobin after the disappearance of parasites. Dr. Ewing reports several cases of fatal anemia in his Montauk series, after the disappearance of the parasites. Dr. Thayer has reported a very interesting case of comatose fever, which he regards as post-malarial auto-intoxication. Marchiafava and Bignami state that, "in addition to the post-malarial fevers, there are individual cases of a post-malarial morbid condition, which must be put by the side of those which come after other infectious diseases (typhoid, pneumonia, scarlet fever, etc.)."

Another phase of malaria which the case reported suggests is

IMMUNITY

in malaria as it relates to the negro. We not infrequently see it stated that the negro is relatively immune from malarial infection, and sometimes that he is absolutely so. These statements are not in accord with my experience of eight years with the negro in the southern Mississippi Valley. Maurel, from his studies on this subject, concludes that no human race is immune from malaria, not even the black race.

On the plantation where I reside there were 24 whites last year. Of this number, 41 per cent. were infected with malaria. Of this number, however, 5 did not have malaria, and were away from home during the greater part of the malarial season. Excluding these, the malarial morbidity would be about 53 per cent. among the whites. There were 184 blacks; of these 61 per cent. were infected.

I have tabulated 20 cases of pernicious fever which I have seen since June, 1900, showing:

No.	Age.	Race.	Term of Residence Days.	Month.	Type.	Result.
1	34	White.	34	June. . . .	Hemoglobinuria	Recovery.
2	4	Black.	4	June. . . .	Cardialgic	Death.
3	7	Black.	7	July. . . .	Comatose.	Death.
4	60	Black.	20	July. . . .	Hemorrhagic, Melaena. . . .	Death.
5	15	Black.	2	July. . . .	Bilious continuous	Death.
6	3	Black.	3	August. . . .	Cerebral, convulsions	Recovery.
7	35	Black.	12	August. . . .	Hemoglobinuria	Recovery.
8	55	White.	55	August. . . .	Hemoglobinuria	Recovery.
9	4	Black.	3	August. . . .	Comatose.	Recovery.
10	11	Black.	5	August. . . .	Comatose.	Recovery.
11	5	Black.	5	August. . . .	Cerebral, convulsions	Recovery.
12	55	White.	55	August. . . .	Cerebral, delirium.	Recovery.
13	10	Black.	2	September. . . .	Comatose.	Recovery.
14	10	White.	1	September. . . .	Comatose.	Recovery.
15	18	White.	18	September. . . .	Hemoglobinuria	Recovery.
16	10	Black.	10	September. . . .	Comatose.	Recovery.
17	38	White.	10	September. . . .	Hemoglobinuria	Death.
18	30	White.	12	November. . . .	Hemoglobinuria	Death.
19	60	Black.	12	February. . . .	Comatose.	Recovery.
20	5	White.	60	February. . . .	Comatose.	Recovery.

Of this number it will be seen that 12, or 60 per cent., were negroes. The mortality among negroes was 33 1/3 per cent.; among whites, 25 per cent. Indeed, I have seen the comatose, convulsive, cardialgic, gastric, choleform, hemorrhagic and hemoglobinuric types occur in negroes, and in some death resulted.

The next phase which this case suggests is

GANGRENE IN MALARIA.

Several cases have been reported by Petit and Verneuil, in reviewing the occurrence of gangrene in connection with Raynaud's disease. These and others have been referred to by Barlow and Monroe. Osler reported a case of multiple spontaneous gangrene occurring in an estivo-autumnal infection, in the *Bulletin of the Johns Hopkins Hospital*, in February, 1900. He classifies three groups of gangrene: 1. In connection with Raynaud's disease. 2. Multiple spontaneous gangrene, in association with the acute infections, like measles, typhoid fever, typhus fever, scarlet fever, diphtheria, and malaria. 3. Multiple spontaneous gangrene of the limbs in young and middle-aged people without any obvious cause.

The case I report is one of multiple gangrene in an estivo-autumnal infection. I saw another case of spontaneous gangrene last year, which, from the clinical history, I regard as one in association with malaria. I regret very much that I could not get a blood-smear for examination, being without cover-glasses and slides at the time.

On March 27, 1900, the father applied to me for medicine for his child, C. S., a mulatto girl 4 years of age. He said that she had been having chills and fever for some time and was swollen about the face, hands, and feet. I prescribed iron, quinin and bitartrate of potash, under the influence of which, he said, the swelling disappeared. On May 16 I was called to see the child, and found her fairly well nourished. She had quite a high fever, with very rapid pulse and respiration. There was no edema and no eruption on the skin. The mucous membranes were anemic. The lungs were negative. There was no enlargement of the heart, and no cardiac murmurs. There was vomiting. The bowels were constipated; the spleen was enlarged, being palpable below the costal margin. The urine showed no albumin nor sugar. The child was put upon vigorous antimalarial treatment. On May 20, four days after my visit, the father reported that her fever had subsided and she seemed much better, but one foot was cold and apparently dead. The next day I visited the patient and found the right leg very cold as high as the knee, the toes in a state of dry gangrene, the dorsum of the foot black, with a few purpuric spots on the leg. The left leg appeared normal. There was no fever. A careful examination showed no endocarditis; there was no history of rheumatism. There were some pains about the knee on moving the leg. Another careful examination of the urine failed to show any albumin or sugar. I ordered a nourishing diet and tonic treatment, and advised the father to inform me when the line of demarcation formed. I did not hear any more from the patient until about two weeks afterward, when I happened to be passing the house, and asked how the child was getting along. The reply was, "Dat foot done come off." I went into the cabin and found the child sitting on a pallet on the floor, in very good spirits. On removing an old apron in which the leg was wrapped, I saw that spontaneous amputation had occurred at about the upper third of the leg.

BIBLIOGRAPHY.

1. On summer-autumn fevers: Marchiafava and Bignami, new Sydenham Society's Translation.
2. Celli: Malaria.
3. Thayer: Lectures on the Malarial Fevers.
4. Selected Monographs: Barlow's Translation of Maurice Raynaud's Observations on Spontaneous Gangrene; New Sydenham Society.
5. New York Medical Journal, February, 1899.
6. Bulletin of the Johns Hopkins Hospital, February, 1900.
7. Progressive Medicine, March, 1900.
8. Report of the Malaria Commission to the Royal Society.

DISCUSSION.

DR. E. H. MARTIN, Clarksdale, Miss.—I have not many criticisms to make, but I do wish to draw attention to a distinction between the forms of pernicious malaria spoken of; for

instance, hemoglobinuria and the comatose forms are all classed as pernicious malaria. Hemoglobinuria may be pernicious, but never in the sense we regard the other. Hemoglobinuria is malignant in that the germs present cause an absolute destruction of the blood cells. The other form is pernicious simply from an overdose of the toxin of malaria. In other words, one is pernicious from an overdose of the toxin and the other from the malignancy of the germs causing a destruction of the blood cells. In the negro we see but few cases of pernicious malaria in proportion to the total number of cases. In adults we see but few cases of pernicious malaria. In small children, especially in the small white children, we have a great many cases; it is quite an uncommon thing for a white family in the swamp lands of Mississippi to raise more than two out of five children, on account of the pernicious malaria, and the reason, I think, is evident; the blood of a child is very rich in the pabulum which nourishes the malarial germs; the vital resistance is low, and comparative immunity unattained. A vast number of the malarial germs develop, producing toxin, an overproduction of which kills the patient. I do not think that the mechanical theory can hold good; there is no reason why a few germs, and broken-down malarial organisms, should stop up the capillaries of the brain of a white child and not do the same for a white adult or negro child; yet it is a fact that these children have the "congestive" type almost exclusively. This is the idea that we get in speaking of pernicious malarial fever, that the toxins develop rapidly and in large quantity, and are administered to the patient in lethal doses.

Immunity in the negro is real, but only relative. It is a well-known fact that the number of white people that are affected with malarial troubles is less in proportion to the number of negroes affected than it was supposed to be; this is on account of the use by the whites of gauze doors, mosquito bars, the administration of quinin, etc. The class of "poor whites" who live as the negroes do are nearly all sufferers from acute or chronic malaria.

There is some hemoglobinuria among the negroes; but, where there were 400 negroes and 100 whites, there were 30 cases of hemoglobinuria among the whites and but 1 among the negroes. So, the negro has a relative immunity. Again, this relative immunity may be lessened by residence in non-malarial regions; a negro returning from a residence in New York, for instance, has lost it to a certain extent, but, after he has been in a malarial district for a time this relative immunity is regained. The negro is not fully immune, but is more nearly so than the white man; and the white man who has resided long in a malarious locality has a greater partial immunity than a newcomer.

DR. WILLIAM BRITT BURNS, Memphis, Tenn.—In referring to the immunity in the negro, the word relative, as mentioned by Dr. Martin, is a good one to employ. The negro is not immune to malaria, but he is relatively immune. The negro has malaria in mild form; it is true he has malaria in malignant forms, yet they respond to treatment quickly, much more so than the white people. I have seen several cases of comatose malaria in the black negro, and I have seen them get well; they all get well. I have never seen a case of comatose malaria in the negro terminate fatally, but I have seen many cases among the whites, especially in children. I have no doubt but what the accumulation of the malarial toxins in children is very rapid, and it is a common saying in the delta of the Mississippi valley, in Arkansas and Louisiana, that we must not allow a child to have a second or a third chill; it is the commonest thing to see, in a very large percentage of the cases, a child with congestion or congestive chill who dies. I have seen very few get well in infancy after the paroxysm of congestion came on. I have seen negroes, comatose for hours, get well.

DR. D. M. HALL, Memphis, Tenn.—I do not live in a malarial district as those who have preceded me, and therefore I have not seen as many cases of malaria, especially of the malignant type. My experience, however, leads me to believe that the negro is relatively immune. I have seen a great many more cases of malaria among white people than among negroes.

I think that if you take an equal number of whites and negroes, under similar conditions of exposure, etc., a greater number of whites will have malaria than the negroes. I do not know why this is so. We have in the South negroes living in places where the mosquitoes are in droves and where the conditions are good for the production of the disease, and they live there with very little infection; yet, I am sure the white people could not live there without being infected. In one city in the South there is a little creek which runs through the town and a great deal of sewage is thrown into it despite the efforts of the board of health. This is a great breeding-place for mosquitoes and they are there in swarms; near this creek seems to be a favorite place for the negroes; they live there on its banks, in cabins, and they seem to be in good health. Yet the white people who live near there are greatly infected with malaria. Therefore, I do not think the negro is immune, only relatively so. I think he can be overcome by malaria, but this depends upon the quantity of the poison introduced into the system. It requires more of the poison, in a more virulent form, to overcome the negroes than it does the white people.

DR. DELANCEY ROCHESTER, Buffalo, N. Y.—I presume it is well known that the mosquito is one of the sources of malaria; I should like to ask the gentlemen present if they recognize any other mode of entrance of the malarial parasite into the body besides the mosquito.

DR. W. G. HARRISON, Talladega, Ala.—The negro is relatively immune to most every disease, except lung troubles. I remember, when a student of medicine, that Dr. Westmoreland, in his lecture, stated that the negro was greatly immune to most of the bad and alarming features of tertiary syphilis. I have watched closely since, and I am sure that what he said was true. I think the negro has some relative immunity to all diseases except those of the lungs. They die from pneumonia and tuberculosis most readily. I saw a family in February where five died from pneumonia in one week; in the same community I had five or six cases among the white men and none died; all had the same kind of medical attention, the same physician, the same nursing, etc. Negroes die rapidly from pulmonary diseases of every kind.

Scientific medicine is being introduced into the Southern malarial swamps. The physicians in Louisiana, Alabama, Georgia, etc., are rapidly learning the use of certain apparatus for the better diagnosis and treatment of disease. It is now necessary that any man who practices in the South should equip himself with a microscope and all other instruments for clinical diagnosis. I have seen many cases, some of them among the negroes, who were taken sick with some inflammatory trouble and they seemed in a fair way to recover, when there was an unexplained rise in the temperature; the microscope revealed the malarial parasite; without it the rise in the temperature could not have been explained. Recently, within the last six weeks, a little girl of 2 years was brought to the hospital with cholera infantum; she got along slowly. Under the use of the saline solution she gradually got better, but, after one week, without any apparent reason at all, there occurred an elevation of temperature. I expected a return of the vomiting and watery passages, but they did not appear. After thirty-six hours an examination of the blood revealed some of the organisms of malaria. Under the proper administration of quinin the patient promptly got well. Regarding the statement that negroes are more susceptible to all diseases, I can only say my experience is very different. They easily contract smallpox, and most of them have latent syphilis, but I am sure the race possesses a distinct relative immunity against all the fevers, stomach and intestinal derangements, rheumatism, gout and many other maladies.

DR. VICTOR C. VAUGHAN, Ann Arbor, Mich.—I saw many cases of malaria in Cuba in 1898, and a large number of these were of the comatose variety. While I can not give any exact figures, I am under the impression that the comatose form of malaria, as we saw it in Santiago, was relatively quite as common among the negro soldiers as among the whites. Some time ago I went through the army records investigating this

matter, and while I can not give exact figures, I am confident that the general statement that I have just made is correct. The last speaker surprised me when he stated that the negro is immune to many diseases to which the white man is susceptible. The question of relative immunity between races is an important one, and I do not believe that definite conclusions should be stated until we are sure of the data on which they are founded. Only a few years ago it was believed that the negro was quite immune to yellow fever, but later epidemics have shown that this is not true. When yellow fever prevailed in our army in Santiago in 1898, two companies of the 24th Infantry, composed of negroes, were detailed as nurses on account of their supposed immunity to the disease. Practically all of these negroes became infected, and the mortality was quite as great as it was among the whites. It should be stated that this regiment had long been stationed in the western part of the United States, and possibly the negroes composing it had lost any racial immunity that they may have originally possessed. I have always been of the opinion that the white man is hardier and as a rule less susceptible to the infectious diseases than the negro, and I am surprised to hear it stated here that the negro is largely immune to most of the diseases to which we are susceptible. In this country tuberculosis and pneumonia are the most fatal of the infectious diseases, and certainly the negro is quite as susceptible to either of these as the white men, and the mortality is greater among the blacks. I am also convinced from extended observation that the negro succumbs to typhoid fever quite as readily as does the white man.

DR. G. W. GOINS, Breckenridge, Mo.—I wish simply to ask a question whether, in the practice of any present, many cases with a malarial symptomatology have occurred which simulated more or less typhoid fever and which did not yield to the administration of quinin.

DR. LOUIS SCHWARZ, Cincinnati, Ohio—Several of the gentlemen have spoken of a malarial imprint upon other diseases. This has been noticed in the progress of many of the diseases, particularly such that usually run an uninterrupted course. In the Millcreek valley, in which I live, many of the acute infections are observed to be disturbed in their behavior by malaria, and this belief is strengthened by the fact that quinin generally corrects the interrupted course. One reason why, in the treatment of the negro, better results are obtained by remedial agents, is perhaps because doses are more heroically given. This is particularly true where quinin is indicated for malarial affections. The statement that the negro enjoys an immunity from syphilis seems singular and can only be explained by invoking the belief that his immunity exists because he is born with the taint. The frequency with which unlooked-for fatal results occur in the negro is a matter of common observation on the part of those who have much to do with him. Sometimes, with no apparent reason for the fatality, death occurs, and it has been assumed by many that he loses his nerve just at the time he most needs it. It is known that the negro is very impressionable; this is best demonstrated by his conduct at religious exercises where in his excitement he is led to exhibit his fear in the most peculiar manner. It may be that a sense of impending danger lowers his resisting power and explains why he dies when there is no reason for it. Dr. W. S. Christopher, of Chicago, once applied the term "negroitis" to this manifestation.

DR. JOSEPH BRAYSHAW, Berlin, Ill.—In Central Illinois we have a few cases of malaria that so much resemble typhoid fever that a diagnosis can only be made by means of the microscope; but we have a large number of typhoid fever cases that so clearly resemble malaria that were it not for the Widal reaction and the absence of the malarial parasite a diagnosis would be impossible. We frequently have typhoid fever without pea-soup stools, without rose spots, and without very much abdominal tenderness, while we frequently have abdominal tenderness to a marked degree in malaria. The typhoid condition and typhoid tongue are frequent in our cases of malaria.

DR. T. B. FUTCHER, Baltimore, Md.—I should like to emphasize one important complication in malaria which Dr.

McElroy has already referred to in his paper, i. e., the occurrence of multiple gangrene in malaria. I was fortunate in having had an opportunity of seeing the case of gangrene of this nature which was under the care of Dr. Osler. The patient was a man, 23 years old, who was admitted to the surgical side of the Johns Hopkins Hospital, with dry gangrene over the dorsum of both feet, the heel of the right foot, and symmetrical patches over the fingers of both hands. There were, also, one or two large patches over the gluteal regions, and two small areas over the occiput. The case was at first a puzzling one from an etiological standpoint. There was slight elevation of temperature, and this, with the existence of an enlarged spleen, led to the examination of a specimen of fresh blood, which revealed numerous typical malarial crescents. This suggested the possible relationship between the malarial infection and the gangrene. Accordingly, the patient was transferred to the medical side and immediately placed upon anti-malarial treatment, quinin being administered in large doses. The patches of gangrene were treated locally. Improvement began at once, and the patient soon recovered from the local lesions. This was the first case of malarial gangrene that we have had at the Johns Hopkins Hospital out of a total of over 2000 cases, and it is undoubtedly an extremely rare complication. The case illustrated the importance, in cases in which the etiology is obscure, of a careful examination of the blood.

I should like to add that my impression in regard to the relative frequency of malaria in the two races in Baltimore, without having definite figures at hand, is that the negro race is practically as frequently affected as the white.

DR. W. D. HAINES, Cincinnati, Ohio—We have all seen *delirium tremens* follow in the wake of an accident, fracture, etc.; in the so-called moderate drinkers, on the other hand, we have seen malarial fever manifest its influence immediately after some surgical procedure. This would seemingly indicate that the prerequisite principle in either instance lay dormant until some exciting cause demanding extra attention on the part of Nature becomes operative. This is somewhat at variance with the usually accepted theory of disease, viz., that the host must prepare for the reception of the guest. We are exposed daily to the infectious principles of diphtheria, typhoid fever, tuberculosis, etc., and still we do not contract these maladies, forsooth the soil is sterile. On the other hand, how frequently we observe patients with lowered vitality or injury become an easy prey to these infections.

The statement made by one of the speakers, Dr. Harrison, astonished me not a little. He said: "The negro enjoys immunity from syphilis." It seems to me as I now recall personal experience in the treatment of this race that I have seen more cases of active syphilis in the negro than all other races combined. I never ask a negro *whether* he has had syphilis, but *when* he had it.

MEDICAL SHOCK.*

O. T. OSBORNE, M.A., M.D.
NEW HAVEN, CONN.

Before discussing what I interpret to be a condition of medical shock, it may be well to say a few words concerning the justification of ever rendering a certificate of death from "heart failure." Far be it from my purpose to excuse, justify, or encourage the use of this term to aid ignorant certifiers of undiagnosed deaths, or to give a mystic covering to illegitimate deaths, whether due to actual malpractice, or to the careless use of active drugs.

It may be said that the certificate of death from medical shock or heart failure is unjustifiable, because, if we seek carefully enough, we will find a cause for the failing heart, such as myocarditis, fatty degeneration, an

endarteritis or atheroma of the coronary arteries, etc., which causes, however, can many times be diagnosed only on the autopsy table.

Neurasthenia, which is a condition and not an entity, is recognized as a legitimate cause of death, but the pathology of this condition is more intangible than the condition of heart failure.

I do not wish to advocate the use of the term "medical shock" for conditions of failing heart, due to distinct pathologic conditions, or due to the gradual break-up or overpowering of the system by very acute or prolonged processes, but I think the term justifiable, when a badly acting and gradually weakening heart is the most urgent cause for anxiety, and its weakness is out of all proportion to the pathologic condition or symptoms present.

I believe it to be a fact that the hearts of our patients cause us more and more anxiety every year. If we doubt this statement, we have but to trace the treatment of acute conditions from the period of bleeding, calomel, tartar emetic, or other depletions, gradually down to the present time to find that first there was a cessation of depletions; next, the avoidance of nauseating doses of anything, denoting a greater care of the stomach; next, forced nutrition; then the general disuse of such drugs as aconite and veratrum viride; next alcoholic stimulation, and now we even note the loss of vasomotor pressure from alcohol and use strychnin in its place. This all means that the hearts of to-day do not withstand fevers or other systemic upsets as they formerly did.

To be specific, we are as justified in using the term "medical shock" as we are in using the term "surgical shock," and that the symptoms of heart failure which develop so unexpectedly in acute cases otherwise progressing satisfactorily, are due to this condition of shock. We all know the symptoms, viz., a more or less rapid heart, but with the least movement or excitement made very rapid; a soft, irregular, perhaps dicrotic or intermittent pulse; incomplete inspirations without much actual dyspnea, interspersed with frequent sighs, and the subjective symptom of precordial oppression. The temperature is generally low, the flesh cool and clammy, and if quick relief is not obtained, death will soon take place, caused by heart failure or medical shock.

The pathology of surgical shock is believed to be a paralysis of the vasomotor system, causing primarily a bleeding of the body into the large abdominal vessels, splanchnic paralysis, and secondarily great loss of heat from the dilated surface vessels.

I believe that medical shock is just as certainly a vasomotor paralysis, except, perhaps, in severe acute malignant affections where there may be a great destruction of red blood corpuscles. As a consequence of this paralysis, the major part of the blood is in the abdominal veins. The blood flows slowly into the dilated and, therefore, non-elastic arterioles; hence, slowly in the capillaries, and returning slowly in the veins, but imperfectly fills the heart cavities. The heart, not only from imperfect filling, but also from impaired nutrition, contracts irregularly, incompletely, and arrhythmically. The aorta has not the pressure ahead or the forcible quota of blood from behind, and consequently does not give enough elastic rebound to force blood properly into the coronary arteries, and the heart muscle is improperly nourished. By the lowered pressure, and the normal blood pressure is so vital, all the functions of the body begin to fail, all digestive processes are imperfect, and undoubtedly normal oxygenation is impaired, molecular death begins to take place, at least almost unaccountably

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Practice of Medicine, and approved for publication by the Executive Committee: Drs. Frank Billings, George Dock and J. M. Anders.