it, and neither surgeon nor patient are compromised; in short, it renders Wardrop's method more legitimate than formerly, because it does not involve the same risks.

Until I examined Mrs. L.—after death, I imagined her aneurism to be either on the innominate or on the subclavian at its origin. The error arose, I suppose, from the size of the tumour, its bulging towards the middle line, and from the shortness of the innominate trunk, which caused the subclavian to ascend more vertically than usual to get between the scaleni.

This variety is common in *dissecting-rooms*. However, it made no difference in the principles of treatment.

Edinburgh, July, 1858.

A Mirror

OF THE PRACTICE OF MEDICINE AND SURGERY

IN THE HOSPITALS OF LONDON.

Nulla est alia pro certo noscendi via, nisi quam plurimas et morborum et dissectionum historias, tam aliorum proprias, collectas habere et inter se comparare.—MORGAGNI. *De Sed. et Caus. Morb.* lib. 14. Proœmium.

ST. THOMAS'S HOSPITAL.

THREE CASES OF STONE IN CHILDREN; LITHOTOMY.

(Under the care of Mr. SOLLY and Mr. LE GROS CLARK.)

WE had the opportunity of witnessing four operations of lithotomy on the 24th instant, performed on young children, three at this hospital, and one at King's College Hospital by Mr. Fergusson. The two first were patients of Mr. Solly's, the subjects of stone for some months, their ages being under seven years. He experienced some difficulty in removing the stone from the first child, on account of its very much larger size than was anticipated, being about the dimensions of a good-sized walnut, and unusually large for so young a patient.

In the other child the stone was as large as a small marble, and was got out of the bladder with the aid of the scoop. Both of these patients had chloroform, and were in a complete state of anæsthesia, the first child taking a little time to recover himself, as there was the loss of some blood, which, with a free inhalation of the chloroform, for a minute or so caused a little anxiety, but the use of cold water to the face and slapping the cheeks brought back a natural condition.

The third patient, also a young child, was operated upon without chloroform by Mr. Le Gros Clark, and bore it very well. A stone was removed, the size of a large broad-bean, which was with difficulty detected before the operation. It was, however, distinctly felt by Mr. Clark, Mr. South, Mr. Solly, and Mr. Macmurdo, and the operation was performed with the more confidence.

A fourth case was present in the hospital under Mr. Macmurdo's care, but for some reason which we did not ascertain the removal of the stone was deferred till another occasion.

The patient at King's College Hospital was five years and a half old, and had been suffering from stone for the last eleven months. It was readily detected, and removed by Mr. Fergusson under the influence of chloroform. The stone was circular, the size of a marble, and studded with frosted crystals of, we presume, the oxalate of lime. As was stated at the Pathological Society at its last session, crystals on the surface of urinary calculi are somewhat rare, and when they do occur they are most generally the oxalate of lime.

We notice all these cases of stone thus early, because there is a reasonable probability that they will turn out well—a result that is almost general in children, unless circumstances may have occurred at the operation to induce conditions unfavourable to success. One that struck us in the first child was the large size of the stone, and the difficulty of catching hold of it and then extracting it, as some slight amount of laceration of the parts would necessarily ensue. We recollect very well an instance of a stone being removed so large that the perinæum was torn quite through into the rectum in its extraction, the

sphincter ani giving way with other intervening structures, and we believe the patient died.

Although the result of lithotomy in the young is so favourable, we think many surgeons will agree in the opinion that it is in reality a much more difficult operation than in the adult. The membranous portion of the urethra is very slender in the child, is easily torn, the prostate recedes from the finger on its being introduced, and sometimes when the incisions are being made the knife does not enter the bladder, and the stone will be searched for in vain. Such difficulties are not so often encountered in the adult.

Within the past few weeks the number of stone cases have been greater than usually met with. Besides the cases reported last week, Mr. Curling, Mr. Tatum, Mr. Birkett, and some others, have submitted patients to the usual operation with success, the subjects of stone being mostly children.

The case of Mr. Fergusson's, recorded in our "Mirror" of last week, in which five stones were removed, is doing well, and will make a good recovery, notwithstanding his advanced age.

GUY'S HOSPITAL.

DISEASE OF THE HIP-JOINT, WITH DISLOCATION OF THE HEAD OF THE THIGH-BONE; RESECTION; DEATH.

(Under the care of Mr. HILTON.)

WE have already freely entered into the consideration of the propriety of the operation of excision in disease of the hip-joint and pelvis, in previous "Mirrors," and will not, on the present occasion, again discuss it. As an addition to the two cases of excision recorded last week (to which we would refer the reader), we append another, performed in February last, which terminated unfavourably. The patient had been the subject of disease for nearly two years, and had undergone the various processes of inflammation, ulceration, suppuration, and, finally, dislocation of the thigh-bone on to the dorsum ilii. During his stay in the hospital, his sufferings were extreme, bed-sores formed, and he was not expected to live. Mr. Hilton, therefore, performed the operation as holding out a chance for life, and removed the head of the bone below the trochanter minor. Although the parts looked healthier than was anticipated, and the acetabulum unaffected, the poor man was so enfeebled by suppuration and the irritation from the bed-sores that he survived but eleven days. He seemed to improve after the operation, and had less pain; but we think a better chance of success would have been afforded had something been resorted to at an earlier period, before the system began to give way. That the termination was due to other causes than the operation itself, very few surgeons, we think, will be inclined to dispute.

The following notes of the case are reported by Mr. C. Hilton Fagge:—

W. W.—, aged fifty-four, was admitted on November 10th, 1857, with disease of his hip-joint. About a year and a half ago, after working as usual, he found at night that his right leg was numbed and painful, and the symptoms of diseased hip gradually developed themselves. His right leg is drawn up and wasted, and great pain follows pressure of the head of the thigh-bone against the acetabulum. On January 2nd, he says, he has not improved since his admission. On the 5th his back was very sore, and he had one bed-sore, for which he was placed on a spring bed.

Jan. 11th.—Is about the same. An incision was made to-day behind the great trochanter, and some serum, containing some cheesy matter, evacuated.

15th.—Does not seem much relieved by the discharge of pus from the incision made on the 12th.

18th.—No alteration in his condition. He refers all his pain, even when the limb is moved, to the knee; but there is not the slightest heat or inflammation there.

23rd.—The discharge is increased and very offensive. He does not suffer so much, and looks less anxious.

Feb. 1st.—The discharge continues in great quantity, but, with the exception of occasional attacks of pain, he is much more comfortable; there is also a slight blush of erysipelas over the hip. He is to take an emetic, and a purgative afterwards, and a mixture of aromatic spirits of ammonia and compound infusion of gentian, three times a day. A lotion of nitrate of silver to be applied over the part. The redness had nearly disappeared two days after.