Fig. 26. The germ of the 1st upper molar of *M. giganteus*. m'' , germ of 2nd dentition; m' , rudiment of 1st.

27. Premolars, canine, and abnormal incisors of *Perameles* (i_4 and i_5 fused).

28. Section through the dental lamina of *Perameles* between incisors 4 and 5 of the upper jaw, showing enlargement supposed to represent the true 5th incisor.

29. Palatal aspect of the premaxilla of a young *Petrogale xanthopus*, showing the three adult incisors in relation to the jaw and the vestigial canine. S , premaxillo-maxillary suture.

May 16, 1893.

OSBERT SALVIN, Esq., F.R.S., Vice-President, in the Chair.

The Secretary exhibited on behalf of Mr. Rowland Ward, F.Z.S., a complete skin of Grévy's Zebra (*Equus grevyi*), from a specimen recently shot by Col. Arthur Paget in Somali-land.

The following extracts from a letter, from Professor E. C. Stirling, M.D., C.M.Z.S., to Professor Newton, dated "Adelaide, April 10th, 1893"¹, were read:—

"... I am writing now to give you early information of a great 'find' of *Diprotodon* remains which I have recently been exploiting, and the results will clear up the remaining points of doubt about this great beast. Some months ago I received information that large bones were visible in a watercourse in a large dry salt lagoon called Lake Mulligan, situated some 20 or 30 miles to the north of a still larger dry salt lagoon which you will find marked as Lake Frome on any good map of South Australia. The accounts were so favourable that I sent up a good man to inspect and report. His accounts were also favourable, and I induced our Museum people to institute a systematic search—and we sent up four men with a complete outfit under the charge of a man who had some considerable experience of the kind of work. This party has been at work for nearly three months with very gratifying results from the reports I got. According to them there are *hundreds* of skeletons available and accessible, and many of them have already been excavated. Several nearly complete skeletons have been exhumed, and all the bones of the body have been found... It appears that the animal had five well-developed toes, the five terminal phalanges of which are almost equal in size. The carpal and metacarpal bones have also been found, though I am not specially informed whether the digits of the manus are also five in number. This, however, is implied from the enclosed sketch which I am able to send you.

"There is a tail of 9 vertebræ, which in one instance was 1 foot

¹ N.B.—Professor Stirling's telegram to Professor Newton published in 'The Times' of 25th April and 'Nature' of 27th April (vol. xlvii. p. 606) is of later date.

2½ inches long, and I am informed that the marsupial bones have also been found. We have also several distinct impressions of the skin of the fore foot.

"Several skeletons of a large Wombat about the size of a bullock have been also unearthed—probably *Phascolomys gigas*; and by the last advices the discovery has been made of a frail, tenderly-built animal about the size of a sheep, of which the nature is as yet unknown.

"The above account will give you some idea of the extent of this deposit of bones, which I think will undoubtedly prove to be one of the most important of its kind yet made in Australia. I am in hope that we shall be able to continue the search for some two or three months longer, by which time we shall have reaped a very rich harvest.

"I ought to mention that the locality is about 600 miles north of Adelaide, in a very inhospitable country, and 180 miles from the railway. In this season of drought all carting and travelling has to be done by Camels, and the temperature averages 110° F. in the shade, occasionally rising to over 120°. Myriads of flies and frequent sand-storms are other trifling difficulties which have to be undergone by the party.

"So far my engagements have not permitted me to get to the spot; but I hope to do so shortly. In any case it will be some time before we can get the bones down to town, as Camels are the only means of communication. When we do get the material to work upon we ought to have something interesting to say, but in the meantime I think you will be glad to have this preliminary report, however fragmentary . . . I send you herewith a rough sketch of one of the specimens *in situ* which will better explain some of the points I have mentioned, more especially with regard to the feet. Photographs of others have been taken, but I have not yet received copies; and indeed the impurity and scarcity of the water available has been so far a great hindrance to the development of the plates. The last advices I have received are nearly a month old, owing to the difficulties of access to the place, and I expect a good deal has been done since then.

"In writing to Sir W. Flower some weeks ago I was only able to mention that we had dropped into what seemed to be a good 'find,' but I was unable to give him any details. Perhaps, if you see him you will let him know how we are getting on.

" . . This is a very hurried letter, but I only received to-day my last advices about the *Diprotodon*, and the mail which I wish to catch goes to-morrow. Pray, therefore, excuse my imperfect sketch, which I only send as it is because I know you will be interested, and because I should like you to be the first to receive the information such as it is.

"*Addendum.* Professor Tate informs me that the geological formation of this salt-lake district of South Central Australia must be considered Pliocene. Lake Mulligan is, like Lake Eyre, Lake Frome, and other neighbouring lakes marked in the map, a vast

level expanse of salt-encrusted, black mud, only becoming filled after very heavy rains, which are not of very frequent occurrence. Lake Mulligan is relatively small, being only about 8 miles across, and the *Diprotodon* remains are somewhere about midway between the east and west edges. Usually the salt crust is not firm enough for bullock-traffic, and I may safely say that thousands of bullocks have at different times been bogged in crossing or attempting to cross . . . The bones, as will be seen, lie close to the surface, that is two or three feet [beneath] . . . It would appear that an immense herd of these and other animals had got bogged, probably in seeking water in a dry season, just as cattle do now in the North by hundreds. There is every indication that all this region of South Central Australia was formerly occupied by freshwater lakes. We have, for instance, remains of Alligators or Crocodiles from a district not far off, and other evidence of freshwater-life. Of course we are on the look-out for *Thylacoleo*; but, so far, it does not appear to have been met with; but I am quite hopeful that if we can manage to prosecute the search, that this and perhaps other remains will be met with.

"This is about all the information I can give you at present. Again excuse its unsystematic and fragmentary character, but I am writing far into the night so as to catch the mail to-morrow.

"E. C. S."

The following papers were read :—

1. On the Atrium and Prostate in the Oligochæta. By FRANK E. BEDDARD, M.A., F.R.S., Prosecutor to the Society.

[Received May 2, 1893.]

The structures termed "atrium" and "prostate" in the Oligochæta have given rise to considerable differences of opinion as to their nature.

The word "atrium" has been generally used for the dilated chamber into which the sperm-duct opens in the Tubificidæ and Lumbriculidæ; "prostate" has been the term usually applied to the glandular structures in Earthworms which are appended to the male duct. The latter term was first applied by Vaillant (1, who first described a worm with the glands in question, viz. *Perichæta posthuma*), who remarked concerning the gland: "faut il y voir une vésicule séminale accessoire ou une glande annexe comparable à une prostate?" The word "atrium" appears to have been first used by Claparède (2, p. 21) in the case of *Tubifex*. Vaillant was of opinion that the glands, which he described for the first time, in *Perichæta* were *sui generis* and not comparable to any structures in other Oligochæta. Vejdovsky is the first writer who has attempted any comparison between the structures in question

throughout the *Oligochæta* (3, p. 143). The following is a brief epitome of what he has said upon the subject: he considers that the glands described by Perrier in *Perichæta*, *Acanthodrilus*, and *Digaster* are possibly the equivalents of what he (Vejdovsky) terms the "Cement-Drüse" in the *Tubificidæ*; that the glandular tube in *Eudrilus* which Perrier called "*vesicula seminalis*" is to be looked upon as the homologue of the atrium in the aquatic *Oligochæta*; so also is the gland in *Pontodrilus*.

The next contribution to the subject is by myself; I pointed out (4) that the genus *Moniligaster*—an earthworm according to the definition of most naturalists—has a terminal gland connected with the sperm-ducts which agrees in all essentials with the atrium of the aquatic genera. In a further contribution (5) I dealt with the prostates of Earthworms in general, giving reasons for regarding them as the homologues of the atria of the lower *Oligochæta*.

Among Earthworms there are two principal forms of "prostate" met with. In *Acanthodrilus*, *Pontodrilus*, and other genera the glands are represented by long tubular bodies; in *Perichæta* &c. there are a pair of lobulate bodies often occupying the same position with regard to the ends of the sperm-ducts. One question to be decided was whether these two kinds of glands were related to each other; the next question was whether these glands were homologous with any structure in the lower *Oligochæta*. As to the first question, the tubular gland of *Acanthodrilus* was shown to differ only from the branched gland of *Perichæta* by the fact that the glandular cells of which it is largely composed are in the latter segregated into masses instead of forming a continuous coating. The answer to the second question is rendered easier by a consideration of the structure of the gland appended to the sperm-duct in *Eudrilus*. Perrier's account of the structure and relations of this gland were not, as I myself showed (6), quite accurate; the sperm-ducts open into the interior of the gland at about its middle. In this feature the gland of *Eudrilus* differs from that of *Acanthodrilus*, which is quite independent of the sperm-duct, or from that of *Pontodrilus*, where the sperm-duct only opens into the gland near to its external aperture. The identity of minute structure, however, appears to favour a comparison of the glands in *Eudrilus* and *Pontodrilus*; the only difference concerns the thick muscular coat of the gland in *Eudrilus*; but I pointed out that the genus *Trigaster* of Benham seems to be an intermediate form in this respect. "The identity of structure," I remarked, "between the glandular bodies appended to the termination of the vas deferens in *Eudrilus*, *Typhæus*, &c., leads to the inference that they are homologous; while the relations of the vas deferens to this body in *Eudrilus* clearly favours the supposition that it corresponds to the atrium in the '*Limicolæ*.'" The comparison of the gland of *Perichæta* to the prostates of the *Tubificidæ* seemed to me to be rendered impossible by reason of the fact that in the former the cells which it was sought to compare were covered by the peritoneum, while in the *Limicolæ* (I did not particularly mention *Tubifex*) and in the

genus *Moniligaster* "the prostates are formed by a metamorphosis of certain peritoneal cells." The real equivalent therefore of the "prostates" in the *Limicolæ* and in *Moniligaster* are to be found in the peritoneum of *Acanthodrilus*, *Perichæta*, &c. "In Earth-worms therefore," I concluded, "there are two organs which have been termed prostates—(1) the atrium of *Acanthodrilus*, *Perichæta*, &c.; (2) the atrium + prostate of *Moniligaster*."

These opinions were upheld with slight modifications in a subsequent paper (5, p. 117 &c.). One important difference between the atria of the higher and those of the lower Oligochæta I sought to explain by the primitive position of the atrial pores. Assuming that they originally were developed as invaginations of the clitellar region, it would follow that the lining membrane would consist, as does the clitellum, of two layers of cells; the resemblance of the cellular lining of the atrium in *Acanthodrilus* &c. to the clitellar epithelium has been commented upon by others as well as by myself. In the lower Oligochæta, on the other hand, the clitellar epithelium is one-layered; hence the lining membrane of the atrium is one-layered also. My later discovery, made since the paper to which I am now referring was written, that *Moniligaster* has a clitellum like that of the lower Oligochæta, still further confirms this way of looking at the facts. In the lower Oligochæta the atrial epithelium is ciliated—another difference as I then thought it; I have, however, lately found that in *Eudriloides brunneus* (n.sp.) the same ciliation at any rate partially occurs. I need not therefore recapitulate my attempt to explain what is now not a difference between the lower and higher genera of Oligochæta. Other points in this paper will be referred to again.

The views expressed in my papers were controverted by Benham (10). He points out in the first place a confusion of terms of which I was guilty. Mr. Benham writes:—"Beddard takes up a rather curious position in regard to the prostate of *Moniligaster*. For him the peritoneal coat, outside the muscular wall of the atrium, is the 'prostate' and is homologous with the "Cement-Drüse" (or prostate) of *Tubifex*. Now this prostate in *Tubifex* has been shown by Vejdovsky to be formed by a proliferation and outgrowth of the atrial epithelium at a certain point, which bursts through the muscular wall of the atrium and projects into the body-cavity. The atrial epithelium is derived from the epidermis, so that the 'Cement-Drüse' is epiblastic; whereas the glandular covering of the 'atrium' of *Moniligaster*, *Stylaria*, *Rhynchelmis*, &c. is mesoblastic,—it is in reality a modification of the peritoneal cells. Hence Beddard would regard the epiblastic 'prostate' (Cement-Drüse) of *Tubifex* as the homologue of the mesoblastic covering of the atrium in *Moniligaster*!" I did make this comparison at first; and it seemed to me to be justified by the curious fact that in *Tubifex* the Cement-Drüse was not covered by peritoneum, the rest of the atrium being covered; the disappearance of the peritoneum at this particular point appeared to me to indicate that possibly the data of Vejdovsky were not perfectly accurate.

In my fuller paper upon the subject I did not insist upon this comparison, which, however, I shall here again bring forward, though for different reasons.

Benham in his paper retains the name prostate for the glandular appendices of *Acanthodrilus*, *Perichæta*, &c., allowing, however, that the terminal part of these glands, which in *Perichæta* receive the sperm-duct, may be compared with the atrium of the lower *Oligochæta*. If this view is to be pushed to its logical conclusion, we are brought face to face with the somewhat puzzling conclusion that in *Eudrilus* half of the atrium is atrium and the rest prostate; that is to say, two parts of a continuous tube which are perfectly identical in structure are morphologically different.

A careful comparison of the various structures which have been called atrium by myself throughout the *Oligochæta* seems to me to prove their identity beyond the possibility of a doubt; an attempt to distinguish between the atria of different families would lead to the enunciation of somewhat impossible conclusions. Before comparing in some detail the atria of different forms, a few preliminary points may be disposed of. I have urged, as already mentioned, the similarity in structure between the atrium and the clitellum; this resemblance seems now to me to be without the significance which I have attempted to attach to it. The atrium is, as far as we know, an involuted region of epidermis; in many regions of the epidermis, not only in the clitellar region, there are unicellular gland-cells developed; the most prominent of these perhaps are the sometimes quite large glands which are associated with the genital papillæ in the genus *Perichæta*. I regard the glandular layer of the atrium as simply a thick layer of such unicellular glands; and the fact that we find every possible stage between a single mass of such gland-cells (*Tubifex*) attached to a portion of the atrium and a complete layer (*Acanthodrilus*) seems to justify this conclusion. The resemblance to the clitellum is then simply due to the fact that in both sections of the epidermis there are glandular cells present in great quantity.

In comparing the atria of the different families of the *Oligochæta* I shall commence with the higher groups. The tubular atria of the *Eudrilidæ* must be comparable to the somewhat similar atria of the *Acanthodrilidæ* &c. In *Pontodrilus* the sperm-ducts open into the atria at the junction of the muscular and glandular parts; in the *Eudrilidæ* into the glandular part itself. As, however, there are so many variations as to the exact place at which the sperm-duct does communicate with the atrium, this can hardly be regarded as a difference of more than trifling importance.

It will, I think, be generally allowed that the tubular glands of *Acanthodrilus*, *Pontodrilus*, &c., are homologous with the complicated lobate glands of *Perichæta*, *Cryptodrilus* &c. On *a priori* grounds only it seems impossible to distinguish glands which occur in such nearly related forms as *Cryptodrilus* and *Megascolides*. Perhaps if we had only to deal with *Acanthodrilus* and *Perichæta* the *a priori* grounds would not be so strong. Fortunately, how-

Spermiducal glands of various Oligochæta.

A. *Sutroa*. B. *Moniligaster*. C. *Pontodrilus*. D. *Acanthodrilus*.
E. *Eudrilus*. F. *Perichæta*. G. *Tubifex*.

v.d., Sperm-duct; *p.*, peritoneum; *m.*, muscular layer; *a.*, glandular layer;
b., lining epithelium; *sp.*, spermatozoa.

ever, there are stronger arguments than those to be derived from considerations of this kind. I have pointed out that in *Megascolex newcombei* the atria have the compact appearance of the corresponding organs of *Acanthodrilus*, to which also they show some slight approximation in outward form; this is further shown in the slight branching of the lumen of the gland. A better instance still has lately come under my observation. In the remarkable genus *Diplocardia*¹ of Garman (11) the atria appear on a dissection to resemble those of other Acanthodrilidæ; they are long tubular structures terminating in a short muscular duct. They have, however, a curious spotty appearance, small white rounded masses being imbedded in a darker ground-substance. They are also not quite so regularly tubular as they are in all other Acanthodrilidæ which I have dissected; towards the junction with the muscular duct the tubes are wider, and everywhere the margin is slightly crenate. In transverse sections the reason for these divergences from the normal Acanthodrilid structure was apparent: the white spots are aggregations of the pear-shaped glandular cells, which no longer form a regular investment of the lining epithelium of the tube; and here and there the lumen has inconspicuous diverticula into which the pear-shaped cells pour their secretion. The whole gland in fact shows a commencement of that branching which is so characteristic a feature of the atrium in the Perichætidæ. A little more development of the tendency to the aggregation of the pear-shaped cells and we should have an atrium exactly like that of the Perichætidæ.

If it be reasonable on *a priori* grounds alone to regard the glandular appendices of the sperm-ducts in the Megascolicidæ as homologous throughout, it is at least equally reasonable to seek for detailed resemblances in the atria in the different genera which make up the family of the Tubificidæ. At first sight there seem to be considerable differences. In *Tubifex* itself, and *Limnodrilus* and some other genera identical with *Tubifex* in these particulars, the atrium is long and has the well-known "Cement-Drüse" attached to it; this gland is limited in extent, though varying somewhat in different genera, and is superadded to the lining epithelium of the atrium; Vejdovsky (3) has shown that it is developed from the lining epithelium. In *Telmatodrilus* (12) there is a series of these outgrowths of the atrial epithelium, which deck the entire atrium, being symmetrically arranged. Finally, in my genus *Branchiura* (7) the atrium is closely invested by a thick layer of pear-shaped cells, which are separated from the lining epithelium by a muscular coat of circular fibres. *Ilyodrilus* appears to possess an atrium of the same kind but *minus* the muscular fibres. In *Tubifex* the muscular coat exists but it is less developed than in *Branchiura*; it is, moreover, deficient at the point where the "Cement-Drüse" is grafted on to the atrium. The only remaining genus of Tubificid which shows any differences from these

¹ I am indebted to the kindness of Dr. Beuham for the opportunity of examining this worm.

types is *Bothrioneuron* (13), where the "Cement-Drüse" is attached to the summit of a short diverticulum of the atrium.

It seems to me impossible to resist the conviction that *we have here a series of stages in the production of a limited "Cement-Drüse" out of a continuous glandular investment of the atrium.* In *Moniligaster*, whose atrium as regards the points under discussion is so like that of *Branchiura*, the glandular cells which invest it externally perforate the muscular layer to reach the inner epithelium; I imagine that this will be found to be the case with *Branchiura* too. A reduction of this investing layer to a restricted region of the atrium brings about the Cement-Drüse of *Tubifex*; a greater separation between the masses of glandular cells would produce the state of affairs characteristic of *Telmatodrilus*. It is noticeable that the absence of peritoneum upon the atrium of *Branchiura* is seen in *Tubifex* in the region of the Cement-Drüse; why this should be I cannot suggest; but that it is so is a further reason for the justice of my comparisons.

In the family Lumbriculidæ there are indications of a similar series of stages in the conversion of a complete glandular covering of the atrium into a more partial one; but the stages are fewer than in the Tubificidæ, in fact there are only, so far as our present knowledge of the group goes, two such stages. The more usual condition is seen in the majority; in *Rhynchelmis*, for example, the atrium is lined by a layer of cells, outside which are masses of pear-shaped cells which are aggregated to some extent into separate masses; the after all rather slight indications of discontinuity between the masses of cells which clothe the atrium in *Rhynchelmis* are still further emphasized in *Sutroa*. In this Annelid (8) the atrium is invested by very distinctly separate masses of glandular pear-shaped cells; the entire atrium is covered by a thin muscular layer which might be regarded as the peritoneal investment; between this supposed peritoneum and the glandular coat of the atrium lie masses of developing sperm; the structure therefore, whatever be its morphological nature, is, in function, a sperm-sac.

It seems to me that the two families Tubificidæ and Lumbriculidæ are nearly allied, in any case no one has disputed the morphological identity of the organs termed atria in both; it is therefore reasonable, at least pending further information, to assume that the atria correspond in detail, and in this case the Lumbriculidæ, like the Tubificid genus *Branchiura*, have an atrium which is frequently devoid of peritoneum. The intermediate condition between the atrium in the Lumbriculidæ and that in the higher Oligochæta is afforded by the Moniligastridæ. *Moniligaster* itself has an atrium which seems to be identical in all essentials with that of *Branchiura* and the Lumbriculidæ; it has two layers of cells which are separated by a well-developed muscular layer. I have shown (9) that the outer layer of cells sends processes through the muscular layer, which therefore are in a position to void their contents into the lumen of the atrium. I myself have only seen this arrangement

in one specimen of a species recently described by me (9) as *Moniligaster bahamensis*; in the other species investigated there was no trace of any perforation of the muscular layer by the outer layer of cells of the atrium: I imagine, however, that these prolongations of the cells were originally overlooked, owing to their inconspicuous character; they were especially obvious in the species *Moniligaster bahamensis* on account of the ducts being filled with granules of secretion, which were deeply stained by borax carmine. I compared this atrium with that of the *Lumbriculidæ* and various other aquatic *Oligochæta*, considering that the outer glandular layer of cells in both was the modified peritoneum. Benham, in the paper to which I have referred, agreed with me in this matter, but held that the cells, from the very fact that they were peritoneal, could not exert a glandular function; he doubted whether there was any known case of "coelomic epithelium being converted into a gland which pours its secretion to the exterior." This doubt appears to me to be quite justified, but then there is the undoubted fact that the processes of the cells in question *do* pour their secretion on to the exterior through the lumen of the atrium. The explanation of the difficulty seems to be that I was wrong in regarding this glandular epithelium as mesoblastic at all; I have never seen it myself, but Mr. Benham has shown me sections through the atrium of *Moniligaster* in which there was a distinct, though thin layer of peritoneum *outside* the glandular cells. The presence of this does not, however, invalidate my comparisons, though it puts them in a different light, for Benham remarked *apropos* of *Rhynchelmis* that he could "detect in sections a delicate membrane surrounding the pear-shaped cells which is no doubt the true coelomic epithelium."

The structure of the atrium in *Moniligaster* has also been studied by Rosa, who is disinclined to favour any "limicoline" affinities in that genus; he principally uses the characters of a new *Moniligastrid*, *Desmogaster* (14), to strengthen his position. In *Desmogaster* the atria have the following structure:—They are tubular glands slightly coiled in an S-shape; the lumen is much restricted, and is floored by a layer of cylindrical cells; outside this, the walls of the atrium are mainly muscular, the fibres being disposed centrally in a circular direction and longitudinally outside: among the external bundles of fibres are groups of gland-cells which by means of long ducts (formed merely by a prolongation of their own substance) reach the lumen of the atrium; but these groups of cells are not exposed at the surface of the atrium on account of the longitudinal muscular fibres; the entire organ is covered externally by a delicate layer of peritoneum. As Rosa points out, this seems to indicate a gradual moving outwards of the pear-shaped cells from their primitive position as lining the lumen of the atrium. My own criticism of this paper (as concerns the facts about the atria), published three years ago, need not here be referred to, as the data upon which it rested are not sufficient. I am now quite of opinion that Rosa was perfectly right in looking upon the

atria of *Desmogaster* as offering an intermediate stage between *Moniligaster* and the higher Oligochæta. In the Oligochæta we have an atrium consisting of two layers of epithelium, covered either by a simple thin peritoneum only, or (in the Eudrilidæ) consisting of thick muscular layers themselves covered with the peritoneal epithelium. In a few simplified genera of Megascolicidæ the lining of the atrium is reduced to the inner layer of epithelium only: these are *Ocnerodrilus*, *Gordiodrillus*, *Pygmæodrillus*, and *Kerria* (an Acanthodrilid). It is, I imagine, fairly probable that these forms are really, so far as concerns the atrium, degenerate in character; they show no real resemblance to the "Limicolæ" as I was at first inclined to believe; for as I now show that the same constituents are present in the atrium of the lower as of the higher Oligochæta, the single layer of cells is a difference from both.

At one time the family Geoscolicidæ was defined by having no atria like other families such as the Megascolicidæ; many genera, however, do possess organs connected with the external pores of the sperm-ducts which seem to be referable to this category. Rosa, however, in an account of the minute anatomy of these organs in *Kynotus* (15), is disposed to regard them as morphologically different from the atria of the Megascolicidæ. Unfortunately the description of the structure of the organs in question in *Kynotus* is unaccompanied by any illustrations, so that it is a little difficult to follow. The terminal male apparatus consists of a muscular bulb ("bursa propulsatoria") attached to the walls of the segment, on to which it opens by a retractor muscle; from this muscular sac arises a glandular tube of some length; the muscular sac is divided into two chambers by a transverse septum; from the dorsal one of these is given off the glandular appendix; the appendix is at first lined by a single layer of cells, and it is into this part of it that the sperm-duct opens; further back the lining epithelium is composed of two layers, the outer one being a mass of pyriform glandular cells. Rosa is of opinion that this structure is not the homologue of the atria of the Megascolicidæ; its resemblances, he thinks, are superficial and, such as they are, are due to convergence. The main reasons which Rosa gives for this opinion are three:—(1) The fact that these "pseudoprostates" are found only in a much specialized group of the Geoscolicidæ; (2) their difference in structure from the true "prostates" of the Megascolicidæ; (3) the great resemblance which they show to the sacs accompanying the copulatory setæ of the same worms. The second reason is further expanded into the actual points of difference; these are the following:—the presence of a retractor, the complicated structure of the terminal sac, the mode of connection of the sperm-ducts with the glandular appendix, and the disposition of the glandular part.

It appears to me that the differences are not so great as Rosa would have us believe; it is true that if we compare, as he suggests, *Pontodrilus* with *Kynotus*, the differences are more striking than the resemblances. I, however, decline to regard the

genera *Pontodrilus*, *Microscolex*, &c. as the lowest forms of the Cryptodrilidæ except in the sense that they are degenerate forms; it seems to me that it is the Eudrilidæ which present, as regards the atria, the most obvious likeness to the Geoscolicidæ. In them, as in the Geoscolicidæ, the atrium opens into a terminal "bulbus" which may even, as in *Eudrilus*, possess a pad developed out of its lining epithelium apparently not unlike the "scudo ovale" described by Rosa; the glandular appendix which in the Eudrilidæ opens into this is sometimes (as in *Heliodrilus*) hardly differentiated, as it is in many others, into a muscular and a glandular portion. In Eudrilids the position of the pore of the sperm-duct into the "appendix" is so variable that there is no difficulty in comparing the two families in respect of the fact that in both the sperm-duct opens into the glandular appendix. Even among the other subfamilies of the Megascolicidæ (I am here following Rosa's classification) there are not wanting indications of a close approximation between the atria and those of these Geoscolicids: in some species of the genus *Perichæta*, for example, the atria open into a sac variable in size, which itself opens on to the exterior; it appears to me that this sac is the equivalent of the terminal sac in the Geoscolicidæ. As to the presence of a retractor, which Rosa thinks distinguishes the atrium of the Geoscolicidæ, many Megascolicidæ have bands of muscles which appear to me to be perfectly comparable; for instance, in my genus *Octochaetus* (9) among the Acanthodrilids there are bands of muscles which, though perhaps not exactly inserted on to the atria, are attached to the body-wall in their immediate neighbourhood. A better example still is furnished by two species of *Eudriloides* lately examined by me; in these there are a series of muscular strands actually inserted on to the muscular termination of the two atria. To assert that these are in their nature different from those of the Geoscolicids seems to be too strong.

It seems therefore that the facts allow of no other view save that the various structures termed atrium by myself are homologous. The extremes are united by too complete a series of intermediade forms to permit of any doubt upon the point. There are, however, as has been pointed out, differences in detail between the atria of different groups; these amount to so much in the entire series that the complex gland of *Eudrilus*, or *Perichæta*, would perhaps unhesitatingly be regarded as different from the simple atrium of *Stylodrilus*, were there no intermediate stages. Though there would appear to be no great difficulty in deriving the atrium of one family from that of another, it is not so apparent which are the ancestral and which are the derived forms. To determine or attempt to determine this, opens up the whole question of the classification of these Annelids; and in my opinion a consideration of the facts relating to the atria confirms for the most part the scheme of classification adopted by myself. In order to determine which form of atrium is most primitive, it is necessary to enquire into the origin of the atrium.

Rosa's view appears to me the most probable; but this, however, he restricts to the Geoscolicidæ. In certain Geoscolicidæ, for example in *Microchaeta benhami*, there are a series of paired glands in the neighbourhood of the male pores; in some species, for example in *Kynotus michaelsonii*, setæ are associated with these glands which are in all respects similar to the penial setæ of other Earthworms. His view is that one pair of the glands have become converted into the "pseudoprostates" of other Geoscolicidæ, which I regard, as already pointed out, as identical with the atria of other Oligochæta. These glands are identical in structure with atria, and theoretical considerations also favour their homology. There are many instances among segmented animals of the reduction or concentration of metamerically repeated organs; we are more likely to be correct in assuming in such cases a reduction than a multiplication. It is easy on this assumption to understand the complete independence in the Acanthodrilidæ of the sperm-ducts and the atria; and, furthermore, the remains of additional glands such as occur in *Dichogaster damonis*; in this worm there are three pairs of tubular glands of which one only is connected with the sperm-duct. It is even possible that the glands (which I have called "capsulogenous") of the Perichætidæ are referable to the same category; and it may be pointed out that the papillæ upon which they open are sometimes regularly paired and correspond more or less accurately in position to the male pores. On this hypothesis of the original development of the atria out of copulatory glands, it is clear that those of the higher Oligochæta are nearer to the primitive form of these glands than the lower; and, furthermore, that the retention of more than one pair and their independence of the sperm-ducts are so far primitive characters. The Acanthodrilidæ come obviously very near to the base of the series, though a little further off than certain of the Geoscolicidæ. The Perichætidæ and the Cryptodrilidæ can readily be derived from the Acanthodrilidæ, and it must be remembered that some of the former have tubular atria and may therefore be nearer to the ground form. I do not propose in fact to discuss the relative positions of the different families, as I should have to take other characters into consideration with which this paper does not deal.

The question of terminology has now to be considered; it evidently stands in need of revision. Are we to retain the terms "atrium" and "prostate," or one or neither? They have been used in so many senses that it will lead to misunderstanding if they are again used without careful definition. The use of the term atrium has the objections that Mr. Benham has urged; for in *Acanthodrilus*, for example, the atrium is not a chamber leading into any other chamber or duct; on the other hand, to speak of the "atrium" of *Tubifex* as a "prostate" would give a wrong impression, for the idea of a prostate is a glandular appendix to the male ducts. To speak of the sac itself as an atrium and of the coating of pear-shaped cells as the prostate is disadvantageous; for, excepting in *Tubifex*, the two parts of the terminal apparatus

of the male ducts are combined into one structure. And in *Tubifex* their separation is more apparent than real. It is, in fact, far easier to object to current terms than to invent suitable new ones. In considering this question the origin of the so-called atria must be borne in mind; there are, as has been pointed out, strong reasons for believing them to have been produced by a reduction of glandular sacs often furnished with modified setæ found in the neighbourhood of the male pores and of the spermathecae; primarily, we are to suppose, they have no connection with the male pores; it would, however, be inconvenient to speak of them when limited to a single pair into which the sperm-ducts open as "copulatory glands," for by doing so their specialized condition would be lost sight of; it would be the same thing as calling the sperm-ducts nephridia. The various difficulties will perhaps be obviated if we speak of them in those cases where they have a clear relation to the sperm-ducts as "spermiducal glands." This term has the advantage that it has not been before used in the group and hence has no preconceived meaning attached to it; it is also to a considerable extent actually descriptive of the structures in question.

LIST OF LITERATURE REFERRED TO.

1. VAILLANT, L.—"Note sur l'Anatomie de deux Espèces du genre *Perichæta*, &c." Ann. Sci. Nat. (5) x. p. 225.
2. CLAPARÈDE, E.—"Recherches anatomiques sur les Oligochètes." Mém. Soc. Phys. Genève, t. xvi. 2me partie.
3. VEJDovsky, F.—System und Morphologie der Oligochaeten. Prag, 1884.
4. BEDDARD, F. E.—"Note on the Reproductive Organs of *Moniligaster*." Zool. Anz. 1887, p. 678.
5. BEDDARD, F. E.—"On the Structure of three new Species of Earthworms, with Remarks on certain Points in the Morphology of the Oligochæta." Quart. Journ. Micr. Sci. vol. xxix. p. 101.
6. BEDDARD, F. E.—"Contributions to the Anatomy of Earthworms: Nos. I., II., III." P. Z. S. 1887, p. 372.
7. BEDDARD, F. E.—"A new Branchiate Oligochæte, *Branchiura sowerbii*." Quart. Journ. Micr. Sci. vol. xxxiii. p. 1.
8. BEDDARD, F. E.—"A Contribution to the Anatomy of *Sutroa*." Trans. Roy. Soc. Edin. vol. xxxvii.
9. BEDDARD, F. E.—"On some new Species of Earthworms from various parts of the World." P. Z. S. 1892, p. 666.
10. BENHAM, W. B.—"Atrium" or "Prostate"? Zool. Anz. 1890, p. 368.
11. GARMAN, H.—"On the Anatomy and Histology of a new Earthworm (*Diplocardia communis*, gen. et sp. nov.)." Bull. Illinois State Lab. vol. iii. p. 47.
12. EISEN, G.—"Oligochætological Researches." Rep. Comm. Fisheries for 1888, p. 879 (1885).

13. STOLC, A.—“Monografie Ceských Tubificidu.” Abh. böhm. Ges. 1888.
14. ROSA, D.—“Viaggio di Leonardo Fea in Birmania e regioni vicine. XXV. Moniligastridi, Geoscoleци ed Eudrilidi.” Ann. Mus. Civ. Genova, (2 a) ix. p. 368.
15. ROSA, D.—“*Kynotus michaelsonii*, n. sp. (Contributo alla Morfologia dei Geoscolicidi).” Boll. Mus. Torino, vol. vii. No. 119.

2. Descriptions of Fifteen new Species of Shells of the Family *Pleurotomidæ*. By G. B. SOWERBY, F.L.S.

[Received April 28, 1893.]

(Plate XXXVIII.)

PLEUROTOMA WALCOTÆ, n. sp. (Plate XXXVIII. figs. 7, 8.)

Testa turrata, breviuscula, rugosa, pallida, antice zona pallide purpurea cincta; spira turrata, acuta, gradata, sutura inconspicua; anfractus 8, ubique spiraliter striati, supra medium angulati, infra angulum convexi, longitudinaliter costati (costis in anfr. ult. circ. 12), spiraliter lirati (liris in anfr. penult. 4); anfractus ultimus spiram subæquans, infra angulum leviter convexus, basin versus subcontractus, vix rostratus; apertura latiuscula, fauce purpureo-grisea; canali brevissimo, leviter recurvo; columella rectiuscula; labrum acutum, crenulatum, superne sinu profundo latiusculo emarginatum.

Long. 20, *maj. diam.* 10 millim.

Hab. Spencer's Gulf, South Australia.

The shell is of the typical form of the section *Drillia*. The purple-grey zone on the lower part of the body-whorl seems characteristic. The type specimen, belonging to the South Australian Museum, has been kindly lent me for description and figuring by W. T. Bednall, Esq.

PLEUROTOMA HUBERTI, n. sp. (Plate XXXVIII. figs. 11, 12.)

Testa turrata, acuminata, rugosa, fusca, albo-costata, zona albida cincta; spira elata, acuta; sutura vix impressa; anfractus 10, convexi, obtusissime angulati, ubique spiraliter striati, longitudinaliter costati (costis in anfr. ult. circ. 12), infra angulum spiraliter lirati; anfractus ultimus spiram paulo brevior, zona albida cinctus, infra medium contractus, breviter rostratus; apertura latiuscula; canali breviusculo, leviter recurvo; labrum incrassatum, superne angulatum, ad angulum late et subprofunde sinuatum.

Long. 16, *maj. diam.* 8 millim.

Hab. Cape Grenville, N.E. Australia.

Of the typical form of the section *Drillia*, the sinus forming a

broad channel in the wing-like expansion of the lip. The longitudinal ribs are nearly white, between which the brown colouring is partly disposed in transverse linear streaks.

PLEUROTOMA ALBOVARICOSA, n. sp. (Plate XXXVIII. figs. 9, 10.)

Testa elongato-fusiformis, glabra, fusca, albo interruptim strigata; spira elongata, turrata, acutissima; anfractus 8, convexi, longitudinaliter costati, costis rotundatis, albidis, zona fusca transmeatis; sutura callosa, albida; anfractus ultimus elongatus, costis 6 munitus, fusco bifasciatus, infra medium constrictus, spiraliter striatus, longirostratus; apertura elongata; columella fere recta; labrum tenue, superne mediocriter sinuatum, canali longiusculo.

Long. 22, *maj. diam.* 6 millim.

Hab. East Africa? (ex coll. Keen).

A prettily painted shell, with a smooth shining surface. The ribs are whitish, crossed on the upper whorls with one and on the body-whorl with two brown zones; the interstices between the ribs are entirely brown. The suture is covered with a whitish callosity.

It may be doubted whether this species is rightly placed in the section or subgenus *Surcula*, but to judge only from the shell, and to propose a new section for it, would be only to add to the already far too numerous ill-defined subdivisions.

PLEUROTOMA EDITHÆ, n. sp. (Plate XXXVIII. figs. 3, 4.)

Testa subfusiformis, sordide albida, fusco zonata; spira elongata, acuta; anfractus 10, lævigati, superne leviter concavi, obtuse angulati, infra angulum costati, costis elevatis, obtusis, latiusculis; sutura callosa; anfractus ultimus zona lata fusca cinctus, supra medium obtuse angulatus, supra angulum concavus, infra leviter convexus, basin versus attenuatus, spiraliter obscure liratus; apertura oblonga, fauce fusco-tincta; labrum acutum, superne angulatum, ad angulum late et profunde sinuatum; columella arcuata, supra callosa, canali brevissimo.

Long. 18, *maj. diam.* 7 millim.

Hab. Hong Kong.

A shell of simple character. I place it provisionally, but with some doubt, in the section *Clavus*, which may or may not prove to be clearly separable from *Drillia*.

PLEUROTOMA INCLINATA, n. sp. (Plates XXXVIII. figs. 25, 26.)

Testa subfusiformis, solidiuscula, luteo-albida fusco-strigata et balteata; spira elongata turrata, acuta, leviter inclinata; anfractus 10, convexiusculi, costis numerosis acutiusculis leviter obliquis muniti, spiraliter minutissime striati; anfractus ultimus spiram vix æquans, superne leviter concavus deinde convexus, prope marginem costa crassa instructus, ad basin attenuatus

haud rostratus, spiraliter liratus; apertura oblonga, latiuscula; columella callosa; labrum acutum, late et profunde sinuatum.

Long. 20, *maj. diam.* 8 millim.

Hab. Mauritius.

This species belongs to the *Clavus* group. It has been placed in various collections as *P. albicostata*, Sowerby, with which I find, however, it has little affinity. It comes rather nearer to *P. obliquata*, Reeve.

PLEUROTOMA KEENI, n. sp. (Plate XXXVIII. figs. 13, 14.)

Testa turrita, solida, pallide fusca, nigro-fusco strigata, albo granulata; spira acuminata (apice?); anfractus obtuse angulati, costis numerosis confertis, longitudinalibus subobliquis biseriatim granulatis muniti; interstitiis nigro-fusco strigatis; anfractus ultimus supra angulum leviter concavus, infra angulum convexiusculus, infra medium leviter attenuatus, haud rostratus; apertura oblonga; columella callosa, alba, polita; labrum fere rectum, supra angulum mediocriter sinuatum.

Long. 24, *maj. diam.* 5 millim.

Hab. — ? (ex coll. Keen).

A species of somewhat remarkable character, apparently belonging to the section *Crassispira*. The ribs are beaded with white granules and bordered with dark brown streaks. The specimen has lost several whorls of the spire.

PLEUROTOMA HUNGERFORDI, n. sp. (Plate XXXVIII. figs. 1, 2.)

Testa anguste turrita, grisea, ad apicem fusca; spira elongata, acuta; anfractus 8, læves, supra medium leviter concavi, infra costis paucis lævigatis muniti; anfractus ultimus breviusculus, supra medium obtuse angulatus, inferne levissime attenuatus, haud rostratus; apertura oblonga, latiuscula; labrum acutum, sinus profundiusculus, mediocriter latus.

Long. 12½, *maj. diam.* 4 millim.

Hab. Hong Kong.

A very simple shell, without sculpture, and with but few smooth longitudinal ribs. It is hardly possible to say in which section this species should be placed, whether it is a *Drillia* or a modification of *Clavus*.

PLEUROTOMA INFRAFUSCA, n. sp. (Plate XXXVIII. figs. 5, 6.)

Testa acuminata, albida, antice rufo-fusca; spira turrita, acutissima; anfractus 10, primi 2-3 læves, sequentes angulati, supra angulum concavi, infra vix convexi, costis numerosis rotundatis, ad angulum elevatis muniti, maculis minutis fuscis ad angulum picti; anfr. ultimus spiram subæqualiter longus, multo latior, basin versus leviter attenuatus, haud rostratus; apertura latiuscula; labrum acutum, vix sinuatum, canali brevissimo.

Long. 19, *maj. diam.* 7 millim.

Hab. Mauritius ? (ex coll. Sir D. Barclay).

The most striking character of this species is its bright reddish-

brown base. It will probably come into the section *Clavus*. The specimen is now in the collection of General Tripe.

DAPHNELLA FUSCO-PICTA, n. sp. (Plate XXXVIII. figs. 15, 16.)

Testa angusta, fusiformis, albida, nigro-fusco conspicue flammulata; spira elongata, ad apicem acuta; anfractus 9-10, convexi, undique eximie cancellati; sutura angustissime canaliculata; anfractus ultimus oblongus, spiram leviter superans, vix inflatus, inferne leviter attenuatus, haud rostratus, flammulis longitudinalibus albo bifasciatum interruptis pictus; apertura oblonga, utrinque attenuata; labrum leviter crenulatum, sinu profundiusculo emarginato, canali brevissimo.

Long. 17, maj. diam. 5½; apertura longa 7½, lata 2 millim.

Hab. Hong Kong.

A prettily marked and delicately sculptured shell, minutely cancellated throughout. It somewhat resembles *D. flammea* (Hinds), but differs considerably from that species in sculpture (fine close-set longitudinal striæ crossing the spiral ones). In sculpture it closely resembles *D. diluta* (Gould), which species is very different from it in form. Another allied species is *P. delicata* (Reeve), from which it differs both in form and sculpture.

DAPHNELLA SPENCERÆ, n. sp. (Plate XXXVIII. figs. 17, 18.)

Testa acuminato-fusiformis, albida, fusco maculata; spira acuminata, acuta; anfractus 8, convexiusculi, supremi longitudinaliter plicati, undique spiraliter striati; maculis fuscis conspicuis ornati; anfractus ultimus spiram superans, leviter inflatus, maculis fuscis regularibus rotundatis biseriatim dispositis pictus; apertura oblonga; labrum arcuatum, acutum, sinu profundiusculo angustiusculo emarginato, canali brevissimo.

Long. 10, maj. diam. 4; apertura longa 4, lata 2 millim.

Hab. Hong Kong.

A beautiful little shell, of which I have only seen a single specimen. It is finely spirally striated throughout, the upper whorls being longitudinally plicated. The colouring consists of conspicuous rounded brown blotches, which upon the body-whorl are disposed in two rows, one above and one below the middle of the whorl.

DAPHNELLA ELATA, n. sp. (Plate XXXVIII. figs. 19, 20.)

Testa turrata, luteo-albida, maculis pallide fuscis promiscue conspersa; spira perelongata, acutissima; anfractus 10, primi 2 fusi læves, sequentes leviter angulati, liris spiralibus numerosis acutis cingulati, longitudinaliter minute interplicati; anfractus ultimus spiram vix æquans, leviter convexus, infra medium constrictus; apertura oblonga, parva; labrum extus crenulatum, sinu breviusculo emarginato, canali brevissimo.

Long. 15, maj. diam. 4; apertura longa 5, lat. 1½ millim.

Hab. Mauritius.

This species has a much longer spire than others of the group;

in sculpture it resembles *D. lactea*, but the spiral ridges are closer; in form it approaches *D. flammea* and *D. subula*.

DEFRANCIA INFRACINCTA, n. sp. (Plate XXXVIII. figs. 21, 22.)

Testa oblonga, solida, carnea; spira mediocriter elata, ad apicem acuta; anfractus 8, convexi, linea fusca tenuissima ornati, spiraliter lirati, longitudinaliter costati, costis numerosis, rotundatis; anfractus ultimus vix inflatus, infra medium leviter constrictus, fascia fusca conspicua pictus; apertura breviuscula; labrum extus incrassatum, intus serratum, superne profunde sinuatum, canali brevissimo.

Long. 10, maj. diam. 4 millim.

Hab. Mauritius.

A rather rough shell, with numerous rounded ribs, crossed by close-set raised ridges. The conspicuous band near the base surmounted by a thin brown line seems characteristic.

DEFRANCIA MAURITIANA, n. sp. (Plate XXXVIII. figs. 23, 24.)

Testa turrata, albida, maculis fuscis promiscuis picta, undique costis numerosis confertis granulatis sculpta; spira elata, leviter convexa, ad apicem acuta; anfractus 9, gradati, levissime convexi; sutura impressa; anfractus ultimus spiram vix æquans, haud inflatus, rectiusculus, ad basin constrictus; apertura parviuscula, oblonga; labrum crenulatum, incrassatum, superne sinu profundo emarginato, inferne abrupte constrictum, canali brevissimo.

Long. 10, maj. diam. 3 millim.

Hab. Mauritius.

A conspicuously painted species, with a strongly granulated surface; the constriction of the base is peculiar.

CYTHARA GUENTHERI, n. sp. (Plate XXXVIII. figs. 27, 28.)

Testa solidiuscula, utrinque acuminata, albida, pallide fusco tincta et vittata, postice purpureo tincta; spira elato-conica, gradata; anfractus 10, angulati, undique spiraliter striati, longitudinaliter costati; anfractus ultimus spiram superans, convexiusculus, basin versus attenuatus; apertura elongata, angusta; columella levissime curvata, albo-callosa; labrum extus incrassatum, ad marginem acutum.

Long. 25, maj. diam. 11; apertura longa 15, lat. 3 millim.

Hab. Holothuria Banks, N.W. Australia, 38 fathoms.

A beautiful delicately sculptured shell, hitherto without name, in the British Museum Collection.

CYTHARA RINGENS, n. sp. (Plate XXXVIII. figs. 29, 30.)

Testa abbreviata, angulata, utrinque acuminata, alba; spira conica, ad apicem acuta; anfractus 7, angulati, longitudinaliter costati, spiraliter subtilissime striati; sutura anguste canaliculata; anfractus ultimus spiram superans, angulatus, supra angulum leviter concavus, deinde vix convexus, basin versus

regulariter attenuatus; costis longitudinalibus circ. 8; apertura elongata, angusta, columella granulata et plicata; labrum incrassatum, intus serratum, supra angulum leviter sinuatum.

Long. 7, maj. diam. 3 millim.

Hab. Hong Kong.

Of the typical form of *Cythara*. The serrated lip and columella present the appearance of two rows of teeth, almost like those of the *Cypree*.

EXPLANATION OF PLATE XXXVIII.

- Figs. 1, 2. *Pleurotoma hungerfordi*, p. 489.
 3, 4. — *edithæ*, p. 488.
 5, 6. — *infrafusca*, p. 489.
 7, 8. — *walcotæ*, p. 487.
 9, 10. — *albovaricosa*, p. 488.
 11, 12. — *huberti*, p. 487.
 13, 14. — *keeni*, p. 489.
 15, 16. *Daphnella fusco-picta*, p. 490.
 17, 18. — *spenceræ*, p. 490.
 19, 20. — *elata*, p. 490.
 21, 22. *Defrancia infracincta*, p. 491.
 23, 24. — *mauritiana*, p. 491.
 25, 26. *Pleurotoma inclinata*, p. 488.
 27, 28. *Cythara guentheri*, p. 491.
 29, 30. — *ringens*, p. 491.

3. A Nominal List of the Mammals inhabiting the Bornean Group of Islands. By A. H. EVERETT, C.M.Z.S.

[Received May 2, 1893.]

As no complete catalogue of the Mammals of Borneo and Palawan appears to be available for the use of Oriental Zoogeographers, whilst a considerable number of new species have been recorded of recent years, the notices of which are scattered among various scientific serials, it is hoped that the present list, although a merely nominal one, may not be without its use.

I have elsewhere¹ detailed the evidence which seems to render it necessary that Palawan should be treated as forming an integral part zoologically of the Bornean group of islands. The area of the Bornean group as a whole may be defined, therefore, for the purpose of the present list, by the same limits as those which I have adopted in a list of the Birds of Borneo and Palawan², viz., by a line which starts from a point immediately to the west of St. Julian I. in the Tambelan Archipelago, and, being drawn south of the Great Natuna (Bungoran I.), passes northward of Labuan and thence follows the 100-fathom line so as to embrace Balabac, Palawan (Paragua), the Calamianes, and the Cuyo Islands, and, returning along the same line of soundings on the southern side of

¹ P. Z. S. 1889, p. 220.

² Journ. Straits Branch R. As. Society, xx, p. 92 (1889).

Palawan, is drawn immediately to the eastward of the islands of Cagayan Sulu and Sibutu,—whence it is continued through the Macassar Straits south of the Paternoster, Lauriot (Laset Ketjil), and Solombo islets, and in a north-westerly direction through the Karimata Strait back to the island of St. Julian.

In attempting thus to define the area within which the neighbouring smaller islands may be regarded as being affiliated zoologically to Borneo, there exists no guide in many cases beyond their greater proximity to the latter island than to the other large masses of land by which they are more or less closely environed, supplemented by the evidence (often very imperfect) afforded by the soundings shown on our Admiralty charts. The limits adopted above must therefore be looked upon as being in some degree approximate. Nevertheless, as there are not any islands of considerable size of which the fauna is wholly unknown bordering upon the line of delimitation, it is not probable that material extensions or contractions of the boundary here assumed for the Bornean group as a whole will be necessitated by the result of future field-work.

Turning to the accompanying list, it may be remarked that no family of Mammals is peculiar to the group and that only three genera are so, viz. *Nasalis*, *Trichys*, and *Rhithrosciurus*, which, so far as we know at present, are all confined exclusively to the island of Borneo itself. Peculiar species, however, are fairly numerous, and they are indicated in the list by the prefix of a single or double asterisk in accordance with their being confined to Borneo or to the Palawan Sub-group respectively. Although the general facies of the mammalian fauna of the Sub-group is clearly Bornean, it is to be noted that no species appears to be peculiar to the group as a whole, a fact which suggests the inference that, closely connected as Borneo has undoubtedly been with Balabac and Palawan, and isolated as they have been together from the mainland of Asia, there has also been much isolation of Borneo and Palawan *inter se*.

A few of the species enumerated have been no doubt introduced by man. Such are the Common Mouse and the Black and Brown Rat, with the Buffalo; and it seems probable that the Javan Sambur (*Cervus hippelaphus*) and the Elephant owe their presence to the same agency. A few other species which have been credited to Borneo will be found to have been omitted below. Thus *Hylobates concolor*, Harlan, is almost certainly identical with *H. muelleri*, Martin. *Macacus maurus*, F. Cuv., long believed to inhabit Borneo, has been shown¹ to be restricted to the southern peninsula of Celebes and to the neighbouring island of Bouton. There is a stuffed specimen of *Canis aureus*, Linn., in the Leyden Museum which is labelled as having been collected by Diard in Borneo, but it is more than doubtful whether the locality is correct². The Javan Rhinoceros (*R. sondaicus*) has been supposed to exist in Borneo, but my inquiries on the subject have failed to elicit any reliable evidence

¹ Weber, Zool. Ergebn. Niederl. Ost-Ind., Leiden, 1890, p. 103.

² Blanford, Faun. Brit. India, Mammalia, i. p. 141.

that this is the case; but the discovery of some subfossil molars in Sarawak, which have been identified as belonging to this species on good authority¹, render it probable that it may yet be discovered in the comparatively unexplored interior. It is remarkable, in this connexion, how extremely rare the remains of Pleistocene or Quaternary mammals appear to be in Borneo, and this scarcity must be real, taking into account the extensive surface-mining which has long been practised in Western and South-eastern Borneo, and the value which natives set on such objects as fossil teeth for use as charms, medicine, &c.

Notwithstanding the numerous additions made in recent years to the list of the Mammals of Borneo and Palawan by Mr. John Whitehead and of Borneo by Mr. Charles Hose, many species doubtless await discovery, especially among the smaller mammals and the Cetacea, of which latter our knowledge remains at present practically a blank. Among the Chiroptera particularly, it may be expected that a number of species already known in the neighbouring countries will be detected as ranging into some part of the Bornean group, such, for instance, as *Miniopterus schreibersi*, *Kerivoula picta*, *Vespertilio hasselti*, *Vesperugo abramus*, *V. noctula*, *V. pachypus*, *Hipposiderus speoris*, *Xantharpyia amplexicaudata*, &c.

In concluding these brief remarks I wish to express my best thanks to Mr. Oldfield Thomas, of the British Museum, to whose courtesy and assistance I have been frequently indebted both in the identification of species and in various other ways.

PRIMATES.

Fam. SIMIIDÆ.

- Simia satyrus*, Linn.
- *morio*, Owen².
- Hylobates leuciscus*, Schreb.
- * — *muelleri*, Martin.

Fam. CERCOPITHECIDÆ.

- Semnopithecus maurus*, Schreb.
- * — *chrysomelas*, Müll.³
- * — *cruciger*, Thos.³
- * — *hosei*, Thos.
- * — *everetti*, Thos.
- * — *rubicundus*, Müll.
- * — *frontatus*, Müll.
- * *Nasalis larvatus*, Wurm.
- Macacus arctoides*, Geoffr.⁴
- *nemestrinus*, Linn.
- *cynomolgus*, Linn.

Fam. LEMURIDÆ.

- Nycticebus tardigradus*, Linn.

Fam. TARSIIDÆ.

- Tarsius spectrum*, Pallas.

CHIROPTERA.

Fam. PTEROPODIDÆ.

- Pteropus edulis*, Geoffr.
- *hypomelanus*, Temm.
- * *Cynopterus spadiceus*, Thos.
- *marginatus*, Geoffr.
- * — *lucasii*, Dobson.
- *ceadatus*, Temm.
- * — *maculatus*, Thos.
- Eonycteris spelæa*, Dobson.
- Carponycteris minimus*, Geoffr.

Fam. RHINOLOPHIDÆ.

- Rhinolophus luctus*, Temm.
- *trifolius*, Temm.
- *affinis*, Horsf.
- *minor*, Horsf.
- Hipposiderus diadema*, Geoff.
- *galerita*, Cantor.
- *cervinus*, Gould.
- * — *doriæ*, Peters.
- *bicolor*, Temm.

Fam. NYCTERIDÆ.

- Megaderma spasma*, Linn.

¹ Busk, P. Z. S. 1869, p. 409.

² Doubtfully distinct from *S. satyrus*. Said to occur in Northern Sumatra.

³ Both doubtfully distinct from *S. femoralis*, Horsf.

⁴ Günther, P. Z. S. 1876, p. 425.

Fam. VESPERTILIONIDÆ.

- * *Vesperugo stenopterus*, Dobson.
- *imbricatus*, Horsf.
- *tenuis*, Temm.
- * — *tylopus*, Dobson.
- * — *doriae*, Peters.
- Scotophilus kuhli*, Leach.
- Harpiocephalus suillus*, Temm.
- Vespertilio adversus*, Horsf.
- *muricola*, Hodgson.
- Kerivoula papillosa*, Temm.
- *hardwickei*, Horsf.

Fam. EMBALLONURIDÆ.

- Emballonura semicaudata*, Peale.
- *monticola*, Temm.
- Taphozous melanopogon*, Temm.
- *longimanus*, Hardw.
- *affinis*, Dobson.
- Chiromes torquatus*, Horsf.
- Nyctinomus plicatus*, Buch. Ham.

INSECTIVORA.

Fam. SORICIDÆ.

- Chimarrogale himalayica*, Gray.
- Crocrida fuliginosa*, Blyth.
- * — *fetida*, Peters.
- * — *doriae*, Peters.
- Pachyura indica*, Geoffr.
- *hosei*, Thos.

Fam. ERINACEIDÆ.

- Gymnura rafflesi*, Vigors.
- Hylomys suillus*, Müll.

Fam. TUPAIIDÆ.

- Ptilocercus lowi*, Gray.
- Dendrogale murina*, Müll.
- Tupaia javanica*, Horsf.
- * — *minor*, Günth.
- * — *melanura*, Thos.
- *ferruginea*, Raffles.
- * — *splendidula*, Gray.
- *tana*, Raffles.
- * — *chrysuræ*, Günth.
- * — *dorsalis*, Schleg.
- * — *picta*, Thos.
- * — *montana*, Thos.

Fam. GALEOPTHECIDÆ.

- Galeoptithecus volans*, Linn.

CARNIVORA.

Fam. MUSTELIDÆ.

- Putorius nudipes*, F. Cuv.
- Mustela flavigula*, Bodd.
- Mydaus meliceps*, F. Cuv.
- ** — *marchei*, Huet¹.
- Lutra sumatrana*, Gray.
- *cinerea*, Illiger.

Fam. URSIDÆ.

- Ursus malayanus*, Raffles.

Fam. CANIDÆ.

- ? *Cyon rutilans*, Müll.

Fam. VIVERRIDÆ.

- Herpestes brachyurus*, Gray.
- * — *semitorquatus*, Gray.
- Cynogale bennetti*, Gray.
- Arctictis binturong*, Raffles.
- Hemigale hardwickei*, Gray.
- * — *hosei*, Thos.
- Arctogale leucotis*, Blyth.
- Paradoxurus leucomystax*, Gray.
- *hermaphroditus*, Pallas.
- *philippinensis*, Jourdan.
- Linsang gracilis*, Müll.
- Viverra zangalunga*, Gray.

Fam. FELIDÆ.

- Felis planiceps*, Vig. & Horsf.
- * — *badia*, Gray.
- *temminckii*, Vig. & Horsf.
- *bengalensis*, Kerr.
- *marmorata*, Martin.
- *nebulosa*, Griffith.

RODENTIA.

Fam. HYSTRICIDÆ.

- * *Trichys guentheri*, Thos.
- * *Hystrix crassispinis*, Günth.
- ** — *pumila*, Günth.
- *muelleri*, Jent.

Fam. MURIDÆ.

- Mus rattus*, Linn.
- *decumanus*, Pallas.
- * — *infraluteus*, Thos.
- *muelleri*, Jent.
- * — *sabanus*, Thos.
- *hellwaldi*, Jent.
- *jerdoni*, Blyth.
- * — *alticola*, Thos.
- *musechenbroeckii*, Jent.
- *eplippium*, Jent.
- *musculus*, Linn.
- * — *margarettæ*, Thos.
- * *Chiropodomys major*, Thos.
- * — *pusillus*, Thos.

Fam. SCIURIDÆ.

- Sciuropterus pulverulentus*, Günth.
- *horsfieldi*, Waterh.
- *setosus*, Temm.
- *genibarbis*, Horsf.
- ** — *nigripes*, Thos.
- Pteromys nitidus*, Desm.
- * — *phaemelas*, Günth.
- * *Rhithrosciurus macrotis*, Gray.

¹ Huet, Le Naturaliste, 1887, p. 149.

- Sciurus bicolor*, *Sparrrn.*
 — *ephippium*, *Müll.*
 — *prevosti*, *Desm.*
 — *hippurus*, *Geoffr.*
 * — *pryeri*, *Thos.*
 * — *brookei*, *Thos.*
 — *tenuis*, *Horsf.*
 * — *lowi*, *Thos.*
 * — *jentinki*, *Thos.*
 — *notatus*, *Bodd.*
 — *insignis*, *Desm.*
 * — *hosei*, *Thos.*
 * — *everetti*, *Thos.*
 ** — *steerei*, *Günth.*
 — *laticaudatus*, *Müll.*
 — *soricinus*, *Müll. & Schl.*
 — *exilis*, *Müll.*
 * — *whiteheadi*, *Thos.*

UNGULATA.

- Fam. ELEPHANTIDÆ.
Elephas indicus, *Linn.*
 Fam. RHINOCEROTIDÆ.
Rhinoceros sumatrensis, *Cuv.*
 Fam. TAPIRIDÆ.
Tapirus indicus, *Cuv.*
 Fam. BOVIDÆ.
Bos sondaicus, *Müll.*
 — *buffelus*, *Blumenb.*
 Fam. CERVIDÆ.
Cervus equinus, *Cuv.*
 — *hippelaphus*, *Cuv.*
 — sp. inc.
Cervulus muntjac, *Linn.*

- Fam. TRAGULIDÆ.
Tragulus napu, *F. Cuv.*
 ** — *nigricans*, *Thos.*
 — *javanicus*, *Gmel.*
 Fam. SUIDÆ.
Sus vittatus, *Müll.*
 — *verrucosus*, *Müll.*
 * — *barbatus*, *Müll.*
 — *longirostris*, *Nehring.*
 ** — *ahænobarbus*, *Huet*¹.

CETACEA.

- Fam. DELPHINIDÆ.
Delphinus sp. inc.
Orcella brevirostris, *Owen.*
 Fam. BALÆNIDÆ.
Balænoptera sp. inc.
 (*Megaptera*?)

SIRENIA.

- Fam. HALICORIDÆ.
Halicore dugong, *Erxl.*

EDENTATA.

- Fam. MANIDÆ.
Manis javanica, *Desm.*
 — sp. inc.²

4. Brief Notes on Flukes. By PROSPERO SONSINO, M.D.,
 Pisa. (Communicated by Prof. F. JEFFREY BELL.)

[Received May 16, 1893.]

DISTOMUM TRIGONOCEPHALUM, R.

In the collection of Worms in the British Museum (Natural History), which, through the kindness of Dr. Günther and Prof. Jeffrey Bell, I was enabled to examine in 1891 while attending the International Hygienic Congress, my attention was attracted to a bottle containing a worm which was labelled "*Ophistomum mucronatum*, R.," but which, some days previous to my visit, my friend Prof. Blanchard, after a cursory examination, had decided to be a *Distomum*. Prof. Blanchard had not sufficient time to examine the anterior end of the single specimen, which was lying upon the dorsal side of the body, and so he failed to perceive that this

¹ Huet, Le Naturaliste, 1888, p. 5.

² Marche, Luçon et Palaouan. 1887. p. 302.

Distome belongs to the subgenus *Echinostomum*, Dujardin, the oral sucker being surrounded by a disk bordered by a circle of spines. Dr. L. Örley, who some years ago published an account of the Nematodes from the collection of Prof. Siebold in the Museum¹, has not mentioned there the supposed *Ophiostomum*, therefore we may fairly infer that he had not examined it.

In the unique specimen of the worm in question, notwithstanding its rather bad condition, the following characters may be verified:—Body flat, elongated, rather narrowed behind; length 7 millim., breadth 2 millim., with the greatest diameter corresponding to the ventral sucker, which is larger than the oral sucker. The latter surrounded by a triangular disk, bordered by spines. The number of these latter appeared to me to be 24, comprising the 4 lateral and inferior larger. A darker spot behind the ventral sucker is due to the oviduct being full of eggs, which are of large size and not less than 100 micromillim. in length. Yolk-glands rather voluminous from the ventral sucker to the posterior end. Two oval testes at the beginning of the posterior half of the body, one behind the other. Ovarium anterior to the testes.

With such characters I suspected that the Trematode was *D. trigonocephalum*, the ordinary *Echinostomum* found in mammals; but before pronouncing myself definitely, I wished to compare the characters of the specimen in question with those of *D. trigonocephalum*. My short visit to the British Museum not allowing me to make this comparison there, I made it when I returned home. I found in the collection of the Zoological Museum of Pisa no less than four bottles containing *D. trigonocephalum* obtained from three different species of Mammals, viz. *Mustela foina*, *M. putorius*, and *Meles europæus*. Thus I had plenty of material to give me a clear notion of the characters and different appearances of *D. trigonocephalum* from various sources. I am now able to say that the specimen of the Trematode of *Plecotus communis* in the collection of the British Museum is, both as regards size and conformation of the body, as well as the disposition of the spines around the oral sucker, much like *D. trigonocephalum*. There is also a similarity in the eggs. Only the testes of the *Echinostomum* in the British Museum appear to be situated rather more behind; but I do not attach much importance to this. Indeed the number of the spines round the head given by von Linstow² and verified in many specimens by myself is 26 instead of 24. But it is possible that I mistook the number when examining the specimen. I think, therefore, that the specimen in the British Museum is nothing else but *D. trigonocephalum*. In his 'Compendium,' Linstow does not make mention of such a Trematode as parasitic in any Bat. But *D. trigonocephalum* has been found in *Erinaceus europæus*, and it is rather interesting to know that it may also be parasitic in a Bat. Thus it is found in animals belonging to no

¹ Ann. & Mag. Nat. Hist. ser. 5, vol. ix. p. 301 (1882).

² Troschel's Archiv, i. p. 106 (1873).

less than three different orders of Mammals, undoubtedly on account of the food that they may partake of in common.

DISTOMUM OVO-CAUDATUM, Vulpian.

Distomum ovo-caudatum, both of *Rana esculenta* and of *R. temporaria*, has been described since 1859, by Vulpian, as living in the mouths of these animals. Vulpian gave an interesting account of this worm and has not failed to put in evidence the important character of the presence of a basal filament in the egg,—a character very rarely to be seen in the digenetic Trematodes, but frequently observed in monogenetic or ectoparasitic Trematodes. Recently Creutzburg has published an interesting study of the life-history of the same worm, which according to him in its larval form is the famous *Cercaria cystophora*, Wagener, which was once referred by Willemoes-Suhm to *D. lanceolatum*, Mehlis. According to Creutzburg, the intermediary host of *D. ovo-caudatum*, a *Planorbis*, would become infested not by active introduction of the embryo into its body, the embryo being unable to swim in water, but by eating the eggs with the contained embryo, which eggs are found in the faeces of frogs. In a similar way the final host would be infested by eating *C. cystophora* contained in its nurse within the tissues of the mollusk. But as yet Creutzburg has not succeeded in rearing the adult worm in the mouth of a frog, by introducing the tissues of *Planorbis* containing *C. cystophora*.

I am certainly not prepared to criticise Creutzburg's work, which is only preliminary; but I wish to point out some facts observed by myself, which are not in accordance with those noticed either by Creutzburg or Vulpian, respecting the worm in question. I have had good opportunities to observe and study *D. ovo-caudatum*, which I have found rather frequent in both *Rana esculenta* and *R. temporaria* in the neighbourhood of Pisa. I may now draw the following conclusions as to the result of my observations, of which I gave a full account in the 'Monitore Zoologico' of last April:—(1) *D. ovo-caudatum* takes up its abode not only in the mouth, but also in the stomach and the upper part of the intestines. It is perhaps more frequent in the stomach than in the mouth, and it is possible that its occurrence in the mouth is merely accidental, and the result of an emigration, occasioned by the distressed condition of the worm on the death of the host. (2) The basal filament is from 4 to 6 times as long, or even longer, than the length of the body of the egg. If previous observers have given a shorter length, it is probably in consequence of their having seen only a truncated filament, or of not having properly focussed the microscope so as to see the entire length of the filament, which gradually thins towards the end. (3) The embryo is not only armed with a crown of spines (*bâtonnets*) on its anterior extremity, but evidently possesses even cilia. Whether these cilia aid it to swim in water when it issues from the egg, spontaneously and completely mature, or are simply for locomotion in the intestines of *Planorbis*, I am not prepared to say, not having seen the

embryo alive and swimming in water, but only the embryo coming out from the shell by pressure. In this case the embryo has always died and has often presented the body spoiled and without the ciliar tegument, so that I was only once able to observe a specimen with the cilia in motion.

FLUKES OF EXOTIC SNAKES.

In a bottle in the collection of parasitic worms in the Museum of Pisa I found, in 1890, a *Distomum* obtained from the renal pelvis of *Python molurus*. I described it under the name of *Distomum simile*¹, regarding it as a new species, resembling in size and structure *D. lanceolatum*, with the exception that the two testes, instead of being anterior to the ovary, were situated behind that organ. Other examples of a similar Distome, obtained from the liver of an animal of the same species, I thought, but could not be certain, owing to decomposition of the specimens, might also be *Distomum simile*. At the time I could not compare *Distomum simile* with the *Distomum horridum* of Leidy, because I had not access to Leidy's description. But lately, through the kindness of Dr. Stiles, I have received a specimen of Leidy's *Distomum horridum* (obtained from a Boa Constrictor), and from this I have been able to satisfy myself of the identity of *D. simile* and *D. horridum*. I have since read Leidy's paper² describing the parasite, and although in some respects his description is somewhat incorrect, as he speaks of four testes instead of two, and of vitellaria as ovaria, I can recognize in it the identity of *Distomum simile* with *Distomum horridum*.

Another species which appears very similar to *D. horridum* is the *D. sauromatis*, described as new by M. J. Poirier. The principal difference between *D. simile* and *D. horridum* on the one hand, and *D. sauromatis* on the other, is that in the latter the two testes are lobate instead of being round as in the former. This variation is so slight that it may merely depend on the diverse habitats of the individuals and not on difference of species. I cannot, however, definitely pronounce on this point, not having at hand a specimen of Poirier's *Distomum*, which lives in the lungs of *Elaphis sauromates*. But the identity of *Distomum simile* with *D. horridum* is very interesting, as it shows that the same species of *Distomum* lives in two different species of Snakes inhabiting different countries.

DISTOMUM BARALDII.

Distomum baraldii is a parasite which I have lately described as a new species found in *Zamenis viridiflavus*³, and which I have

¹ See 'Processi verbali della Società Toscana di scienze naturali residente in Pisa,' Adun. 4 Maggio, 1890.

² Journal Acad. Natur. Sciences, series 2, i. p. 303 (Philadelphia).

³ *D. baraldii*, given by me as a new species, may prove to be identical with *Distomum nigrovenosum*, Bellingham, found in *Tropidonotus natrix*. At least my new species much resembles the description and figure given by my friend Dr. Monticelli of Bellingham's fluke in an unpublished paper with which I have made acquaintance by the author's kindness.

often obtained in Pisa from the mouth and gullet of the Snake referred to. The colour of this small Distome is quite characteristic, being white in the anterior and black in the posterior half. But the most interesting point in connection with this Distome is that I often found in *Zamenis viridiflavus* an imperfectly developed Distome encysted in the submucosa of the mouth, in the pericardium, and in the mesenterium; and as I obtained the cysts sometimes empty, sometimes full, and as the full cysts when placed in water rupture quickly and allow the young Distome to escape, and there is some resemblance between this young Distome and *D. barallii*, I suspect that the encysted and the mature Distomes belong to the same species, and that, contrary to the usual course in the evolution of the Distomes, the same animal plays the part both of intermediary and final host to this parasite.

5. On a Second Collection of Mammals sent by Mr. H. H. Johnston, C.B., from Nyasaland. By OLDFIELD THOMAS.

[Received May 16, 1893.]

In the 'Proceedings' of this Society for last year¹ I had the honour of giving an account of a fine collection of Mammals presented to the National Museum by Mr. H. H. Johnston, Consul-General for Mozambique, and H.M. Commissioner for Nyasaland. That distinguished explorer has now sent home a second series, collected, like the first, by Mr. Alexander Whyte, at Zomba and Milanji. Of these, by the kindness of our Secretary, I am now permitted to give a list, supplementary to the former one, and carrying on a little further our knowledge of the Nyasaland fauna, to the study of which Mr. Johnston and Mr. Whyte have rendered such material aid.

The present collection contains between 70 and 80 specimens, belonging to 30 species, of which a large number are additional to those previously sent. As with the previous set, the great interest of the collection lies in its having been made so near to the localities where Dr. Peters obtained the material on which his classical work on the Mammals of Mozambique was founded. Such a collection as this, therefore, although containing no novelties, is, and will always continue to be, of the greatest service to English naturalists, as giving them the means of accurately comparing specimens from other parts of Africa with examples certainly corresponding with those described by Peters.

With regard to the interesting questions as to the seasonal changes of fur, which so sorely need investigation, it unfortunately happens that the present collection was almost entirely formed in the months of October and November, the very same months when the previous series were obtained. It would therefore be very desirable for Mr. Whyte to try and collect specimens of all the

¹ P. Z. S. 1892, p. 546.

species now and previously sent at other times of the year. Seasonal variation has been but little studied in African mammals, and specimens obtained by so able a collector all from the same place and at different seasons would have the utmost value as enabling us to form a comprehensive idea as to the general influence that the changes of such a climate as that of Nyasaland have on its Mammalian inhabitants.

Pending the formation of such a supplementary collection, the present series, full as it is of rarities from a Museum point of view, demands for publication little more than a nominal list of the species sent.

1. *OTOGALE KIRKII*, Gray.

a. Ad. sk. Nmasi. 9/92.

This skin is not in a good state, and the fur appears to be in process of change. There seems, however, to be little doubt that it represents *O. kirkii*, whether that does or does not prove in the end to be a valid species.

2. *PETRODROMUS TETRADACTYLUS*, Pet.

a. Ad. al. ♀. Milanji. 10/92.

3. *CROCIDURA* sp. inc.

a. Ad. al. Milanji. 10/92.

4. *CROCIDURA* sp. inc.

a. Ad. al. Milanji. 10/92.

5. *HERPESTES GRACILIS*, Rüpp.

a, b. 2 ad. sks. Zomba. 11/92.

These specimens evidently belong to the S.-African type of the species, and show that Peters's *H. ornatus* should be referred to that form rather than to the typical variety, to which I assigned it in 1882.

6. *CROSSARCHUS FASCIATUS* (Desm.).

a, b. Ad. sks. Tschiromo, junction of Rivers Ruu and Shiré. 9/92.

C. fasciatus is recorded from Mt. Kilimanjaro by Mr. True under the name of *C. mungo*, Gmel.¹ That name is evidently taken from the synonymy of the species as given in my monograph of the African Mongoose², Mr. True rightly ignoring my plea for the retention of the name *fasciatus* merely as being classical, well-known, and appropriate. He does not, however, seem to be aware that the name *mungo*, originally given to a mixture of the common Indian Mongoose and the S.-African striped one, has been revived by Mr. Blanford³ as the appellation of the former, in spite of my

¹ P. U.S. Nat. Mus. xv. p. 453 (1892).

² P. Z. S. 1882, p. 90.

³ P. Z. S. 1887, p. 631; Mamm. Brit. Ind. p. 123 (1888).

own argument in favour of *Herpestes griseus*¹. Even if, therefore, this barbarous name has to be used, it clearly cannot be applied to two species; and since Schreber's description of "Die Manguste—*Viverra ichneumon* β,"² on which *Viverra mungo*³ was founded, has much more reference to the Indian than to the African animal, I am now of opinion, with Mr. Blanford, that the name of *mungo* should be used for the former, and, for more valid reasons than in 1882, *Crossarchus fasciatus* (Desm.) for the latter.

7. *CANIS* sp. inc.

a. Yg. sk. Upper Shiré. 11/92.

Too young for determination.

8. *SCIURUS MUTABILIS*, Pet.

a-e. 5 ad. sks. Zomba. 10/92.

Pending the arrival of specimens collected at other times of the year, I shall not express any opinion as to the cause of the considerable variation in colour existing among the present series of this Squirrel.

9. *GERBILLUS AFER*, Gray.

a, b. 2 ad. sks. Fort Johnston. 11/92.

10. *OTOMYS IRRORATUS*, Bts.

a. Ad. sk. Upper Shiré. 11/92.

11. *GOLUNDA FALLAX*, Pet.

a-e. 5 sks. Zomba. 10/92.

12. *MUS (DASYMYS) INCOMTUS*, Sund.

a, b. Ad. sks. Upper Shiré.

This rare species has been hitherto only obtained in Natal, although a closely allied and possibly identical form (*Mus nudipes*, Peters) has been described from Angola⁴.

13. *MUS RATTUS*, var.

a. Ad. sk. Upper Shiré.

b-d. 3 yg. sks. Zomba. 10/92.

14. *MUS DOLICHURUS*, Smuts.

a. Ad. sk. Tschiromo. 9/92.

b. Ad. sk. Zomba. 10/92.

15. *MUS NATALENSIS*, A. Sm.

a. Ad. sk. Tschiromo. 9/92.

b. Ad. sk. Zomba. 10/92.

c-e. 3 ad. al. Milanji. 10/92.

¹ P. Z. S. 1836, p. 56 (ftnote).

² Schr. Säug. iii. p. 430 (1777).

³ Gmel. Linn. S. N. i. p. 84 (1788).

⁴ See Thos. Ann. Mag. N. H. (6) x. p. 180 (1892).

16. *MUS MUSCULUS*, L.

a, b. Ad. sks. Zomba. 10/92.

c. Ad. al. Milanji. 10/92.

17. *MUS* (*LEGGADA*) *MINUTOIDES*, A. Sm.

a-c. Ad. sks. Zomba. 10/92.

18. *CRICETOMYS GAMBIANUS*, Waterh.

a. Ad. sk. Zomba. 10/92.

19. *SACCOSTOMUS CAMPESTRIS*¹, Pet.

a. Ad. sk. Upper Shiré. 11/92.

b-f. Ad. & imm. al. Fort Johnston. 11/92.

20. *STEATOMYS PRATENSIS*¹, Pet.

a-c. Ad. sks. Zomba. 10/92.

d. Ad. al. ♂. Milanji. 10/92.

These rare and peculiar short-tailed African Muridæ, with their doubtful affinities, are of the greatest interest, and it is probable that a clue to some of the most difficult problems of Murine phylogeny will be gained by a study of their teeth. For these reasons specimens of *Saccostomus* and *Steatomys* are most acceptable, and it is to be hoped that Mr. Whyte will also come across examples of the still rarer *Malacothrix* and *Mystromys*, approximately similar in size and proportions, but with very different teeth.

Although similar to each other superficially, these four last-named genera appear to belong to three different subfamilies—*Mystromys* seeming to be strictly Cricetine; *Steatomys*, as Alston has stated², being closely allied to *Dendromys*, as also are *Malacothrix* and *Dendromys*; while *Lophuromys*, *Saccostomus*, and *Cricetomys* ought, I believe, all to be placed in the restricted subfamily Murinæ. Of these last three the first was placed by Alston in the *Dendromyina* and the other two in the *Cricetina*.

21. *DENDROMYS MESOMELAS*, Bts.

a. Ad. sk. Zomba. 10/92.

b-f. 5 in al. Milanji.

These specimens, although I place them provisionally under the oldest name, *D. mesomelas*, certainly seem to differ constantly from the ordinary form of that species in being entirely without the black dorsal line and in being considerably smaller. In these respects they agree with two out of the previous collection, while two others are of the usual large-striped character. They thus correspond with Wagner's *D. pumilio*, described in 1841³, but afterwards⁴ considered by him as a synonym of *D. mesomelas*. Larger series from different localities are, however, needed before one can

¹ For nomenclature see Ann. Mag. N. H. (6) x. p. 264 (1892).² P. Z. S. 1876, p. 82.³ Münch. Gelehrt. Anzeiger, xii. p. 437.⁴ Schr. Säug. Supp. iii. p. 464 (1843).

be certain as to whether the absence of the black stripe is due to old age (as said by Smith), youth (as said by Wagner), season, or locality.

22. *MYOSCALOPS ARGENTEO-CINEREUS*, Pet.

a-j. 10 ad. & imm. sks. Zomba. 10/92.

23. *POTAMOCHÆRUS LARVATUS*, F. Cuv.

a. Ad. sk. & skull. Zomba.

b, c. 2 yg. sks. Zomba. 11/92.

24. *BUBALIS LICHTENSTEINI*, Pet.

a. Ad. sk. & skull. Zomba.

This specimen, equally with that mentioned in the previous paper (p. 553), has the whitish rump referred to by Dr. Matschie as characteristic of his *B. leucoprymnus*¹. Since both individuals were obtained, however, at quite a short distance from the typical localities (Tette and Sena on the Zambesi) of Peters's species, there can be no doubt that they represent the true *lichtensteini*. Whatever therefore may be the differences in the horns between "*lichtensteini*" and "*leucoprymnus*," it is clear that the whiteness of the rump in the northern form cannot be used as a distinguishing character.

25. *OREAS CANNA*, H. Sm.

a. Ad. skull. ♀. Zomba.

26. *HIPPOTRAGUS NIGER*, Harr.

a, b. 2 ad. skulls. Zomba.

27. *TRAGELAPHUS SCRIPTUS ROUALEYNI*, Gord.-Cumm.

a-f. 6 skins. Zomba.

These specimens belong clearly to the Lower Zambesi form, their white markings, although very variable, being far less developed than in the Chobe River Bush-buck, assigned in 1891 to the true *T. scriptus typicus*, Pall.²

28. *NANOTRAGUS SCOPARIUS*, Schr.

a. Ad. sk. ♂. Zomba.

29. *COBUS ELLIPSIPRYMNUS*.

a-d. 4 skulls. Zomba.

30. *CEPHALOLOPHUS GRIMMIUS*, L.

a. Ad. sk. & skull. Zomba.

¹ SB. Ges. nat. Fr. 1892, p. 137.

² P. Z. S. 1891, p. 380.

FEMALE WATER-BUCK & YOUNG, (1893).

J. Smit del. et lith.

Wintern Bros. imp.