blankets were disinfected twice or thrice, the stable periodically disinfected—all of which means of disinfection were carried out at one and the same time, and the grooming kit was steeped in a strong solution of chloride of lime once daily.

Means of Prevention.—Conjointly with treatment of the affected, means of suppression of any source of contagion should be adopted, and that which is to be recommended is clipping each stableful of horses, scrubbing them with soap and water, followed by dressing with a weak solution of some parasiticide; each part of the leather-work of the saddlery may be washed with hot water, and the animals' blankets and such like steeped in a watery solution of a parasiticide, or boiled (in the latter case it will be advisable to remove temporarily the leather-work); at the same time most thorough scrubbing of every part of the stables with hot water and a solution of chloride of lime, assisted if necessary by scraping the walls, mangers, and hay-racks; and the grooming kit should be steeped daily in an anti-parasitic remedy; the bedding of the cleansed stables should be burned, and if considered necessary no fresh material used for three or four nights. The above process of disinfection may be repeated as considered necessary.

Inspections.—On the occurrence of a case of scabies every horse in the stud should be most carefully and systematically inspected as frequently as possible (once a day if that be practicable), to detect any horse showing symptoms in the earliest stage. An error will be committed if grooms or any such person attending to the horses be trusted to report cases, because the majority of such men will bring to the notice of veterinary surgeons only cases that are well marked or show signs of itch; if we mean to suppress an outbreak we must get hold of the cases in the earliest stage.

HÆMOGLOBINURIA OF THE HORSE.

By J. M'FADYEAN, M.B., B.Sc., Royal Veterinary College, London.

To the opening volume of this Journal I contributed an article on Hæmoglobinuria (Azoturia) of the Horse, in which I discussed at considerable length the then prevalent views regarding the pathology of the disease, and described my own observations regarding its lesions. At that date the view generally held regarding the nature of the disease by veterinary surgeons in this country was that which found expression in the article on Azoturia in Professor Williams' text-book of Veterinary Medicine, that is to say, the essence of the disease was supposed to be "a hyper-nitrogenous condition of the blood and system generally, due to over-feeding and want of exercise; the excessive secretion of urine and excretion of urea being physiological results due to the presence of effete products, metamorphoses of nitrogenous food in the blood."

In the article above referred to I contended that this view of the pathology of the disease was erroneous and even absurd, and I showed that while there was no evidence that there was a constant or notable increase of urea in either the blood or the urine, there was easily

discoverable evidence that both the blood and the urine did contain an excessive amount of an other substance, viz., hæmoglobin or some allied pigment. I showed that this pigment exists in solution in the blood plasma, that after death it there frequently assumes the crystalline form, and that during life it is excreted by the kidneys and colours the urine. Furthermore, I showed that in fatal cases of hæmoglobinuria the *post-mortem* reveals a very important and interesting lesion of the muscles.

The object of the present article is a double one. In the first place, I feel that it is necessary, although it involves some repetition of what I have already written on the subject, to controvert for a second time the uraemic or nitrogenous theory, freshly propounded in the proceedings of the National Veterinary Association for the current year. Secondly, I wish to place on record some new observations that I have been able to make regarding the morbid anatomy of this disease.

In what is entitled a Summary of the Reports sent in by members of the National Association, Professor Williams states that "there is in the reports to be found the greatest discrepancy in the analysis in regard to urea, although all recognise its evident increase." How a statement so directly contrary to the fact came to be made it is difficult to imagine. A perusal of the Reports shows that the percentage of urea found in the various samples of urine analysed was as follows:—

SERIES I.—Analysed by Professor Bayne.

<i>Sample.</i>	<i>Percentage of Urea.</i>
1	1·96
2	2·05
3	2·43
4	2·03
5	1·89
6	3·11
7	2·57
8	2·09
9	1·35
10	1·01

Calculation shows that the average percentage of urea in these 10 samples was 2·049.

SERIES II.—Analysed by Professor Cooke.

<i>Sample.</i>	<i>Percentage of Urea.</i>
1	1.40
2	3.02
3	3.94
4	3.00
5	1.83
6	2.29
7	1.13
8	1.84
9	1.08
10	1.52
11	1.02
12	1.51
13	1.65
14	1.84
15	1.38

The average percentage of urea in these 15 samples will be found to be 1.89.

SERIES III.—Analysed by Professor Smith.

<i>Sample.</i>	<i>Percentage of Urea.</i>
1	4.3
2	1.984

The average percentage of these two samples is 3.142.

One other analysis is given in the Reports, viz., one said to have been made of a sample of urine by Mr Greig (Mr Greig Smith?), for Professor Walley. In this case the proportion of urea is stated to have been 44.05 grains per ounce, or about 9 per cent. This I take to be an obvious error, probably a misprint for 4.405 grains per ounce, for the sample of urine had a low specific gravity, viz., 1.041. But even including this case, it will be found that the average percentage of urea in the entire series of analyses given in the Reports is 2.29.

In this connection it ought to be remembered that the analyses were made by the sodium hypobromite method, and that therefore the percentage of urea given in reality indicates the total nitrogenous constituents of the urine calculated as urea.

What, it may now be asked, is the normal percentage of urea contained in the urine of horses? In an analysis by Boussingault, quoted by Colin, the proportion of urea in the urine of healthy horses is given as 13 grains per ounce, which is equivalent to a little over 3 per cent. In Meade Smith's *Text-Book of Physiology* the percentage of urea in healthy horses receiving ordinary diet is stated to vary from 2.5 to 4 per cent. Lastly, in F. Smith's *Veterinary Physiology* the normal amount of urea excreted daily by the horse in health is stated to be about $3\frac{3}{4}$ ounces, or a little over 2 per cent., the average amount of urine passed daily being $8\frac{1}{2}$ pints.

It is thus evident that the analyses quoted do not show that there is an "evident increase" of urea in the urine of horses suffering from the disease under consideration; on the contrary, they corroborate the previously published analyses, and prove most conclusively that in hæmoglobinuria the proportion of urea excreted suffers no appreciable disturbance.

In Professor Williams' summary in the Proceedings of the National Veterinary Association the following sentence occurs: "Since the malady was called azoturia by me in the first edition of my work on veterinary medicine, some dissatisfaction has been expressed and fault found for the reason that the term is vague and throws no light upon the pathology of the disease. This I grant, but at the same time have to state that it is quite as expressive as its substitute. It simply means urine containing a superabundant quantity of nitrogenous products." To this it may be replied that dissatisfaction has been expressed and fault found with the name azoturia, not because it throws no light upon the pathology of the disease, but because it falsely implies that the urine in that disease contains an excess of urea.

It is probable that some people will continue to call this disease by the term azoturia, for when once a name has been for some years in general use it is difficult to eradicate it. Moreover, if we continued to use the word azoturia merely as a name for the disease, equivalent in purpose to the number that distinguishes a policeman, no objection need be taken to its employment, but the matter is different when we are advised to keep the name in use because it indicates an important fact in connection with the pathology of the disease.

The pathology of hæmoglobinuria is in several points still obscure, but at the same time a great deal more is known regarding it than anyone can learn by reading the summary of the evidence collected by the Investigation Committee of the National Veterinary Association.

The following points regarding the disease have for a number of years been known to everyone who has taken the trouble to follow the literature on the subject:—

1. During an attack the liquor sanguinis of the affected horse contains in solution a large quantity of hæmoglobin or some nearly allied pigment. The presence of this pigment can easily be placed in evidence during life by abstracting a quart of blood from the jugular into a clean vessel, and setting

this vessel aside for 24 hours. At the end of that time the blood will have separated into clot and serum, and if the latter be withdrawn by the aid of a pipette it will be found to be of quite a different tint from the serum yielded by the blood of a healthy horse. The latter has an amber tint, but the serum yielded by the blood of a horse that is the subject of hæmoglobinuria when examined in a half-inch test-tube has a beautiful deep ruby tint.

As I pointed out in the article published in 1888, this pigment may sometimes after death be readily demonstrated in another way, viz., by placing a drop or two of blood on an ordinary microscope slide and then putting on a cover-glass. It sometimes happens that in the course of a few minutes the dissolved pigment assumes the solid form, and when the preparation is then examined under the microscope the field of view is almost entirely occupied by flat and acicular crystals.

2. The urine passed during an attack of hæmoglobinuria is abnormal, in that it always contains albumen and pigment. The pigment exists partly in solution, and often in greater proportion in the form of very minute, dusky, opaque, granules. The urinary pigment is derived from the hæmoglobin circulating in solution in the blood plasma, the kidneys taking upon themselves the work of excreting that substance. The hæmoglobin is not excreted as such, but is, at least for the most part, split up, and this is probably the source of both the albumen and the granular pigment.
3. The statement that the muscles of the hips, loins, or other parts are swollen during an attack of hæmoglobinuria is erroneous. The appearance of swelling, when present, is due to the muscles being in a state of rigid contraction, and it is this rigidity that is the cause of the wooden hardness of certain muscles at the onset of the attack. In fatal cases the *post-mortem* reveals in the muscles of the hip, loins, or pectoral region a remarkable degeneration. This degeneration is quite evident to the naked eye, in the form of a pallor of more or fewer of the muscular fasciculi. These exhibit, instead of the normal dark red tinge, a greyish or ashy colour, somewhat similar to that obtainable by macerating a piece of muscle in water. When subjected to microscopic examination the altered fasciculi show a pronounced "waxy degeneration" affecting more or fewer of the muscular fibres. The degenerated fibres have lost their normal colour, the striation also is lost, and the sarcois substance has been converted into a transparent homogeneous material.

In the foregoing paragraphs I have endeavoured to state briefly what is known, and has been known for some years, regarding the morbid anatomy of hæmoglobinuria. The next point is to consider what light the morbid changes mentioned throw on the pathology of the disease.

The first question that naturally suggests itself is, What is the source of this pigment which exists in such abnormal amount in the liquor sanguinis during the acute stage of an attack? One can conceive of two possible sources. It may be hæmoglobin set free by

destruction of red blood corpuscles, or it may be the colouring matter (believed to be identical with or nearly related to hæmoglobin) liberated from the degenerated muscular fibres. This is really the first problem now requiring solution in connection with the pathology of the disease.

It may be observed that either supposition regarding the source of the pigment will to a certain extent fit in with the known facts. It has been shown experimentally that the injection into the blood-stream of substances that act destructively on the red corpuscles, and thus liberate their hæmoglobin, sets up a train of symptoms that closely simulate those met with in the disease now under consideration. As in the latter disease, the kidneys take up the excretion of the liberated hæmoglobin, and there is consequent hæmoglobinuria. Furthermore, some of the liberated pigments may take the granular form, occlude the small arteries and capillaries of certain muscles, and throw these into a state of rigid contraction, with subsequent degeneration of the sarcous substance.

On the other hand, if we assume that the pigment is of muscular origin, and due to the destruction of muscular fibres brought about by some ferment or irritant circulating in the blood, we can still explain the deep ruby colour of the blood-serum, and the appearance of pigment in the urine.

There are several ways in which evidence that would be of value in clearing up this obscure point might be obtained. In the first place, an endeavour might be made to discover whether at the very onset of the attack—as soon as ever the horse is seized with the well-known muscular symptoms—the bladder already contains a considerable quantity of abnormal urine, that is, urine containing albumen and pigment. If this were clearly proved to be the case it would indicate that the source of the pigment must be sought elsewhere than in the muscles. It would, in short, indicate that the affection of the muscles is the consequence and not the cause of the hæmoglobinæmia.

Or again, the problem might be attacked by the experimental way, introducing into the blood-stream substances that act destructively on the red corpuscles, and observing whether the train of symptoms set up and the lesions discovered after death are identical with those encountered in the natural disease.

Lastly, from the onset of an attack observations might be made at intervals with the hæmocytometer, in order to discover whether, *pari passu* with the excretion of pigment in the urine, there is a diminution in the number of red corpuscles. It is regarding this method of attacking the question that I desire to place on record some observations, but before doing so it may be well to quote a passage from my former article on this subject: "There are great difficulties in the way of accepting the theory that the hæmoglobin which is dissolved in the blood plasma and excreted by the kidneys, results from destructions of red cells. In the first place, if such were the case, one would expect to find, by a careful microscopic examination of the blood, evidence of it in the shape of red corpuscles in various stages of destruction. Moreover—and this is a more fatal objection—such extensive blood disintegration as would be necessary to furnish the great amount of free hæmoglobin found in some cases would lead to profound anæmia. This, in fact, is what we do see in the

hæmoglobinuria or red-water in cattle. But even in horses that have survived for several days, and in which there has been during the whole period a rapid excretion of pigment, one does not find a sensible diminution in the number of red corpuscles."

The statement above made, that an attack of hæmoglobinuria does not entail a condition of anæmia, I could not at the time support by accurate figures, but within the last twelve months I have been able to test the matter by the use of the hæmocytometer, and the results thus obtained were of a totally unexpected character.

The hæmocytometer, or corpuscle counter, it may be well to premise, is an ingenious instrument comprising : (1) a miniature glass pipette in which a minute quantity of blood is diluted to a known extent (1 : 100 or 1 : 200) with an artificial serum ; (2) a small glass cell making part of a microscope slide and intended for the reception of the diluted blood. A spot on the floor of the glass cell is ruled with parallel and intersecting lines so as to divide it into microscopic squares. A droplet of the diluted blood is placed over this spot and covered with a cover-glass, and the slide is then placed under the microscope. In a few minutes the red corpuscles have settled to the bottom of the cell, and when the floor of the latter is focussed one sees the small squares bounded by the intersecting lines and at the same time the red corpuscles lying within these. The average number of corpuscles in a large number of squares is ascertained, and since each square is the bottom of a column of diluted blood of known volume a simple sum in arithmetic gives the number of red corpuscles in a cubic millimetre of the blood under examination. It may be added that observations with this instrument have shown that the number of red corpuscles in a cubic millimetre of blood from a healthy horse is from 7 to 8 millions.

My first observations on the point were made in December last on a typical case of the disease admitted to the College Infirmary, and under the care of my colleague Professor M'Queen. When admitted on the 27th of December the mare was unable to stand owing to weakness of the hind quarters, and the urine was of the usual dark colour. On the evening of the 28th December blood was withdrawn at 7 P.M., and an enumeration with the hæmocytometer gave the number of red corpuscles as 14 million per cubic millimetre. Since the patient had already passed a large quantity of pigmented urine I was prepared to find some diminution in the number of red cells, and I was surprised and to some extent incredulous at the result of my first observation, since that appeared to show that the blood contained nearly double the normal number of red corpuscles.

On the same evening at 10.30 I made a second examination of the blood, and found that the number of red corpuscles per cmm. was now almost the same as before, viz., $13\frac{3}{4}$ millions.

A third enumeration was made on the following evening at 8.30 P.M., by which time the red corpuscles appeared to have fallen to $9\frac{3}{4}$ millions.

After other 24 hours I made a fourth examination and found that the red corpuscles had now fallen to the normal proportion, the enumeration giving the number as $7\frac{3}{4}$ millions per cmm.

I was now less disposed to doubt the accuracy of these results ; (1) because I had in the interval examined the blood of a healthy horse as

a control, and found it to contain about 7 millions of red cells per cmm., and (2) because the apparent decline in the number of red corpuscles, from 14 millions to $7\frac{3}{4}$ millions, was accompanied by a marked improvement in the condition of the animal. The mare subsequently made an excellent recovery, although there was after the attack a slight wasting of the muscles of the quarters.

In the month of March last, through the kindness of Mr Wragg, I was able to examine the blood of another typical case. In this instance I made only one observation, which gave a result in close agreement with that obtained in the first case, the number of red corpuscles per cmm. being $10\frac{3}{4}$ millions. This case proved fatal on the following day.

My next and most recent observations were made during the present month, on a case in the College Infirmary under Professor M'Queen's care. This case, like the preceding two, was a perfectly typical one. The first quantity of urine drawn off after the onset of the attack had been of the usual dark colour, and the mare when brought to the college in a float was quite unable to bear any weight on the hind limbs. The first enumeration of the blood corpuscles was made 40 hours after the seizure, at which time the urine had become much paler, though it still contained some granular pigment. This enumeration gave the number of corpuscles as 11 millions per cmm. As a control, blood from another infirmary patient was examined at the same time, and found to have $8\frac{3}{4}$ millions of corpuscles per cmm. This was a horse in good condition which had been under treatment in the hospital for some time for "capped elbow."

A second examination of the blood was made on the following day (70 hours after the seizure), and this showed the number of red cells to be 10 millions per cmm. The blood of a well-nourished working horse examined at the same time gave a little over 8 millions per cmm.

A third and final examination of the blood was made 94 hours after the attack, and on this occasion the enumeration gave a little over $10\frac{1}{2}$ millions per cmm., while two control horses gave respectively $7\frac{3}{4}$ millions and $5\frac{1}{3}$ millions. The last of these was an unthrifty-looking horse that had been under treatment for 6 weeks with fistulous withers.

During the progress of these observations the case gradually became worse, and by Professor M'Queen's advice the mare was destroyed on the 5th day after the attack.

It will thus be seen that this line of investigation has afforded indications very different from those that were expected. As I had never seen marked signs of anæmia produced by even a bad attack of hæmoglobinuria, I was inclined to adopt the muscular theory of the origin of the pigment, as first propounded by Frohner, and I should therefore not have been surprised to find the number of red corpuscles normal, but I was wholly unprepared to find a notable increase in their number. Should these observations be confirmed, as I confidently expect they will be, the muscular lesions so constant in the disease will have to be regarded as secondary in point of time and importance. It will have to be assumed that the starting point of the disease is an over-production of red corpuscles, uncompensated by a steady destruction of them, such as goes on in health, and that this

over-production is somehow determined by the period of rest and the liberal diet, of which there is a history in nearly every case of hæmoglobinuria.

Before concluding this article I would take the opportunity to refer to a recent contribution to the pathology of this disease which has been made by a French veterinary surgeon, M. Lucet.¹ In France, as in this country, very erroneous notions were until lately held regarding the nature of the disease, the commonest view being that it was a congestion of the spinal cord. M. Lucet proposes to call the disease Paroxysmal Hæmoglobinuria, and he considers that cold is the exciting cause. Curiously enough, M. Lucet regards the disease as an acute nephritis, with consequent toxæmia. Of course, for the acceptance of this theory it is absolutely necessary to maintain that inflammation of the kidneys is a primary and never-failing lesion, and M. Lucet boldly takes this course, and asserts that Fröhner was wrong when he said that renal lesions were not constant. This is, to say the least, a rash statement on the part of M. Lucet. If he had contented himself with stating the number of cases in which he had examined the kidneys, and the time that had elapsed in each of these cases between the onset of the attack and the animal's death, one would have known what weight to attach to his assertion. We know that cases seldom prove fatal within 24 hours, and if any one finds a renal lesion in an animal that has survived for a longer period than that we are by no means bound to admit that this lesion was the initial one. In this connection it is perhaps not out of place to mention that out of 45 cases observed by M. Lucet in 1892-93 only one proved fatal. My own observations have taught me that in the most typical cases of hæmoglobinuria the kidney may be quite free from nephritic changes. Moreover, even if it were admitted that nephritic lesions are constant that would not afford any satisfactory explanation of the most striking feature in the disease, viz., the presence of hæmoglobin, sometimes in enormous amount, in the blood plasma. Congestion and œdema of the kidneys, with extensive shedding of renal epithelium, are not rarely found in animals that have succumbed after some days, but these changes are perfectly accounted for by the hæmoglobinæmia, and indeed are the exact parallel of those met with in experimentally produced hæmocytolysis.

EDITORIAL ARTICLES.

—o— ANTHRAX.

WITHIN the memory of people who are not yet above middle age anthrax or splenic apoplexy was accounted a rare disease in this country. It is unfortunately no longer rare, and there are some grounds for fearing that it is on the increase. Within the past few months especially the reported outbreaks in various parts of the kingdom have been exceptionally numerous, and some degree of

¹ Recueil de Méd. Vét., August 1893.