The Treatment of Insomnia.—WILLIAM BROADBENT (*The Practitioner*, 1906, No. 1, p. 1), in discussing this subject, states that one of the most frequent causes of insomnia is gastrointestinal derangement, especially when associated with flatulence, and while distention of any part of the digestive tract may give rise to dreams and broken rest, or definite wakefulness, it is gastric dyspepsia which gives nightmare, and dilatation of the stomach which produces the most obstinate sleeplessness; consequently, the treatment of insomnia due to these causes should be directed to the etiological factor and all digestive disorders should be rectified. The diet should be simplified and the stomach must never be distended. The amount of bread and of other farinaceous foods, of green vegetables, and of liquid with meals should be restricted, water, hot or cold being drunk every morning. One or two tablespoonfuls of brandy or whiskey in a claret glass of hot water as the only drink at luncheon and dinner will often be useful in dislodging gases. Treatment should be directed to the prevention of gastric or intestinal fermentation, for which purpose salol, naphthol, creosote, carbolic acid or sulphocarbolates may be given, not forgetting calomel or other mercurial, usually a very important consideration. In old persons vascular degeneration may have gone so far that no physiological state of the cerebral circulation conducing to sleep may be attainable; functional derangements, destruction of sleep, may be due to structural changes which cannot be modified; or an idiopathic condition of restlessness may be established; here, after palliatives have been tried, we may give sedatives without hesitation. Late in life there is no danger of inducing a drug habit, and a moderate dose of an opium preparation may conduce to the comfort of the patient and even prolong life. The chloral group is less to be relied upon than opium; the latter may be given nightly with colocynth and hyoscyamus to prevent constipation. In the young, caution is necessary; opium should be avoided and the patient should never be allowed access to sedatives at will. When the sleepless habit has become established sleep may be induced for three or four nights from time to time by one or other of the chloral group with a view toward breaking the habit, or if the patient's self-control is not to be relied upon, paraldehyde may be substituted. Broadbent believes that such drugs as sulphonal, trional, and veronal are doing incalculable harm.

The Physiological Action of Tea as a Beverage.—LAUDER BRUNTON (*Quarterly Journal of Inebriety*, 1906, No. 2, p. 19) sums up a paper with the following statements: Tea when properly prepared and taken in moderation is both useful and agreeable. When taken in too great quantity, or along with butchers' meat, when too strong, when infused too long, or still more when boiled and stewed, it is apt to produce digestive troubles. When taken in excess it may produce nervous symptoms of the most serious character, and facilitate, if it does not actually produce, mental degeneration.

Expectorants.—H. EICHHORST (*Deut. med. Woch.*, 1906, No. 17, p. 649) considers that the irritating cough of tracheitis may be best relieved by 10 drops, three or four times daily, of codeine phosphate $4\frac{1}{2}$ grains and bitter almond water $2\frac{1}{2}$ drams. If the attacks of coughing are associated with nausea or if the appetite is poor he prescribes a mixture

containing codeine phosphate $4\frac{1}{2}$ grains, raspberry syrup 5 drams, phosphoric acid $7\frac{1}{2}$ grains, water 6 ounces; of this the patient takes two teaspoonfuls three times a day. He does not employ narcotics, other than codeine, because of their unpleasant by-effects. These are especially harmful when the bronchial secretion is copious, because of the possibility of producing cyanosis. True expectorants should have a solvent action upon tenacious sputum; in this connection the author recommends the following: potassium iodide 45 grains, simple syrup 5 drams, to which powdered ipecac $7\frac{1}{2}$ grains infused in 6 ounces of water is added; dose 2 drams every two hours. The depressant effects of apomorphine and pilocarpine contraindicate their employment, but a powder consisting of benzoic acid $4\frac{1}{2}$ grains, camphor $\frac{1}{4}$ grain, and powdered sugar 3 grains, may prove useful. In chronic processes with foetid expectoration the balsams are indicated, especially terebinthine and myrtol. In bronchial catarrh of the infectious type benzoesol and benzoguaiaicol are excellent, and the catarrhal inflammation may be lessened by the administration of 20 to 30 drops of the fluidextract of hydrastis four times daily. Steam inhalations also act as an expectorant and in certain instances rhythmic compression of the lower thorax will aid in the expectoration of tenacious mucus.

Marmorek's Serum by the Rectum in Tuberculosis.—M. PETIT (*Revue de Therapeutique*, 1906, No. 10, p. 337) has employed the following technique in the administration of this agent. Every two days, after a preliminary cleansing irrigation. 0.75 of a minim of the serum is injected, after which the patient rests for a time. Of twenty-seven patients so treated, seventeen have been improved and five no longer exhibit physical signs; five have died—all of whom were in the third stage of the disease. Under the treatment the first sign of amelioration is a disappearance of the dyspnoea, which generally takes place after about five injections; later the sputum disappears and finally the physical signs are modified. The general condition remains unchanged. Hemoptysis is not a contraindication, and the author has observed no unfavorable reaction of any kind; in his opinion the treatment seems specific and curative; at any rate it offers no disadvantages. In discussing the paper M. Le Gendre stated that the evidence presented could not, in his opinion, be considered seriously, since the cures or improvements had persisted not more than three or four months, and in consideration of the natural remissions which so often characterize the disease the results hardly constituted a sufficient proof of the efficacy of the treatment.

Thiocol in Non-tuberculous Pulmonary Affections.—M. NIGOUL (*Le Progrès médical*, 1906, No. 14, p. 215) considers that in thiocol we possess a very practical means of administering creosote in such affections as pneumonia, acute and chronic bronchitis, bronchiectasis, etc. The minimum dose, when the object is to drain the bronchi of the secretion, is 30 to 45 grains per day. A smaller dosage will suffice when it is not desirable to relieve the mucous membranes of inflammatory products. In a reported instance of emphysema, daily amounts of from 15 to 25 grains were sufficient to prevent exacerbations of bronchitis and pulmonary congestion. The drug also exercises an advantageous effect upon the digestive system which aids in augmenting the bodily resistance