1913	M. Z. Gr.	n	Gr.	B-R
Sept. 6	14 ^h 7	2	11 ^m 9	0 ^m 0
7	9.0	2	11.9	0.0
8	10.6	2	12.0	+0.1
10	10.7	1	11.8	-0.1
12	11.9	1	12.2	+0.2
18	11.0	—	<12	—
20	11.5	—	<12	—

Bergedorf, 1913 Okt. 31.

1913	M. Z. Gr.	n	Gr.	B-R
Sept. 23	9 ^h 1	2	12 ^m 4	+0 ^m 3
24	9.5	2	12.3	+0.2
24	12.0	2	12.2	+0.1
25	11.2	1	12.7	+0.5
26	8.3	3	11.7	-0.5
27	9.0	2	12.6	+0.4
28	11.1	2	12.5	+0.3

1913	M. Z. Gr.	n	Gr.	B-R
Sept. 29	11 ^h 2	2	12 ^m 5	+0 ^m 2
30	10.1	2	12.8	+0.5
Okt. 1	12.0	2	12.7	+0.4
2	8.5	1	12.9	+0.5
4	9.3	—	12.9	+0.4
21	6.2	—	<13.2	—

K. Graff.

Osservazioni della cometa *Neujmin* (1913c)

fatte all'equatoriale di 38 cm del R. Osservatorio al Collegio Romano. [Seguito, vedi A. N. 4682.]

1913	T. m. Roma C. R.		$\Delta\alpha$	$\Delta\delta$	Cf.	Oss.	α app.	$\log p \cdot \Delta$	δ app.	$\log p \cdot \Delta$	Red. ad l. app.	*	
Sett.	26	7 ^h 37 ^m 31 ^s	+1 ^m 21 ^s 08	+ 4' 8".2	15,3	B	23 ^h 37 ^m 51 ^s 43	9.559 _n	+ 7° 34' 34".8	0.724	+3 ^s 54	+22".4	1
Ott.	5	10 18 1	-1 58.05	- 2 31.3	10,4	B	23 34 0.48	8.667 _n	+10 19 57.6	0.661	+3.47	+23.2	2
	19	7 24 17	-0 25.23	- 6 30.1	10,3	B	23 32 58.91	9.406 _n	+13 25 54.1	0.646	+3.42	+25.0	3
	20	7 13 22	-0 14.80	+ 4 16.8	15,2	M	23 33 9.33	9.426 _n	+13 36 41.0	0.647	+3.41	+25.0	3
	21	7 28 54	-0 55.98	-11 11.9	10,3	B	23 33 21.31	9.363 _n	+13 47 24.8	0.638	+3.40	+25.1	4
	23	7 17 29	-0 26.29	+ 9 4.3	12,3	M	23 33 50.98	9.383 _n	+14 7 41.1	0.635	+3.38	+25.2	4
	24	7 54 12	+1 21.19	-0 35.8	15,3	B	23 34 9.45	9.237 _n	+14 17 41.5	0.620	+3.38	+25.2	5
	28	7 18 9	-0 7.44	+ 3 55.5	12,3	B	23 35 39.67	9.327 _n	+14 54 17.5	0.619	+3.37	+25.7	6

Stelle di confronto.

*	α 1913.0	δ 1913.0	Autorità
1	23 ^h 36 ^m 26 ^s 81	+7° 30' 4".2	Lpz II 11731
2	23 35 55.06	+10 22 5.7	Lpz I 9397
3	23 33 20.72	+13 31 59.2	Kü 10461
4	23 34 13.89	+13 58 11.6	Lpz I 9384
5	23 32 44.88	+14 17 52.1	Lpz I 9377
6	23 35 43.74	+14 49 56.3	Lpz I 9396

Coll'equatoriale di apertura di 38 cm l'astro apparve sempre senza alcuna nebulosità. — Sett. 26. Grandezza 11^m0. — Ott. 5. Gr. 11^m0. — Ott. 19. Gr. 12^m6. — Ott. 20. Gr. 12^m9. — Ott. 21. Gr. 12^m9. — Ott. 23. Gr. 12^m4. — Ott. 24. Gr. 12^m7 (cielo legg. fosco). — Ott. 25. Gr. 12^m8 (B). — Ott. 27. Gr. 13^m2 (B) (cielo legg. fosco). — Ott. 28. Gr. 13^m2. — M = E. Millosevich, B = E. Bianchi. Roma, 1913 Ott. 28. E. Millosevich.

Beobachtungen des Kometen 1913e (Zinner).

1913	M. Ortszeit	$\Delta\alpha$	$\Delta\delta$	Vgl.	α app.	$\log p \cdot \Delta$	δ app.	$\log p \cdot \Delta$	Red. ad l. app.	*	
Auf der Sternwarte des Collegio Romano in Rom von Prof. E. Millosevich.											
Ott. 24	6 ^h 57 ^m 19 ^s	-0 ^m 5 ^s 61	+6' 31".2	12,2	18 ^h 45 ^m 49 ^s 93	9.405	-5° 22' 37".8	0.802	+2 ^s 21	-0".5	1
25	6 11 33	+0 46.73	-1 17.2	20,2	18 50 15.66	9.249	-6 14 29.7	0.811	+2.23	-0.5	2
28	6 22 25	+0 25.66	-3 54.4	20,2	19 4 40.04	9.286	-9 1 19.0	0.826	+2.32	-0.4	3
Ott. 24. Nucleo 10 ^m , piccolissima coda debole, N 70° E. — Ott. 25. Nucleo 10 ^m , nebulosità debolissima, testa r, coda piccola e debole per N 75° E. — Ott. 28. Codetta N 70° E lunga 3', nucleo 9 ^m 5.											

Ott. 24. Nucleo 10^m, piccolissima coda debole, N 70° E. — Ott. 25. Nucleo 10^m, nebulosità debolissima, testa 1', coda piccola e debole per N 75° E. — Ott. 28. Codetta N 70° E lunga 3', nucleo 9^m5.

Am Fadenmikrometer des Kressmann-Refraktors, Heidelberg-Königstuhl von Dr. M. Münder.

Okt. 24	7 23 30	-1 5.72	+1 56.1	20,6	18 45 57.46	9.408	-5 24 13.1	0.847	+2.21	-0.4	4
25	6 36 21	+0 32.24	-11 59.1	20,6	18 50 23.02	9.284	-6 15 59.6	0.854	+2.22	-0.5	5

Am Heliometer der Sternwarte Bamberg von Prof. E. Hartwig.

Okt. 25	6 26 40	+0 50.87	-2 5.6	45,4p	18 50 19.77	9.246	-6 15 18.3	0.858	+2.20	-0.7	2
25	6 50 52	+0 55.08	-3 0.2	45,4p	18 50 23.98	9.325	-6 16 12.9	0.856	+2.20	-0.7	2
27	6 28 36	+1 39.35	-14 11.2	45,4p	18 59 48.05	9.250	-8 5 18.1	0.866	+2.28	-0.5	6
29	6 38 4	-1 49.28	+6 3.8	45,4p	19 9 45.81	9.277	-9 59 25.2	0.872	+2.38	-0.2	7
31	6 45 44	+0 8.08	+3 44.0	45,4p	19 20 12.78	9.298	-11 56 57.0	0.879	+2.44	-0.2	8
Nov. 4	6 42 51	-1 8.66	+6 1.6	45,4p	19 42 30.15	9.274	-15 56 52.6	0.893	+2.61	+0.1	9

Nov. 4. Der Komet wegen Mondschein sehr schwer zu sehen.

Mittlere Örter der Vergleichsterne.

*	α 1913.0	δ 1913.0	Autorität
1	18 ^h 45 ^m 53 ^s 33	-5° 29' 8".5	Strb 6333
2	18 49 26.70	-6 13 12.0	Ott 6426
3	19 4 12.06	-8 57 24.2	Ott 6584
4	18 47 0.97	-5 26 8.9	Strb 6341

*	α 1913.0	δ 1913.0	Autorität
5	18 ^h 49 ^m 48 ^s 56	-6° 4' 0".0	Ott 6433
6	18 58 6.42	-7 51 6.4	Ott 6528
7	19 11 32.71	-10 5 28.8	Cbr M. 6690
8	19 20 2.26	-12 0 40.8	Cbr M. 6774
9	19 43 36.19	-16 2 54.3	Wa 7449

Inhalt zu Nr. 4694. A. A. Michailoff. Zur Theorie der Sonnenfinsternisse. 233. — Mitteilungen über die Kometen 1913 b, c, d, e. 245. — E. Millosevich. Osservazioni della cometa *Neujmin* (1913c). 251. — Beobachtungen des Kometen 1913e (Zinner). 251.

Geschlossen 1913 Nov. 15. Herausgeber H. Kobold. Druck von C. Schaidt. Expedition: Kiel, Moltkestr. 80.