Exhibition of Specimens.

DR. NEVILLE showed the following specimens, giving a description of each:—(a) Sloughing fibro-myoma of uterus; (b) tubo-ovarian cyst; (c) tubal pregnancy, showing decidual cells and hydatidiform degeneration of villi; (d) sarcomatous degeneration of a fibro-myoma.

SECTION OF SURGERY.

President—SIR L. H. ORMSBY, P.R.C.S.I.

Sectional Secretary—J. LENTAIGNE, F.R.C.S.I.

Friday, May 6, 1904.

PROFESSOR E. H. BENNETT in the Chair.

Demonstration in Osteology.

DR. JOHN KNOTT exhibited to the Academy a series of the long bones of the human skeleton, showing the various epiphyses, and pointed out the peculiarities of their ossification and final junction with their respective shafts. The clavicle, which a novice would naturally place among the group of long bones, presented several peculiarities of ossification. There is no epiphysis for the acromial extremity; the centre of ossification for the shaft is the first to appear in foetal life, while the (sternal) epiphysis is usually the last in the skeleton to unite with its shaft. This epiphysis is also distinguished by its peculiar thinness—making it the “battered sixpence” of anatomical similitude, and placing it quite beyond the possibility of the conservative resection of early life. The upper epiphysis of the humerus, which exactly includes the head and tuberosities, has long been familiarly known; but the lower has, till very recently, been always misrepresented. The best English text-books of surgery used to represent an “epiphysary fracture” as including both condyles in the lower fragment. The specimen exhibited showed that the epiphysis was a mere osseous crust, the removal of which hardly diminished the prominence of the condyle. The upper end of the ulna presents a supplemental epiphysis, placed on top of the recognised one, which has remained hitherto undescribed, except by the exhibitor at a former meeting of the Royal Academy of Medicine. The lower epiphysis of the ulna presents no peculiarity, nor does either of those of the radius. The upper end of the femur presents

three epiphyses—for head and each trochanter—approximately in the same stage of fusion with the shaft. That for the head is thinner than is usually thought ; when detached it is but a concave shell of bone. That for the lesser trochanter is very seldom found, its abrupt prominence and small surface of junction causing it to be almost always lost. The epiphysis of the lower end of the femur is of immense importance in connection with growth and development. As its centre of ossification appears immediately before birth, its presence constitutes the best test of the maturity of the fœtus—no other epiphysis of the skeleton developing an osseous centre during intra-uterine life ; and the future gigantism or dwarfism depends more on this epiphysis than on the whole of the rest of the skeleton. Most anatomists place its junction with the shaft at the age of twenty ; this is too early for many cases. Till it fuses with the shaft growth continues, and the height of a giant will be found to depend very largely on the length of the femur. The terminal epiphyses of the tibia, being thin, call for great care on the part of the surgeon in the operation of excision. Those of the fibula present no peculiarity, except the well-known one of the early ossification of the lower one.

Removal of the Cæcum and Ascending Colon.

MR. ALEXANDER BLAYNEY read a paper on this subject, and detailed the notes of a case which was sent to him with symptoms resembling those of recurrent appendicitis. A tumour could be felt somewhat above the appendix region, which proved to be a malignant growth infiltrating the beginning of the ascending colon. The cæcum with a small portion of the ileum and the greater part of the ascending colon were removed. The opening in the colon was closed by a double row of sutures, and the lower end of the ileum was implanted on the transverse colon, a Murphy's button being used to effect the junction. There was some leakage from the colon for a few days. The button was removed from the rectum at the end of a fortnight. The patient recovered rapidly, and was quite well six months afterwards, when the paper was read.