discomfort which such an examination may cause, I have been in the habit of asking a patient whether he really objected to the procedure, and whether he would be willing to undergo it again. If a patient expressed a willingness to undergo another examination, if necessary, or if he stated that he did not have an undue amount of discomfort, it was concluded that the procedure had not been particularly painful. I think that it is going to be possible, however, to utilize intratracheal anesthesia to great advantage in gastroscopy under nitrous oxid.

DR. G. REESE SATTERLEE, New York: I was much pleased to-day to see the general trend of speakers toward embryologic influences as a cause of some of these intestinal conditions. I wish to emphasize again the fact that the water-trap stomach is not an enteroptosis. This condition has not received proper consideration or diagnosis. We found in our own experience that many cases were not recognized at all, or were treated for neurasthenia, hypersensitive stomach, ulcer of the stomach, anemia or some other indefinite condition. Dr. Graves has spoken about the gastropotosis cases. We are not discussing the gastropototic cases. The stomach that we have spoken about is not a gastropototic stomach, although enteroptosis may coexist, but it is distinctly one due to an embryologic condition with the pyloric arm situated high. We do not want to leave the impression that these cases are all operative. The eighteen operative cases referred to were all followed very carefully with the Roentgen ray and at operation, and in every one of them the lesser curvature was below the umbilicus. Often the pancreas was visible above the lesser curvature.

Dr. Mills brought out one very important point, which also bears out the embryologic theory; he says that there is no normal stomach; that the stomach follows the shape of the body. That is what we find. We think that after a while we are going to be able to diagnose these cases without the Roentgen ray, on account of the long, narrow thorax and the narrow upper part of the abdomen.

One more point, in regard to treatment, showing what can be done in these cases medically. I had one woman with a marked water-trap stomach, a long stomach and a long pyloric arm. She was told to use an abdominal belt, to raise the foot of the bed as high as she could with comfort at night, to take advantage of the night's rest and let the stomach and the colon go up toward their proper place and to lie down on the right side of the body after a meal. I saw her about eight months afterward. She said that she did not have the money to get the belt, but that she had raised the foot of the bed and had rested regularly on her right side after a meal. When she had the symptoms of residue in the stomach which at night gave her pain and nausea, she would turn over on the right side and empty the stomach, and the pain and nausea would cease; thus, by this simple procedure, without any belt, without any operation, she had gained 18 pounds, and was doing well.

DR. L. T. LEWALD, New York: In reference to the question of operation not confirming the findings of the Roentgen-ray examination made one or two days previously, I cannot understand how such a discrepancy could occur, except that I have seen a similar apparent discrepancy when the surgeon operates on a hernia, and the hernia has been reduced by a change of position. The point is simply this: As soon as the patient assumes a horizontal position with the stomach empty it may pass to the upper part of the abdomen, where the surgeon may find it. I have seen instances of this sort in which it could be easily drawn down outside the incision, as far as the symphysis pubis, confirming the Roentgen-ray findings as determined by radiographs taken in the vertical position.

DR. R. WALTER MILLS, St. Louis: I am a firm believer in Stillé's views on the subject of the habitus. I believe that its observation constitutes the most important single sign in medicine, considering the universality of its application.

The words enteroptosis, ptosis, atony and dilatation are not used correctly. There should be a perfectly clear and definite understanding with regard to each of these terms, for they mean different conditions.

The findings at operation do not correspond with those found before. One can save himself much difficulty and humiliation by recognizing that this depends chiefly on the fact that owing

to general anesthesia, or the opening of the abdomen, tonus is destroyed. I tried to emphasize the point that the stomach is an organ that has this peculiar attribute of tonus, on which under normal conditions in life its form largely depends. The organs of the cadaver, do not possess it; hence, filling the stomach in cadavers gives misleading results; it does not give us the pictures we find in life. That this same tonus is abolished at operation may be suggested by observations which lead us to conclude that the stomach in life must be contracted for one reason, because when empty it has practically no gas content; at operation this is certainly not the case. Each sacculi and haustra of the intestines will show its individual gas content, if present, in the plate. I repeat that the most important thing in reaching conclusions is a clarity with regard to such points. At the basis of everything is the ability to individualize and realize that there is no normal type of stomach, but that each normal person has a stomach that fits his body, just as his nose usually fits his face.

THE NEW IMMIGRATION AS IT AFFECTS ORTHOPEDIC SURGERY *

WALLACE BLANCHARD, M.D.

Assistant Clinical Professor in Surgery (Orthopedic), Rush Medical College; Orthopedic Surgeon to The Home for Destitute Crippled Children and to The University Hospital
CHICAGO

A larger number of severe rachitis deformities of the legs have been rapidly corrected by bloodless osteoclasis in Chicago than in any half dozen other cities in the world. The records of the Chicago free orthopedic hospitals and dispensaries show that the new immigrant gives them eight times their normal ratio of rachitic deformities, four times their normal ratio of scoliotic deformities, and twice their normal ratio of tuberculous deformities. Chicago is perhaps the only city in the world having winters of ice and snow that harbors twenty thousand new emigrants from tropical or semitropical countries every year, and very bad rachitic deformities are almost peculiar to the children of recent immigrants from hot to cold climates.

The people from Northern Europe huddle closely. Their children double our tuberculous joint deformities. The German and Russian Jew girls are especially prone to scoliosis. Dr. Peter Roberts, in his book¹ presents a map of the United States, with a line drawn from Atlantic City to Cairo, Ill., and thence to the northwest corner of Minnesota, and says that 80 per cent. of the new immigrants remain north and east of those lines, which mark off only one-sixth the area of the country.

In the year 1911 the highest immigration point was reached with one and a quarter millions. The majority of these people flock to our large cities and settle in the poorer quarters. Their sick and injured fill our free hospitals and dispensaries. Statistics show that in the year 1911 Chicago held eighty-four thousand new immigrants from across the Atlantic Ocean, and it is estimated that one in every five or six comes from a hot country.

The records of three thousand orthopedic cases treated in Chicago hospitals and dispensaries show some interesting points as to the susceptibility of different nationalities to different deformities.

* Read in the Section on Orthopedic Surgery of the American Medical Association, at the Sixty-Fourth Annual Session, held at Minneapolis, June, 1913.

¹ Because of lack of space, this article is abbreviated in THE JOURNAL. The complete article appears in the Transactions of the Section and in the author's reprints.

1. Roberts, Peter: The New Immigration, The Macmillan Co., 1912.

Approximately, one-third of all the cases treated were tuberculous deformities, one-third were rachitic deformities, and about one-third were congenital and acquired other than rachitic, and included the infantile paralysis deformities.

English, Irish, Scandinavians and Poles show an average of about 48 per cent., and the Germans 35 per cent. Occasional cases of badly twisted rachitic, so-called "grape-vine" legs are seen in the children of colored people from the far South. Notwithstanding the

Fig. 1 (Case 1).—Lizzie H., aged 4. Bow-legs. The extreme deformity made locomotion difficult.

Fig. 2.—Roentgenogram of Case 1 after rapid bloodless osteoclasis and overcorrection at one point only.

Fig. 3 (Case 1).—Lizzie H., six weeks after osteoclasis, and showing that overcorrection has neutralized the other outbends, producing symmetric legs.

Fig. 4 (Case 2).—Emitt F. Bow-legs, with an inward twist in the shaft of the left tibia and a toeing-in of left foot.

Fig. 5.—Roentgenogram of Case 2 after osteoclasis. Shows overcorrection at apices of deformity bends and rotation out of lower fragment of left tibia.

Fig. 6 (Case 2).—Emitt F., six weeks after rapid bloodless osteoclasis, and showing symmetric and useful legs.

My statistical findings may not be exact, but they vary only slightly from those previously presented by Dr. F. C. Test.² Of all the patients classed as American, 42 per cent. present tuberculous deformities, while the

susceptibility of the colored race to pulmonary tuberculosis, only 17 per cent. of joint tuberculosis is shown. The Italians, Greeks, Syrians and other tropical people are almost exempt from tuberculous joint disease, with a showing of only 4 per cent. The Americans lead slightly

2. Test, F. C.: *Chicago Med. Recorder*, February, 1904.

in congenital deformities, with 12 per cent. The German and Russian Jewish girls develop scoliosis in large numbers. They make up 66 per cent. of all the recorded cases. The most surprising record is given by the Italians, Greeks, Syrians and other tropical people, with 80 per cent. of the rachitic deformities.

Why the new immigrant gives us nearly half of our free treated tuberculous joints, two-thirds of our scoliosis and a great majority of our bad rachitic deformities and how the emergency is met will be briefly considered.

The people from northern Europe largely seek work in our clothing and other factories and do piece work at home. These are essentially indoor people, and their children furnish a large share of the cases of tuberculous joint disease. At the Home for Destitute Crippled Children, Chicago, all the acute tuberculous joint cases are treated in an open roof-garden and the convalescent patients are sent to the Country Home, which is largely an open-air structure on a hill, twenty miles from the city. Five years ago the Home had a long line of then

conditions of climate, food and habits had the effect of producing rachitic children. Curiously enough, the children born abroad did not develop rickets. Even babies in arms, but a month old at the time of leaving their old home, escaped deformity. Only the Chicago-born children were attacked, and they suffered so excessively that at one time 80 per cent. had rickets. The rickets was of a severe type, producing bow-legs and knock-knees of great deformity, and also in many cases distortions of the chest, deformed spines, rachitic coxa vara, and curvature of the clavicles and bones of the arms. This condition prevailed for several years. Then, as these people became adapted to the climate and diet of their new home, the rickets began to abate, and in about seven years from its onset it was practically extinct in the colony.

This is a typical rachitic history that is being constantly repeated and amplified.

The unprecedented accumulation of severe rachitic deformities of the legs forced the Chicago orthopedists

Fig. 7 (Case 3).—Ada S. Knock-knees, so severe as to make locomotion difficult.

Fig. 8.—Roentgenogram of Case 3 after supracondylar bloodless osteoclasis and redressment.

Fig. 9 (Case 3).—Ada S., six weeks after bloodless osteoclasis, showing the knock-knees corrected.

incurable chronic sinuses of tuberculous joint disease. My paper³ on "The Passing of Bismuth Paste," shows that I was the first to report any large number of old sinuses of tuberculous joint disease cured by sinus flooding.⁴ The Home cases soon demonstrated the dangers of bismuth and of flooding acute sinuses.

That 66 per cent. of all the cases of scoliosis should be in Jew girls is easily explained. The Jewish mother in a strange land fears to let her young daughter out of sight. The girl sews early and late on shop work, until a vicious posture becomes fixed.

The Abbott treatment for scoliosis has been used for one year and seems to promise comparatively quick and satisfactory results.

A remarkable history is presented by a Syrian colony that was left stranded in Chicago at the close of the World's Fair in 1893. In Palestine they had known nothing of rickets. But a few months of the changed

to get away from the slow and uncertain methods of mechanical devices and repeated osteotomies. So it comes that I am able to report on nearly two thousand bad rachitis bow-legs, knock-knees and anterior bent tibias straightened by rapid osteoclasis; that is, by eight-second bloodless correction in the osteoclast, eight minutes for redressment of the deformity and application of the plaster of Paris, five weeks in plaster, and walking in six weeks on symmetric legs.

The large cities of Europe have nearly stationary inhabitants and none have the conditions that produce bad rachitic deformities. I recently spent six months in the large orthopedic clinics of England, Germany, Austria, Italy and France, and did not see altogether as many bad rachitic deformities as I might have seen at my own clinic in a week.

Professor Adolf Lorenz, of Vienna, strongly favors osteoclasis, though his osteoclast is not to be compared for accuracy with the Grattan. The Grattan osteoclast gives nearly absolute precision in the breaking-point, and

3. Blanchard, Wallace: *Med. Rec.*, New York, May 18, 1912.

4. Blanchard, Wallace: *Am. Jour. Orthop. Surg.*, August, 1908.

the pressure-bar may be moved considerably in either direction from the center without affecting the rule. The term "rapid osteoclasia" is used because osteoclasia that is not rapid is always a failure and brings the osteoclast into disrepute.

Fig. 10 (Case 5).—Lucy V., with a combination of anterior bent tibias and knock-knees, making "grape-vine legs."

Fig. 11 (Case 5).—Lucy V., after rapid bloodless osteoclasia, showing symmetric legs.

After osteoclasia the bone unites firmly in half the time taken after an osteotomy. Bones bend more or less in the osteoclast before breaking, and the bending contributes toward symmetry of the leg, while osteotomy leaves only an angular correction. I have frequently demonstrated in the correction of bad anterior deformities of the tibia that cuneiform osteotomy on a child 6 to 10 years of age shortens the leg about 1 inch, while osteoclasia of a similar leg lengthens the leg about 1 inch. Two inches difference in height is of great importance to a child already dwarfed by rickets. When the tibia is to be lengthened considerably, the Achilles tendon must be tenotomized as a preliminary step, to allow it to lengthen correspondingly.

In bow-leg with several contributing curves, the apex of the most prominence is usually selected for the operative point and overcorrection is made not only sufficient to correct the curve attacked, but continued sufficiently to neutralize the two minor deformity curves, so that when the plaster of Paris sets it looks like a case of knock-knees. In the same manner, when a supracondylar osteoclasia has been done for the correction of knock-knees, the legs are held overcorrected and when the plaster has set it looks like bow-legs. An extended experience with rapid osteoclasia has demonstrated better results in six weeks than can be usually obtained in from six months to three years by the older methods.

Osteoclasia should not be done on rachitic bone until it has eburnated. Eburnation represents entirely different characteristics from the slow process of ossification. Normal bones of young children are pliable until ossification, while rachitic bones are friable until eburnation. Eburnated bone is strong, brittle and unyielding. Eburnation may be delayed by disease or insufficient nourish-

ment for weeks or years, but with favoring conditions of fresh air, raw and fatty foods it is quite certain to come, and when it does, it comes suddenly, perhaps in a month, and possibly in a week. Normal bone of a child of 4 years will bend a long way under pressure of the osteoclast before it goes to an osteokampsis or partial fracture, while eburnated bone breaks hard and complete like cast iron. These rules of eburnation cannot be found in either German or English literature, but they are founded on practical experience with several thousand cases.

The mild, non-rachitic bow-legs of small children that are frequently corrected in the course of normal development have nothing in common with the cases reported in this paper.

The emigrant woman has nearly twice as many children as the American-born woman, statistics showing 4.7 against 2.9 per cent. Bad rachitic deformities, tuberculous joints and scoliosis in the young of these new emigrants means a large element of helpless pauperism for the future. Congress has taken steps toward the establishment of a government station in Chicago for the temporary care and protection of immigrants. If the government and the community could be roused to the necessity of caring for the welfare of the immigrant mother for the first year or two in her new home,

it would be a great uplift for the future American citizen.

15 East Washington Street.

Fig. 12.—The Grattan osteoclast in position to fracture the tibia and fibula, with the pressure bar against the outside of the leg for correction of bow-leg or anterior bent tibia.

ABSTRACT OF DISCUSSION

DR. REGINALD H. SAYRE, New York: There is no question that the changes in the environment, mode of life and diet of a race have a great bearing on their nutrition. Some years ago a number of articles appeared in the medical journals

stating that rickets was unknown in Japan and that if we were all to live on rice, as the Japanese do, we should not have rickets. I notice now, however, that it is being discovered that the Japanese have rickets. How much of this is to be attributed to the great strain that has been put on that nation as the result of the war with Russia, and the consequent impoverishment and reduction in the diet of its people, may be a subject for consideration. There is no question that since immigration from Europe to this country has increased the number of cases of rickets has vastly increased. In New York it was a rare thing to find a case of rickets when I first began the practice of medicine, while at the present time deformities of all sorts caused by rickets are frequently encountered.

The Italians and negroes furnish the greatest number of cases of this disease. It seems to me that I saw a great many cases of rickets in Bologna. In fact, the orthopedic hospital there is called the Rachitic Institution, showing that rickets has furnished a large quota of the patients. I understood, however, from Dr. Blanchard's paper that there were, apparently, few cases of that kind there when he went through the institution; when I was there they had rickets of all kinds and descriptions.

Fig. 13.—The Grattan osteoclast in position to fracture the femur close to the condyles for the correction of knock-knee.

I agree with Dr. Blanchard that the Grattan instrument is a useful osteoclast. I do not understand, however, what he means by "rapid osteoclasis," because I think that, if you exert the force for more than a short time you will damage and lacerate the tissues extensively. All the bone-breaking machines with which I am familiar are based on the principle of a rapidly exerted pressure, rather than a slow, squeezing force, which would certainly do great damage.

In working close to a joint, I believe that you can do better with an osteotome than with an osteoclast, because it is necessary to get a certain amount of leverage in an osteoclast. It seems to me that it is better to make the correction at the point of the deformity rather than at an intermediate point.

DR. WALLACE BLANCHARD, Chicago: I have spoken of "rapid osteoclasis" in this paper because in my earlier observation of work done with the osteoclast, I found that the pressure was exerted slowly and that the instrument did a great deal of damage simply for that reason. Therefore, I always want to make a point of the fact that eight seconds is the longest time that pressure with an osteoclast should be exerted.

LOCAL AND SPINAL ANESTHESIA IN GYNECOLOGY AND OBSTETRICS *

GEORGE GELLHORN, M.D.

ST. LOUIS

That there is in surgery a field for anesthetics other than by way of the ordinary ether drop-method is so indisputable a premise that I need not waste the few moments at my disposal with argumentation. Suffice it to say that there are no statistics in existence which give a true picture of the alleged harmlessness of the open ether-inhalation narcosis. The exact number of fatalities due to ether will never be known, nor has the rôle of ether in the causation of postoperative complications which endanger life and health of the patients ever been exhaustively considered. Few men have the courage to publish their failures, and the medical journals, as Sellheim remarks, "serve as a medium for recounting accidental success rather than the reverse."

Of the large number of new modes of anesthesia and modifications of old ones, gynecology has been particularly benefited by the two methods which I wish to discuss more in detail—local and spinal anesthesia.

We know both from clinical observations and anatomic researches that the cervix uteri is but meagerly supplied with sensory nerve-endings. Scarifications of the vaginal portions or excisions of small pieces for microscopic examination can usually be done without causing any pain. Under these circumstances it seems to be somewhat out of proportion to employ the heavy artillery of an ether-inhalation narcosis when the injection into the cervix of a small quantity of cocaine, novocain, alypin or other drugs will produce an analgesia sufficient for the performance of most of the minor gynecologic operations on the uterus. There is, of course, the objection that the patient retains her consciousness and may therefore suffer a psychic trauma. This objection is largely theoretical. It is undoubtedly a difficult matter to break away from long-established habits. If local and spinal anesthesia—for the same objection has been raised against the latter method and may as well be anticipated here—if, I repeat, local and spinal anesthesia had by chance been discovered earlier than ether narcosis, we may well assume that most of us would look on a state of unconsciousness as a disquieting and questionable innovation.

We must further bear in mind that, though the patient is conscious during local or spinal anesthesia, her sensibilities are not nearly so acute as those of the operator or the audience. Any anesthesia should begin at least a day prior to the operation. Excessively nervous patients should receive bromids, valerian or other sedatives as soon as they enter the hospital. Veronal should be given on the evening preceding the operation to insure sound sleep, and hypodermic injection of morphin-atropin, morphin-scopolamin, or the like, should be administered one-half hour before the time set for operation. Before the patient enters the operating-room, the eyes should be covered with a mask, and the ears filled with cotton saturated with olive-oil. By this preparation the patient's apprehension is dulled, and, if there is no sensation of pain during the operation, there can be no psychic shock. A single trial of these methods will disarm even the most profound skeptic. After all, preparations of

* Because of lack of space, this article is abbreviated in THE JOURNAL. The complete article appears in the Transactions of the Section and in the author's reprints.

* Read in the Section on Obstetrics, Gynecology and Abdominal Surgery of the American Medical Association, at the Sixty-Fourth Annual Session, held at Minneapolis, June, 1913.