injection. The injections must be made with all aseptic precautions. After the injection an alcohol dressing is advisable in order to counteract any possible infection. I begin with an injection of 0.25 c.c. and gradually increase the strength to 1 c.c. daily (0.01 gm. sodium arsenite). In one case of pernicious anemia I injected 2 c.c. (0.02 gm. sodium arsenite) every second day, without causing any toxic symptoms. Clinically the results with the modified solutions were very gratifying. The injections are made on the upper arm or between the shoulder blades.

Despite the favorable reports in regard to the treatment of certain skin affections and syphilis by the hypodermic use of atoxyl and the cacodylates, which contain, respectively, 37.7 per cent., 35 per cent. of pure arsenic, these organic and less poisonous combinations of arsenic have not proved nearly as efficacious as the conventional inorganic arsenic preparations in the more aggravated skin lesions, and especially in Hodgkin's disease and pernicious anemia. Another objection to the hypodermic use of the organic arsenic combinations in these diseases is found in the fact that the dose has to be a large one to be of any therapeutic use, and this circumstance frequently has caused severe arsenic intoxications with serious consequences, as the literature of the last year goes to show.

224 West Twenty-fourth Street.

AN UNUSUAL COMPLICATION OF THE OPERATION FOR CONGENITAL DISLOCATION OF THE HIP

LEONARD W. ELY, M.D.

NEW YORK

The following case seems interesting on account of a rather infrequent complication:

Patient.—O. A., a girl of 8, was brought to the Roosevelt Hospital on Oct. 23, 1908, on account of a peculiar gait which had always been present since infancy. She had never suffered pain.

Examination.—Fairly well-nourished child, walking with a decided waddle. Heads of both femurs were dislocated, the right being fairly well developed, the left not so well. Well-marked lumbar lordosis was present. Length of right leg, 53.5 cm.; length of left leg, 53 cm.

Operation.—About November 23 the child entered the hospital and shortly thereafter was put to bed with extension of thighs at a right angle to her trunk. On December 7, under ether anesthesia, the Lorenz operation was done on both hips. The adductors were broken down rather easily, and reposition was accomplished on both sides without difficulty. The stability on the right side was rather poor, but on the left good. The head on this side was prominent and in its normal place under the femoral artery. In order to prevent relaxation, the thighs were held in extreme abduction while the dressing was put on, abduction being carried well behind the plane of the body. During the application of the plaster, the house surgeon, who was holding the legs, called attention to the condition of the left leg and foot. They were bluish-red and pulseless. This was considered to be due to the constriction of the dressings at the knee, and they were loosened at that point. As the circulation in the limb had not improved, however, by the time the plaster had been applied, the cause of the difficulty was sought. No pulsation could be felt in the popliteal vessels, nor at any point below the plaster, and the foot was becoming darker and rather cold. Next the plaster was cut away over the head of the femur to relieve pressure there. The femoral artery did not pulsate. Then all the plaster was removed from the af-

fected limb, still without producing any improvement whatever. Not until the hip was redislocated did the circulation start up.

In describing a case of gangrene after reduction of one of these hips Lorenz was wont to ascribe it to a compression of the femoral artery between the plaster and the prominent head of the femur, and this explanation has always been accepted as the correct one, as it certainly would be the most obvious. In this case, at least, if the plaster had been left on and any accident had occurred, we probably should have considered it due to this cause. In view of the facts, however, some other theory must be entertained. Evidently the dressing had nothing to do with it. We are inclined to think that the head of the superabducted femur (that is, superabducted from a position of right-angled flexion) compressed the femoral artery against Poupart's ligament and thus shut off the blood supply.

Whatever theory is entertained, this case demonstrates the necessity of watching carefully the circulation in a leg after the reduction of a congenital dislocation, both at the time of operation and afterward.

THE POSSIBILITIES OF INFECTION FROM TABLE UTENSILS AT SANATORIA*

J. WOODS PRICE, M.D.

SARANAC LAKE, N. Y.

In the following experiments forks, spoons, teacups and milk glasses in constant use at the Reception Hospital, Saranac Lake, were used. This institution receives patients suffering with pulmonary tuberculosis, a large majority of whom are in an advanced stage of the disease, and some of whom are in the last stages.

The experiments were made without the previous knowledge of the nurses or maids of the institution, and several months apart, to avoid the possibility of the dishes receiving an unusually good washing.

For the first experiment thirty forks and twenty-five each of spoons, cups and tumblers were selected immediately after breakfast, and after ordinary good washing but not sterilization, the *modus operandi* of which was as follows: The utensils were first thoroughly cleansed with a mop or cloth, with soap and water as hot as the maid's hand's could stand, and then rinsed in running water as hot as it could be delivered through a pipe leading from a tank containing boiling or almost boiling water, and then dried with a cloth.

In order to obtain the material for the experiment the articles selected were, after the usual washing, well rubbed with a sterile 0.5 per cent. sodium bicarbonate solution by means of sterile cotton swabs on glass rods, and these washings used for the inoculation of guinea-pigs. The washings were taken at once to the laboratory and distributed among fifty-six guinea-pigs by subcutaneous injection into the right groin, 0.75 c.c. to each pig, and with the use of sterile syringes. The distribution was as follows:

Washings from forks, 14 pigs
Washings from spoons, 14 pigs
Washings from cups, 8 pigs
Washings from tumblers, 8 pigs
Mixed washings of cups and glasses, 6 pigs.
Mixed washings of forks and spoons, 6 pigs.

One animal died on the second day after the inoculation, and thirteen others from time to time up to the forty-first day, when all living animals were killed. An

* Read before the International Congress on Tuberculosis, Washington, D. C., Oct. 1, 1908.

autopsy was performed on each animal at the time of its death or when killed, and all were found to be free of tuberculous infection. Smears were made on slides from the spleens and right inguinal glands of all those pigs killed on the forty-first day, and examined microscopically for tubercle bacilli, and found to be negative in every instance, nor could tubercle bacilli be demonstrated in stained specimens of the centrifugalized washings before inoculation.

In the second experiment five articles of each sort were selected immediately after use and before washing of any kind, and swabbed with the 0.5 per cent. sodium bicarbonate solution as described above. These washings, which were very cloudy with particles of food, were immediately on arrival at the laboratory injected into eight guinea-pigs in the same manner as in the first experiment, the washings of each set of articles being divided between two pigs.

All these animals lived until the forty-first day, when they were killed and autopsied, and all, excepting the two inoculated with the washings from the milk glasses, were found to be tuberculous. It was an easy matter to demonstrate tubercle bacilli on slides made from the glands and spleens of these pigs, though no tubercle bacilli had been found in a casual search of smears made from the washings previous to inoculation.

The third experiment was a repetition of the first, but with the use of only six each of the utensils and twelve pigs, the washings from the different articles being distributed among three pigs. This test was made eight months later, when the dish-washing was being done by a new set of maids, but under the supervision of the same corps of nurses. The result, however, was the same.

It is fully realized that two experiments are not sufficient for definite conclusions, and that sterilization of eating utensils by heat or boiling, when possible, is not to be discouraged; at the same time when the dish-washing is thorough the danger of infection from this source does not seem to be very great.

My thanks are due to the Saranac Laboratory and the Reception Hospital for facilities placed at my disposal for the experiment, and to Dr. Baldwin, who suggested it.

121 Main Street.

CONGENITAL INSUFFICIENCY OF THE PALATE*

G. W. STIMSON, A.B., M.D.

ATLANTIC CITY, N. J.

My attention was first directed to this condition by Makuen of Philadelphia, who exhibited a similar case before the College of Physicians and Surgeons about a year ago. As far as I have been able to discover, there has been no reference made to it in medical literature up to the present time.

Patient.—L. C., white, female, single, aged 18, referred to me by Dr. Mary Townsend, of this city, consulted me Oct. 22, 1908. Practically her entire complaint was a marked defect in speech, which she said had existed ever since she began to talk; it was that alone which prompted her to seek medical advice. There was slight nasal obstruction and some post-nasal "dropping," but these were not troublesome enough to cause any complaint in themselves. There was no history of

diphtheria or other serious illness; in fact, with the exception of measles at the age of 14, which clearly had had no effect on the condition under consideration, the patient had always enjoyed good health. The family history was negative.

Examination.—The palatal reflex was absent, and the free edge of the soft palate hung so far away from the posterior pharyngeal wall that a postnasal examination could be made with a large laryngeal mirror with perfect ease. Sensation was unimpaired. Occasionally small amounts of tea and coffee come through the nose, but apparently not often enough to be troublesome. The following points were further evidence of faulty development: The uvula was short and bifurcated. The palate, while not extremely high, was undoubtedly more highly arched than the average normal one. On palpation with the finger in the mouth, or on inspection while lifting the velum upward and backward with a blunt probe, it was seen that the posterior edge of the hard palate at the point of its attachment to the soft one, presented the shape of a long narrow letter V with the apex located well forward, and that the soft palate was abnormally short and not of sufficient length to reach the posterior pharyngeal wall. Incidentally, there was a hypertrophic rhinitis with a marked hypertrophy of the tips of the lower turbinates posteriorly, with mucoid degeneration of the mucous membrane covering them.

In normal voice-production when any sounds except those of *m*, *n*, and *ng* are spoken, the soft palate is made tense by the tensor palati muscle and lifted upward and backward by the levator palati until it comes tightly in contact with the posterior pharyngeal wall, thereby shutting off all communication with the nasal cavity and causing the oral cavity to be used as the chief resonating-chamber. But, when the sounds *m*, *n* and *ng* are produced, the soft palate hangs uncontracted, the communication with the nasal cavity remains intact and the latter plays the chief rôle as a resonating chamber.

In this case, so far as results are concerned, the palate is constantly in a state comparable to that of a normal individual during the production of these three sounds; that is, it is "insufficient" to shut out the nasal cavity as it should in the production of all other sounds, thereby allowing it to participate strongly as a resonating chamber not only in the production of the sounds of *m*, *n* and *ng* as it should under normal conditions, and which, as would naturally be expected, she can say as plainly as anyone, but also in the production of all other sounds as well, a distinctly abnormal condition, thus imparting a strong nasal "twang" to her speech and causing her to talk through her nose.

In the development of the normal embryo, at about the fifth week the first visceral arch divides into an upper part, the maxillary arch, and a lower portion, the mandibular arch. The maxillary arches of the two sides unite at their anterior ends with the intervening nasofrontal process to form the upper boundary of the buccal cavity; the mandibulars to form the inferior boundary. In the fetus the palate process of the superior maxilla advances toward the median line until it meets and unites with its fellow of the opposite side. The horizontal plate of the palate bone develops similarly and very shortly after and thus is produced the hard palate. The two halves of the hard palate unite first in front, their union being completed about the twelfth week. Ordinarily it is stated that if union is incomplete the anomaly of cleft palate results.

In this case while union has apparently been practically complete and the patient has been fortunate enough to escape the greater evil of cleft palate, still development has not gone on to the perfection of completion (as is evidenced by the V-shaped posterior edge and the high arch of the hard palate, the short soft pal-

* Read before the Atlantic County Medical Society, Dec. 4, 1908.