ON A CASE OF CHROMIDROSIS.

By ARTHUR WYNNE FOOT, M.D., UNIV. DUBL.;

Senior Physician to the Meath Hospital.

[Read in the Section of Medicine, December 14, 1888.]

AFTER the exhibition of several characteristic coloured drawings of patients with chromidrosis, Dr. Foot remarked :—

In tracing the history of the disease, from the first case recorded by Dr. Yonge, of Plymouth, in 1709, up to the present year (1888), I find that there are forty-six admittedly genuine cases on record—cases which have passed repeatedly through the crucible of criticism, and to which no exception can be taken. Of these forty-six cases six have occurred in men, and forty in women—for the most part young and unmarried. That so small a number as forty-six cases should have been reported in the 180 years since Dr. Yonge first drew attention to it, I consider to be a proof of the rarity of its occurrence. That it is a curious and striking affection is shown by the amount of incredulity which has often been exhibited in regard to it. As recently as the year 1861 the Medical Society of the Hospitals of Paris gave an open verdict on the subject, notwithstanding the strong pleading of M. Hardy in its favour; and in the same year the Imperial Academy of Medicine declined to allow de Méricourt's essay on chromidrosis to be published in their memoirs, on the ground that the authenticity of the cases was doubtful, although the Committee which they had appointed to report on it recommended it merely for publication, without asking for the expression of any opinion on its merits. On April 4th, 1882, a Committee of the Clinical Society of London reported "that chromidrosis has an undoubted existence as a genuine affection."

I am not inclined to follow Féréol in substituting the name

chromocrinia for chromidrosis ; for if the views of Meissner and Unna are correct, that the coil of the sweat gland secretes fat and the end of the duct sweat, disorder of the coil glands would account for all the phenomena.

I am an advocate of the theory that the pigment is due to the oxidation of indican eliminated by the cutaneous glands : its preference for the face may be explained by the free exposure of that area to the great oxidising influences of air, light, and heat. Some ferment also might operate in effecting the conversion of soluble colourless indican into indigo-blue or brown. The possibility of a bacterial origin is not to be overlooked, as the source of the colour in "blue milk" is often the *B. cyanogenus* and in red sweating *Bacterium prodigiosum*. The indican, I consider, is derived from the absorption from the intestinal canal of small quantities of indol. The frequency of constipation in chromidrosis, the sudden disappearance of the colour after purgation, its diminution after regulation of the bowels, and the general aversion to animal food these cases exhibit, are facts which make in the direction of this view. The proportion of indican in the urine increases, *pari passu*, with the amount of exudation on the face, and *vice versâ*. Hoffmann has detected indigo in the bluish-black exudation, and after reducing the pigment to indigo-white it was seen to regain its colour by exposure to the air.