THE LARYNGOSCOPE.

VOL. XXVIII.

ST. LOUIS, NOVEMBER, 1918.

No. 11.

ORIGINAL COMMUNICATIONS.

(Original Communications are received with the understanding
that they are contributed exclusively to THE LARYNGOSCOPE.)

THE PATHOGENESIS OF BRONCHIAL ASTHMA.*

DR. WOLFF FREUDENTHAL, New York.

A. *The Bronchial Attack.* While studying cases of so-called spasmodic or bronchial asthma the question came up in my mind: Is spasmodic asthma in fact a spasm or what is it? The general belief is that an attack of asthma is characterized by a constriction of the bronchioli, lasting for hours or even days. Many theories have been advanced in support of this belief and it seems perfectly plausible to accept the nervus vagus as the main culprit and to explain every symptom by the action of that nerve.

Although I cannot prove that this is not so, I believe that an attack is brought about, not by a spasm of the constrictor fibers coming from the vagus, but rather by a paralysis of the dilator fibers and that this accords with the description given by me of the etiology of asthma in general.

The first thing that strikes the observer is the unusual occurrence of a spasm lasting for such a long time, as it so often does. Is it not easier and more in accord with other pathological facts to explain that attack as the result of a paresis or paralysis? There are similar conditions in the larynx, if you will remember some neurological problems of that region, as for example the bilateral abductor paralysis.

*Read before the American Laryngological, Rhinological and Otological Society at its Twenty-fourth Annual Meeting, held at Atlantic City, May 29 and 30, 1918.

If we suppose that the asthmatic paroxysm sets in with a paralysis of the dilator fibers we have to assume that a constriction of the bronchioli must follow immediately in the same manner as in facial paralysis, where the face is drawn to the non-affected side. In other words, as soon as the inhibitory or bronchodilator fibres of the vagus that supply the bronchial musculature, are paralyzed, a tonus of the opponents sets in, i. e., a condition of persistent activity of the constrictor fibres is established. In the horse it has been proven that these constrictor fibres of the bronchi are in a state of constant or tonic excitation. Yet, according to Luciani,¹ the physiological function of the smooth muscles of the bronchi and of the constrictor and dilator nerves, which serve them, is not yet fully explained. What we do know positively is the existence of constrictor fibres, as proven by Einthoven and Baer, and the existence of broncho-dilator fibres running side by side with the former, as demonstrated by Dixon and Brodie.² We see, therefore, that the vagus may be held responsible in either case.

But there is another system that is of great importance; namely, the sympathetic. The ramifications from one system to the other are numerous and there may be disturbance of the sympathetic when the symptoms incline towards an irritation of the vagus and vice versa. While the action of these nerves is not yet clearly established, there are other symptoms apparently in favor of my theory that we have to deal here with an atony or paresis or paralysis of the broncho-dilators and not with a spasm of the constrictors.

If we take, for example, the abdominal muscles, i. e., the muscles that are concerned in expiration, we find that they are "quite flaccid" (Starling) even during expiration. That could never happen if a spasm of the expiratory muscles had been present.

Another point is the frequent development of pulmonary emphysema in such cases. According to Luciani pulmonary emphysema is promoted by atony or paresis or paralysis of the smooth muscles of the bronchi. An atony never means spasm.

Nor is it necessary to demonstrate a spasm of the inspiratory muscles. For at the beginning of an attack the asthmatic patient has difficulty only in expiration, which means that through a paralysis of the dilatory muscles the others gain the mastery and contract. You will find a corresponding condition in facial palsy. The muscles of one side become paralyzed and immediately their antagonists on the other side get into action and contract.

Later on during an attack, when the lungs become overdistended, there is difficulty in inspiration as well. But it was found by Walker³ on fluoroscopic examination that the lungs do not expand and that the diaphragm remains stationary. This is an important finding, as the diaphragm under normal conditions is the main factor in inspiration, a factor that is eliminated entirely through an atony or paresis or paralysis in an acute attack. A spasm of the inspiratory muscles may, therefore, also be excluded with certainty.

Furthermore, the action of adrenalin and other drugs has to be considered. Laryngologists and other clinicians, seeing the almost immediate effect adrenalin gives when either injected subcutaneously or applied endobronchially in an acute attack of asthma, were of the opinion that it caused a dilatation of the bronchioles. Golla and Symes, however, found that adrenalin usually produces unmistakable constriction of the bronchioles. That was the case in the rabbit and cat. Undoubtedly it holds good for man as well. But when the bronchioles were initially constricted by other drugs, then Golla and Symes⁵ saw the normal effect of adrenalin reversed and bronchial dilatation produced. This fact may be as surprising to others as it has been to me, but nevertheless it remains a fact.

Besides, adrenalin acts only on those tissues which are innervated by the sympathetic nervous system. This again speaks against the activity of the vagus and in favor of the sympathetic. It must be admitted, however, that the vagus may carry some fibres from the sympathetic, as mentioned before, a circumstance that makes this question still more complex.

The action of other drugs on the bronchial musculature has been studied very carefully by D. E. Jackson⁶ of St. Louis. Some of them produce immediately and most profoundly a broncho-constriction when injected intravenously, as arecoline, for instance. Death may easily follow from absolute failure of air to enter the lungs. Other drugs tested by Jackson, like hordenine, have the opposite effect, i. e., hordenine is fully as effective as epinephrin in producing broncho-dilatation. This also occurs after constriction has been produced by histamine. It is to be regretted that owing to war conditions most of these drugs cannot be procured at present.

Returning to the relationship between the vagus and sympathetic we know that in the function of respiration both the respiratory centre and the autonomous nervous system participate, so that by a sort of self-regulation the vagus acts like an antagonist to the automatism of the center. The action of these two systems, when

in a state of equilibrium, according to Eppinger and Hess,⁷ can be compared to a pair of evenly balanced scales, containing equal weights, as represented by atropin on one side and adrenalin on the other.

And it may be concluded that paralysis of the sympathetic and stimulation of the vagus show the same symptoms and are closely related, not only symptomatically, but also etiologically. As long as this fact stands proven by Eppinger and Hess, so long are we justified in speaking of the asthmatic paroxysm as due either to an irritation of the vagus or to a paralysis of the sympathetic. For reasons stated above, the writer is inclined towards the latter theory; to which one more point may be added here.

We have referred repeatedly to the bronchial musculature or that of the bronchioli or alveoli. But it should not be forgotten that the essential distensible air spaces of the lung are the infundibula or the terminals of the bronchioli. However, as the infundibular air is regulated by the musculature of the bronchioli, we were right in mainly considering these. The asthmatic sufferer involuntarily endeavors to keep his infundibula filled; that is, to keep his lungs in an inspiratory position. In that way he reinforces the elastic powers which dilate the bronchioles and thus aid expiration. But since these elastic powers are out of function through some kind of paresis, no deep inspiration can be taken, nor expiration aided in that way.

In conclusion it has to be acknowledged that the constrictor muscles of the bronchioli during an asthmatic attack are not in a spasmodic condition, as heretofore accepted, but that they are in a state of artificial tonus brought about by the paresis of the inspiratory muscles.

These are the writer's views concerning the acute asthmatic attack. The asthmatic state will be described in a later article.

Now we come to the clinical part of our paper, viz.:

Three Types of Asthmatics. In various articles, which the writer has published, he has pointed out the different varieties of the affection called bronchial asthma. During the past years many such patients have been treated according to the methods described in an article published in the year 1915.⁸

The result of such treatment has been excellent in some instances, discouraging in others. This, by the way, holds good for every other form of therapeutics. Not long ago a paper on the treatment of bronchial asthma was read before the New York Academy

of Medicine. On listening to it and to the discussion, one was impressed with the idea that that was the last word in the therapy of asthma. And yet the writer saw many cases which demonstrated the absolute futility of such medication.

As a whole the endobronchial method has given the writer much better results than any other measure, and for that reason alone he would advocate its use in the hands of laryngologists. But endobronchial therapy is still in its infancy, and it will take some time before strict indications for our procedure, especially the application of astringents to the lower air tract, can be laid down. The lumen of the smaller bronchi, at least in asthmatics, is narrowed down to such a degree that there is always a possibility of adhesions following the use of astringents. Formerly, when nearly every general practitioner was in possession of a galvanocautery apparatus, numerous such occurrences were observed after its employment in the nose.

Under normal conditions it is very likely that a powerful caustic, like the galvanocautery, is required to produce such an effect. Where there is an abnormal mucosa, like that of the bronchi in asthmatics, however, it is very probable that any solution of nitrate of silver strong enough to be effective could bring about such an untoward result (instead of silver of nitrate, we may, of course, take any other astringent). That the mucosa in this class of cases is pathologically altered is well known. That the epithelium also is occasionally destroyed we learn from one post-mortem reported by Berkart. Other writers as Ad. Schmidt have found the epithelium well preserved, but the single observation of Berkart is sufficient to put us on our guard. Consequently great care has to be exercised not to close up entirely the bronchi already narrowed by the asthmatic affection.

True enough, applications to the larger bronchi alone occasionally exert a beneficial influence on the lower portions as well and materially improve the asthmatic state. But we have serious doubts as to the possibility of curing a series of cases by such means, as was done by W. S. Syme^o of Glasgow. Syme swabbed the main bronchi and the entrance to the secondary bronchi with a ten per cent solution of nitrate of silver and reports surprisingly good results in eight consecutive cases. Such publications have to be taken with more than one grain of salt. Nevertheless, the endobronchial therapy by means of drugs as well as by the topical application of the faradic and galvanic currents, as advocated by the

writer, has done so much good that they have become part of our armamentarium medicum.

Today only three types will be reported, showing the characteristics of each class. That does not mean that these are the only types of asthmatics seen in practice. There are many more, if one considers the etiology of the disease, a fact discussed in previous articles.¹⁰ These types are described here, because they were of great interest to me, and for no other reason.

Type I. Mrs. F. G., a middle-aged lady, had been referred to me by an ophthalmologist of this city some three years ago on account of lachrymation and sneezing. A year before consulting me she had been operated on intranasally by a well-known laryngologist abroad for empyema of the left frontal sinus. At the same time nasal polypi had been removed. Her main complaint now was the dreadful attacks of sneezing that occurred principally in the morning. On examination masses of granulations were found in the left nasal cavity and there was a thick, purulent discharge coming from the frontal sinus. An external operation having been refused, she was treated in the usual way intranasally. She called on me only every few months, when there was stagnation of pus with the concomitant symptoms of headache, bad odor, etc. It was especially the latter that annoyed her greatly at such times. She had been taught by me to syringe her frontal sinus and did it quite well. But whenever polypi or granulations formed again, such procedures were of no avail. In 1917 she stayed away for eight months and the first thing I heard from her was that she had been suffering from asthma. She had remained in the city all summer, was kept in bed most of the time and given narcotics and other drugs by her family physician. The absolute uselessness of such medication was demonstrated clearly by the almost immediate feeling of relief following the removal of several polypi and washing out of the frontal sinus.

Such cases undoubtedly have occurred in the practice of other laryngologists. I myself have described cases of asthma caused by ethmoiditis and other forms of suppuration in the nose. But this is the very first one which I could trace with absolute certainty to the frontal sinus.

Formerly, and for many years, such an affection was called a reflex neurosis. As a matter of fact it is nothing else but the result of the secretion running down from the nose into the larynx and bronchi, as pointed out by me several times. No doubt, even

in this patient, there was a nervous element that helped to bring about the paroxysm, but that was not the main feature. It certainly developed *post hoc* and *propter hoc*.

It should be added that in some predisposed persons simple irritation of the nasal mucosa is apt to produce an attack of asthma. How far the atony of the infundibula, the bronchioli and all inspiratory muscles has progressed in such cases can only be surmised.

Type II. This case represents an entirely different aspect. Its history is shortly the following: M. C., an agent, 31 years old, had been in perfect health until his nose began to trouble him. Numerous polypi were removed in different clinics, where he may have been handled somewhat roughly according to his story. As a result of these operations "laryngitis" set in and then bronchial asthma. The latter has persisted ever since, that is for three years. He gets the attacks at all times of the day and night and is suffering so badly that he had to give up his business and be supported by his family. There is no specific, nor rheumatic history. He has three healthy children and his wife had no miscarriages.

When I examined him, there were polypi on the right side and an ethmoiditis. The question now was, what to do with a patient like this. Should we proceed as in any other case, removing the polypi, etc., or should we pay attention only to the asthma, leaving the polypi alone? I decided to follow the latter course and treat the asthma. Endobronchial applications were made first of adrenalin, later followed by argentum nitrate, and the endobronchial use of the faradic and galvanic currents. But his nasal condition was not neglected entirely during treatment of the asthma. The secretions were removed and mild applications made to the nose and throat to secure temporary relief of trouble from these parts.

After a few months' treatment he was able to resume work and he began to regain confidence in himself. Then the polypi, which by that time were present on both sides, were removed gradually. Of course, this had to be done carefully in order to avoid any shock to his already shattered nervous system. On and off he received endobronchial treatment while attending to his business, and for more than a year after that he has had no spasmodic attacks. His subjective feeling is expressed in a letter he wrote to one of his friends, who happened to bring it to me. He says: "* * * I feel that since I was treated by Dr. F., life looks bright to me again. I can sleep the whole night without getting up, with

the exception for a drink of water occasionally owing to a dryness in the throat. * * * Letters like this I have received before, but have not published them, as most of these patients cannot be relied upon. Much against my taste I cite part of this letter, as I know the firmness of this man's character so well. He has been cured for more than a year and I feel that he will remain so.

This case offers an interesting problem. For years we have been taught that asthma is often due to nasal polypi and it is a fact that many a case of asthma has been cured by removing these growths. In the opinion of some this case tends to upset this doctrine. The patient developed asthma after he had been treated for nasal polypi and improved while the nose was being more and more occluded by these neoplasms. Although of late my belief in the causation of asthma by nasal polypi has been shaken materially, as rhinologists we cannot do otherwise than remove them whenever they are present. Yet this case teaches us a lesson, i. e., to go slowly in neurotic cases. While in some patients you may, in one sitting, remove as many polypi as your time and the loss of blood will permit, it is necessary to be careful in others, lest shock follows.

The case referred to is the result of such a shock, that undoubtedly could have been avoided by a different mode of procedure. It is also typical of what I have called traumatic asthma, which generally is due either to operations or accidents.

Type III. Mrs. R. L., 39 years old, has had asthma for the past twenty years. At times she was free from it, but it always returned. For three years she has suffered almost constantly, with the exception of short intervals in summer. Slight constipation, poor appetite. As she had been under the care of a practitioner of wide experience, nothing that promised a cure was left undone. Her nose was operated on twice, but she still sleeps with her mouth open. She underwent all kinds of therapeutic fads with vaccines, protein extracts, egg-free diet, oatmeal diet, rest cures, etc. Even some of her teeth were extracted in order to eliminate any possible focus, but all in vain. When I examined her, there were cheesy masses in one tonsil, a marked hypertrophy of the middle turbinal and pus in the middle meatus, apparently coming from the right antrum.

It is but natural that my first impulse was to remove all these pathological conditions in spite of my experience with the case just narrated. But I resisted that temptation and tried to make topical endobronchial applications which, however, I am sorry to

confess, were impossible with her. She was so restless that I never was able under local anesthesia to introduce either a small calibre bronchoscope or the endobronchial spray-electrode. General anesthesia was refused. Finally, it was proposed to her to operate on her nose and throat, to which she would only consent if a cure of the asthma could be guaranteed. Consequently nothing was done and the case proved a total failure therapeutically. The reason was that it never was possible to get a hold on the patient, which I fear was mostly due to her excellent financial circumstances. She refused to believe that she could be cured, always looked with suspicion at everything we did or prescribed and could never be convinced that something could be done to benefit her. This poor rich woman was so much absorbed in her own sickness that only a supernatural occurrence would have diverted her attention from her own person.

The instruments used by me are the same as described in 1915. In order to introduce the endobronchial spray I use a Krauss laryngeal forceps which has been modified so that you can grasp the hard rubber tube easily. Since it is impossible to have one of the latter made nowadays I had a soft rubber tube connected with a De Vilbiss spray or vaporizer and introduce the soft tubing as far as I can get, by means of the modified Krause forceps.

BIBLIOGRAPHY.

1. LUCIANI: Human Physiology. Translated by Welby, London, 1911.
2. DIXON and BRODIE: *Journal of Physiology*, 29, 97, 1903.
3. I. C. WALKER: Studies on the Cause and Treatment of Bronchial Asthma. *Jour. A. M. A.*, Aug. 4, 1917.
4. GOLLA and SYMES: Artificial Respiration. *Journal of Physiology*, 46, 1913.
5. GOLLA and SYMES: Loc. cit., p. 88.
6. D. E. JACKSON: *Jour. Pharmacol. and Experimental Therapy*, 1913, p. 87 and 299.
7. EPPINGER and HESS: Zur Pathologie des vegetativen Nerven-systems. *Zeit. f. klin. Med.*, 1909, p. 205. Die Vagotonie. *Sammlung klin. Abhandl. über Pathol. u. Therap. des Stoffwechsels*, 1910, H. 9-10.
8. W. FREUDENTHAL: On the Direct Galvanization and Faradization of the Bronchi and Their Topical Medicinal Treatment in Bronchial Asthma. *American Medicine*, March, 1915.
9. W. S. SYME: *British Med. Jour.*, June 30, 1917, p. 868.
10. W. FREUDENTHAL: Contributions to the Therapy of Bronchial Asthma. *New York State Jour. Med.*, Dec., 1916.

59 East 75th Street.