As we near the tropics, puberty appears increasingly early, while at, or near the equator, it is often retarded. Poor food and other deleterious factors may also retard puberty. Women in the far North menstruate sparingly, some Esquimaux only during summer. At high altitudes, puberty appears late. However, immigrants passing from one climate into another, frequently mature at the same age as if they had remained at home, even for several generations.

Another interesting observation on man, which was borne out by these experiments on rats, is the fact that, as we approach the tropics, the external sex characteristics, such as the distribution of hair, approach each other in the two sexes.

R. E. WOBUS.

Wiltshire: Basal Metabolism in Menstruation. Lancet London, 1921, cci, 388.

The theory that the life processes of women undergo a periodic variation correlated with menstruation has been frequently suggested, and the effect of menstruation on the life processes of women studied in various ways.

The author undertook a series of experiments comparing the physiologic processes in the menstrual and intermenstrual periods. The points chosen for determination were: (1) the basal metabolic rate of normal women during menstruation and between menstrual periods; (2) the cost to the organism of a certain definite piece of work and (3) the rate of recovery from that work.

The observations were made on five subjects, the basal metabolism being determined each day during the menstrual periods and three or four times between these periods.

The results obtained showed that the basal metabolism was not appreciably affected by menstruation. The cost of work to the organism, and the rate of recovery from work were the same during the menstrual and inter-menstrual periods. The author concludes, that while more experiments must be done before any definite conclusions can be drawn, the processes appear to be identical during the menstrual and intermenstrual periods.

NORMAN F. MILLER.

Clow: Menstruation during School Life. British Medical Journal, 1920, No. 3119, 511.

The author as medical inspector of a large girls' school had the opportunity of studying the menstrual function in 1200 healthy girls. The ages of these girls varied from 9 to 21 years. All of them were interviewed and examined once. The following information was secured: Regular menstruation occurred in 53.8 per cent; irregular menstruation in 46.1 per cent; no pain or discomfort in 73.0 per cent; discomfort or slight pain in 24.6 per cent; severe pain in 2.4 per cent; incapacitated 5.3 per cent; malaise 9.0 per cent; rest required during the period 23.0 per cent; baths taken during the period 23.4 per cent; games and usual exercise continued during the period 40.7 per cent; constipation during the period 9.0 per cent. The girls were advised to act normally during their periods, to bathe and exercise (except swimming) as they would do unless sick from some other cause. Of these girls 734 were seen more than once after having followed this advice and the following results were noted: Those having no dis-