acute, that the least agitation, such as is caused by the passage of the induction current, aggravates the distress.

CASE VIII.—Mrs. B., aged 34, was taken with acute crural neuralgia, when she was almost convalescent from an attack of inflammation of the womb. She suffered for nine weeks without experiencing any relief, when her physician advised the use of electricity. We gave her but three applications, but from the very first the pain was aggravated, and in no slight degree.

We of course discontinued the applications, but she continued to suffer for many weeks, before the disease at all abated.

A case somewhat similar to the above was sent to us by Prof. Willard Parker, of this city, but the results from the use of electrization were almost as unfavorable. It is possible that the use of the galvanic stream might have been followed by better results, for we have in a few instances seen cases of neuralgia yield to this current, when the Faradaic was inoperative.

Hospital Reports.

BOSTON CITY HOSPITAL.

Monthly Report of the more important Operations, by
G. F. JELLY and L. D. GUNTER, House-Surgeons.

CASE I.—*Necrosis of Radius*.—(Service of Dr. THORNDIKE.)—Geo. O., aged 17, began to notice pain in right arm over lower anterior portion of radius, six weeks before entering hospital. An abscess formed, and broke in about a week. Never received any injury. Patient etherized and an incision about three inches in length made over lower part of radius, extending from a small sinus near lower extremity directly upwards parallel with the bone. A loose sequestrum about three inches long was removed with dressing forceps, and on passing in the finger not a particle of denuded bone could be felt, nor has there been any since; and now, three weeks after the operation, the wound is nearly healed.

CASE II.—*Necrosis of Tibia*.—(Service of Dr. THORNDIKE.)—W. D., aged 46, laborer, fell fourteen (14) feet, striking on a pile of iron and broke left tibia and fibula. Fibula broken ($1\frac{1}{2}$) one and a half inches above lower extremity, and internal malleolus was broken off and the fracture was compound.

Two (2) months afterwards there was necrosis of tibia, but fibula had healed.

Opening enlarged and internal malleolus, which was entirely loose and necrosed, removed.

Three weeks after operation, wound closing well; there is considerable motion in joint, though owing to an abscess in opposite ankle there is a free circulation of pus. Astragalus not denuded, nor is the tibia.

CASE III.—*Ununited fracture of Tibia and Fibula*.—(Service of Dr. THORNDIKE.)—M. M., laborer, aged 40, received a compound fracture of right leg, five months before admission; was treated at Massachusetts General Hospital four months, and discharged with non-union of fragments.—Operation, Feb. 14th. Ends of tibial fragments denuded, and a thin section made from each fragment; periosteum dissected up; sound fragments united by a silver ligature, and leg placed in fracture-box. In two days erysipelas became developed in leg and thigh, and he was also attacked with intermittent fever which he had been free from for several months. Two weeks after operation, erysipelas has all disappeared; wound healthy and fragments are adherent. General condition very good.

May.—Three months later, fair union.

CASE IV.—*Hæmorrhoids*. *Death*.—(Service of Dr. THORNDIKE.)—M. B., laborer, aged 60, had hæmorrhoids seventeen (17) years; moderate size. Etherized, and four strong ligatures applied; lost six or eight ounces of blood during the day. A typhoid condition became developed soon after the operation, and he gradually sank and died in a little over a week. Physical signs showed congestion of the lungs.

CASE V.—*Circumcision for Epithelioma of Prepuce*.—(Service of Dr. THORNDIKE.)—J. B., aged 60, laborer, has had epithelial disease of the prepuce for about nine months. It began as a pimple, and has increased to the size of a small English walnut. Neither painful nor tender. Covered with a small dark sloughing. Glands unaffected.

March 13th.—Prepuce, to which the tumor was confined, was removed in the usual way. Skin and mucous membrane stitched together. Patient did very well, and was discharged well, March 23d.

CASE VI.—*Plastic Operation*.—(Service of Dr. THORNDIKE.)—E. T., aged 35, widow, entered hospital with an indolent ulcer of the heel of a year's standing.

March 20th.—Operation. Surface of the ulcer with a narrow rim of the edge removed, leaving an open surface one and one-fourth inch in diameter.

On the outer surface of the middle third of the opposite leg, a flap, two inches in diameter, with a pedicle one and a half inch wide, was dissected up and fitted accurately into the prepared wound of the

heel and secured by silver sutures; the leg being placed in an angular trough and supported by a frame.

Unfortunately the patient had varicose veins, and had been troubled more or less before entering the hospital with diarrhoea. For three days following the operation she suffered very little pain; took ten grains Dover's powder night and morning, and nourishment freely. No violence was done to flap, the position was easy and temperature good.

March 24th.—Legs returned to their natural position, and the flap detached from the heel entirely, although it was adherent over about half a square inch in the centre of the ulcer. Ulcer looked well and was covered with granulations; the greater part of the flap, not attached directly to the heel, was alive, and was allowed to fall back to its original place. General condition rather poor; appetite failing; bowels constipated; very little pain. Extra diet, stimulants, &c. Weak solution of carbolic acid applied to ulcers.

Mar. 26th.—Erysipelatous blush over left knee, and also on the edges of wounds of same leg. Appetite fair, pulse 100; tongue clean and moist; bowels constipated.

31st.—Erysipelatous inflammation extends down to the foot. Appetite poor; tongue brown and dry; pulse 100, rather weak; not much pain; ulcer of heel never looked better. Generous diet, stimulants, &c. *R.* Tr. ferri chloridi, *gtt.* xx. every four hours.

CASE VII.—*Perinephritic Abscess.*—(Service of Dr. CHEEVER.)—H. L. H., aged 5. Four weeks ago complained of pain just over crest of ilium, on the left side, and soon began to walk lame. Five days ago, his mother first noticed a swelling at the seat of pain. On entrance, there was a large fluctuating swelling, between the crest of the left ilium and the ribs, and filling the space from the spine to the anterior superior spinous process of ilium, not painful, but very tender to the touch. Skin tense, but not red. No curvature or tenderness of spine. No pus in urine.

Etherized, the abscess cut down upon, and six ounces of pus, so thick that it would scarcely run, evacuated. The cavity filled the whole space, mentioned above, as the extent of the swelling. No diseased bone discovered.

After the operation, his condition was much improved, and there was very little discharge. On the fourth day after operation, pus was found in the urine. He continued to improve in every way, and three

days later, only a very small amount of pus was found in urine. It is now four weeks since the abscess was opened. The wound has now healed, and his general condition is very good. No pus in the urine.

CASE VIII.—*Operation on Cicatrix of Burn.*—(Service of Dr. CHEEVER.)—H. S., aged 15. Thirteen years ago patient burned her right hand. The fingers are now drawn into the palm, by the cicatrix, and grown together. The joints are movable. The metacarpal bones are better developed than the phalanges. The whole hand, which is about half the size of the other, is clenched in such a manner, as to resemble a well-formed stump, being covered in a bag of closed skin.

Etherized, and incisions were made to free the thumb and the index finger. The terminal phalanx of the latter was removed. They were separated by sponges till suppuration commenced, when the hand was placed upon a posterior splint, and an attempt made every day to straighten the finger and thumb. There is considerable mobility in the joints; there is little pain, and the granulations are looking well. Ultimately a fair result was obtained.

CASE IX.—*Lumbar Abscess.*—(Service of Dr. THORNDIKE.)—Dec. 3d. W. F., aged 11. Boy of scrofulous diathesis, had lateral curvature of spine, and a hard tender tumor in right inguinal region, of three weeks duration. Swelling, pain, tenderness, &c., gradually increased till Jan. 10, when the tumor extended round over the crest of the ilium, nearly to the spine. A small opening was made into the abscess, and several ounces of laudable pus evacuated. No disease of ilium or vertebræ could be detected. Jan. 31st.—Nearly well; discharge very slight.

CASE X.—*Lumbar Abscess; Pott's Disease.*—(Service of Dr. THORNDIKE.)—M. V. B. H., aged 30, a porter. Has had angular curvature of spine five years. Jan. 24th, has a large abscess over dorsum and crest of ilium of two months duration. Evacuated with trocar and canula, and filled with a mixture of one part tr. iodine to two parts water.

Jan. 31st.—Discharge is free, but his general condition has improved but very little.

CASE XI.—*Dislocation of Shoulder.*—(Service of Dr. CHEEVER.)—R. G., aged 56. This morning fell upon right shoulder, producing a sub-coracoid dislocation. Etherized, and reduced by heel in axilla. The arm confined by Velpeau's bandage.

CASE XII.—*Dislocation of Humerus.*—(Service of Dr. THORNDIKE.)—J. Q., laborer,

aged 47. While drunk, fell down stairs, Jan. 17th, striking on head and top of shoulder, inflicting a scalp wound on the former, and dislocating the humerus into axilla. Patient thoroughly etherized, and the dislocation easily reduced by extending the arm and lifting the head of the bone into its place. Jan. 31st, doing well.

Reports of Medical Societies.

ANNUAL MEETING OF THE MASSACHUSETTS MED. SOCIETY—SECOND DAY.

The members of the Society re-assembled on Wednesday, June 3d, at the amphitheatre of the Massachusetts General Hospital, Dr. H. C. Perkins in the chair. The Secretary of the Councillors reported that sixty-six new members had joined the Society during the year, and that twelve had died.

An amendment to the by-laws was passed, providing that there be a meeting of the Censors of the Suffolk District held on the days succeeding the medical examinations of Harvard University, instead of the first Wednesdays in January and July as at present.

The following officers were appointed for the ensuing year:—*President*, C. G. Putnam, Boston; *Vice President*, H. L. Sabin, Williamstown; *Corresponding Secretary*, C. D. Homans, Boston; *Recording Secretary*, C. W. Swan, Boston; *Librarian*, J. C. White, Boston; *Treasurer*, F. Minot, Boston; *Orator*, A. Hitchcock, Fitchburg; *Anniversary Chairman*, J. B. Upham, Boston.

The Treasurer's annual report showed the receipts of the year to have been \$8,533, and the expenditures \$6,511. Of the expenditures, \$2,031 was for publication, and \$1,500 for the dinner at the Music Hall. The funds of the Society amount to \$30,420.78. The expenses of the last annual meeting were \$1,747.75.

The following-named delegates were present from other States:—A. J. Fuller, Me.; A. W. Nelson, Conn.; J. J. Caldwell, J. G. Wilbor, C. A. Lee, New York; John Kirwin, Penn.

A letter was read from Dr. Thomas S. Kirkbride, President of the Association of Medical Superintendents of American Institutions for the Insane, thanking the Society for their invitation, and stating that a committee would be present at the dinner.

Dr. J. Baxter Upham, of Boston, read a paper upon the doings of the Medical Com-
Vol. I.—No. 19a

mission to the International Convention at Paris last year, giving a full abstract of the report of that Commission, which related entirely to medical subjects.

Dr. Echeverria read a report of a case of epilepsy, and showed photographs of sections of the spinal cord.

Dr. J. B. S. Jackson showed a very interesting specimen of complete bony union of intra-capsular fracture of femur, occurring five or six years before death in a lady nearly 70 years of age.

Dr. John M. Harlow, of Woburn, read a very interesting paper, containing the history of the famous case of a man named Gage, a former resident of Lebanon, N. H., who, while blasting rocks at Cavendish, Vt., in 1847, by reason of premature explosion had a tamping iron, three feet seven inches long and one and a quarter inch thick, and tapering to a point, forced through his head. Dr. Harlow attended the man, and gave in detail the daily symptoms of his patient. The iron entered the left side of the face, and came out about the centre of the top of his head. It was supposed at first he would die, but in fifty-nine days after the accident occurred he was able to walk and ride, and was soon nearly as well as before, although his intellect was somewhat affected. This has been doubted by many prominent surgeons. Gage died May 21, 1861, twelve years, six months and eight days after the injury. Since the death of Gage, Dr. Harlow procured his head, and has presented the skull to the Warren Museum of the Harvard Medical College. The skull and iron were exhibited by Dr. H., and they were examined with much interest by the gentlemen present.

Dr. H. J. Bigelow stated his connection with the facts mentioned above, and said he had seen Gage twenty years ago, and was then satisfied of the reality of this wonderful case. He also called the attention of the meeting to a similar accident, by which a tube of iron, five eighths of an inch in diameter and five feet long, passed through a miner's head while blasting coal in Ohio, and which was pulled out by a fellow-miner. The injured man was introduced to the audience, and Dr. Jewett, the attendant physician, recounted the case in detail. The young man's mind has not yet been fully restored.

PRIZES.

The Committee (consisting of Drs. B. E. Cotting, H. J. Bigelow, C. E. Buckingham, D. W. Cheever and C. Ellis) appointed last year to receive and examine dissertations