or four inches in front of the uncovered eye. As it is slowly moved from its first position to a point three or four feet away, the patient is asked how many matches he sees. In most cases the match will present a double image when held quite near the face; the images approach each other and become confused as they are removed, to again separate more and more until the meter distance is reached. The match is again, from this point, gradually brought close to the eye, when the same phenomena, but in reverse order, will be manifest. The second eye is similarly examined and, finally, the room is darkened and a further (control) test is made with a small candle flame. Sometimes three or more images (polyopia) are observed and it is usually possible to exclude one or more of these by interposing a card, so as to cover various segments of pupillary area during examination.

A few words about *pupillary anomalies* in hysterical subjects, because there is much confusion on this point. As a rule, when either or both pupils are unusually contracted or unusually expanded the ordinary reflexes are preserved, that is, they *contract* when light is thrown upon them and when suddenly asked to fix a near object, and they expand when light is withdrawn or when the patient is told to gaze into the far distance. This is, or ought to be, a very simple matter, but in cases of hysterical amblyopia some care should be observed in making the examinations. The patient should be seated facing a half-lighted window; the unclosed eyes are completely covered with a black cloth and he is told to look, and to continue to look, as if gazing upon a distant object which has been previously pointed out to him. In thirty seconds the cover should be suddenly removed and the contractions of the pupils, or its absence, noted. The reflex contraction of the pupils for convergence or accommodation should be tested in a light as dim as is consistent with the observer's ability to see the patient's pupils. Having been told to look at an object across the room for half a minute, he is now asked to quickly fix the end of the finger held four inches from the patient's face. By means of these simple but effective devices one may often avoid the mistake of concluding that he has to deal with a pupil that does not respond to the reflexes mentioned.

I need not remind you that in hysterical amblyopia we frequently find *macropsia* and *micropsia*. Usually the patient complains of this strange symptom, probably due to irregular contracture of the ciliary muscle, but it is often worth while to test for it. A long, lighted candle is held before each eye of the patient at distances of one, four and ten feet, and he is asked whether it gets longer or smaller in size. Notes are made of his answer and the experiment repeated in a day or two.

A very common and, in my opinion, characteristic eye-sign in hysteria is spasm of the orbicularis, the so-called blepharospasm. When this is unilateral and accompanied by photophobia, or spasm of accommodation, it is almost invariably hysterical, and I believe that the majority of the spasms of the orbicularis are of this character, whether in the form of blinking or constant winking of the eyes, or where the spasm is much more marked and involves the facial muscle.

The Phonendoscope as an Ear Trumpet.—Surprising results have been obtained with deaf subjects, especially in reading aloud to them, by removing the external rubber plate of a phonendoscope, and speaking directly into the apparatus, as into a phonograph, with the tubes in the deaf person's ears.

HYSTERIA IN CHILDREN.

Presented to the Section on Neurology and Medical Jurisprudence, at the Forty-ninth Annual Meeting of the American Medical Association, held at Denver, Colo., June 7-10, 1898.

BY J. G. BILLER, M.D.

CHEROKEE, IOWA.

Hysteria is recognized as well worthy of our deepest study and best thought, by those most familiar with it. Notwithstanding this fact, and the volumes that have been written about it, we must acknowledge that in many ways we are woefully ignorant regarding it. We all, probably, feel at times as though, if there were anything which we have not found out by practical experience pertaining to this disorder, we are perfectly willing to remain uninformed.

No matter how willing we may be to call it quits, our omnipresent foe (hysteria) never will consent to that, and by appearing in new phases and new forms, as well as new combinations, forces us to learn more and more regarding it. The more we know of it the less liable we are to use some such an expression as, "Oh! it is nothing but hysteria, and does not amount to anything!" No one would want to say that of any other disease which causes so much suffering and unhappiness, and destroys the pleasures and comforts of as many homes as does hysteria. The temptation thus to speak of it is because we often feel that if the patients would, they could, control themselves, and we back up our ideas with the fact that some simple but disagreeable treatment has often caused the entire disappearance of the symptoms, and for this reason it is thought that if very much were the matter the result would have been different. These brilliant and dashing flank movements are not always successful, and we have to settle down to a regular siege, from which we often withdraw defeated, and too often our defeat is caused by our treating our foe with contempt; the patient becomes dissatisfied and passes, frequently, into the hands of some quack or charlatan, who thrives by accidentally—and probably unconsciously—knowing how to take advantage of some of the tricks of this powerful but susceptible enemy of the human family. Every physician, after having hours, days and weeks of his time taken up with hysterical patients, must oftentimes ask himself: "Where did all this host of hysterical men and women come from?" The answer to the question is: "From the mis-born, and mis-raised boys and girls of the past generation, and the children of the same class in the present generation are hurrying forward to recruit the ranks." I believe a hysterical woman can oftener be cured before she is fifteen years old than after, and it is for this reason that I wish to call your attention to hysteria in children.

This subject, as far as treatment can go, may be divided, primarily, into two classes: the first, those cases which never have hysteria till after puberty, but whose physiologic and psychologic development have been such as to leave the seeds well planted in a soil that has been carefully fertilized for their maturing. This would hardly be called hysteria in childhood, but the principal damage is done then, and that is the time when proper treatment should have been applied.

In the second division, we have the disease fully developed during childhood. It is often surprising to note what apparently slight suggestions will develop a hysterical attack in a young child. I remember a boy 3 years of age, who, upon getting up in the

morning, a week or so after having seen a man who was paralyzed in one leg go by a few times, said his leg was no good, and he could not use it and did not; careful examination was made, and nothing abnormal discovered except that he would not use that leg; he dragged it around all day, but that night, after he was put to bed, his father went in to play with him, and in the excitement of the play his paralysis disappeared, and he seemed never to think of it again. This was a child of very active imagination, and his mind had undoubtedly been impressed until he was unable to move his leg.

I have since then had two cases of partial paraplegia of hysterical origin in children. They were sisters, and of quite dull and heavy minds, whose imaginations were slow to be excited, and when the result was produced, it was just as slow and tedious to overcome. The father of these children had died of tubercular trouble, which started in the back, and one of his legs became nearly useless from sciatic neuritis. The children, after the father's death, were left in the mother's charge; she unwisely had a great deal to say about the father's sickness, and especially about his back and the inability to use his legs. The children were fond of sweet food and were allowed to indulge in cake, pie, etc.; in fact, this was about their whole diet. The living room was lighted by two small windows which were shaded by porches, and was heated in winter by a hard coal burner which was not much smaller than the room; the children's health was carefully watched by the mother; every little pain or nervousness was inquired after and examined into. The elder of these two girls (who were the youngest of a large family) had a spinal curvature, which became tender and painful; a short time elapsed before a cast was applied, and during this period she suddenly became unable to move the legs, and when put on her feet would tumble over if not held up. If held up for a little while she could partially stand by rising on her toes. Reflexes and sensation were normal, and after the application of a well-fitting plaster cast the legs became all right.

About a year later, the younger child, then about eight years old, was brought to see me for severe pain in the head; her temperature was above normal, pulse rapid; this continued for several months, and then began to improve, but all at once the legs became useless; she would not stand on them, and part of the time could not move them. Reflexes were normal; there was some pain in the crural nerves, otherwise sensations were normal. Various kinds and methods of treatment having been tried without success, galvanism was used for two months. At the end of this time she began to improve, and was soon able to be up and around; in a few weeks more she seemed as well as usual. She remained well for a year, when she had a slight tonsillitis, and before she was relieved of this her legs were again paralyzed. The battery was resorted to, and in a few weeks it had the desired result, although, to the best of my belief, no current came from the battery. No. 2 had hardly got her legs to work before No. 1 took cold and had paralyzed legs; the battery was still in the same condition, and it took six weeks to get the girl around.

This last attack was about a year ago. I have urged a better diet and more light in the living room and no conversation about sickness or sick people; but it is hard to convince the mother that anything but inheritance has aught to do with the sickness.

Paralytic hysteria, as represented by the above cases, is not so common as some of the psychic forms, but it is so marked that it attracts attention at once, while the latter often does not attract sufficient attention to be diagnosed as hysteria. I believe some of the peculiar fears of childhood are often manifestations of hysteria. I knew a little boy who would scream and cry so that he had to be taken from the room if any one began singing or playing the organ; this child afterward developed very marked hysterical symptoms.

Night terrors are undoubtedly, at times, of hysterical origin; for instance, a child three or four years of age, awoke one night screaming, and said the rats were after him; the next morning he did not want to eat breakfast in the dining-room for fear of the rats. This kept on night after night, and soon his naps in the day time were troubled with these terrors, and it was almost impossible to do any thing with him for hours at a time; regulation of diet, amusement and medical treatment, were all tried without benefit. The condition was carefully explained to the mother, who was advised to make him control himself. That afternoon he had a nap, and awoke the same as usual with the wild cry of rats; he would not sit on a chair or stay in bed or even be quiet in his mother's arms. No man with delirium tremens ever had greater fear of snakes than that poor child had of rats; the mother told him in a very decided tone that he would be whipped if he did not stop, and then, as he did not mind, he was spanked. This forced him to control himself for a few minutes; then he began again, and the spanking was repeated. This had to be done two or three times, but finally the child was forced to control himself, and was never troubled that way again. This may seem cruel treatment, but it became necessary, in some way, to force the child to exercise sufficient will-power to overcome his excessive imagination. The boy several times since has shown marked symptoms of hysteria, but owing to the care he has had has completely outgrown them.

I do not wish you to understand me as advocating whipping as a cure for hysteria, but in these psychical forms, the child must be brought, in some way, to control itself and taught to continue this, and there is no question but what fear forces not only children but grown people as well to keep their imagination and desires well in hand. The cases which require punishment are, I believe, rare, but all these unfortunate children should have a strong hand to help guide and direct their imaginations.

One of the best instructors in self-government is the association of the child with other children; our public school life is valuable for that if for no other reason; there is no question but that children are often abused and imposed upon, still they soon learn that they can get along better if they behave themselves, and that if they wish to enjoy their play they must be agreeable to their playfellows. This is an excellent discipline, although it may occasionally make the parents a little angry when they think their child has been used unjustly—as it often is—but we must all expect to suffer some for the public good. The rough plays that bring home torn clothes and bruised bodies and limbs are not to be looked down upon. It is the freedom of the boy to indulge in any sort of physical recreation that his desires may dictate that has much to do with making his sex less liable to hysteria than the sex of his less fortunate

sister. The boy, as soon as he is old enough to be around, is dressed so that he can run, jump or turn heels over head; in fact, his movements are not interfered with in any way, either by dress or custom, but how is it with his sister? Whatever of liberty there is left after the method of dressing is taken from her by custom and the fear of being considered immodest. I think the little girl was right, who, when reproved for turning heels over head in the front yard, said, "if people did not want to see her, they must not go by when she was turning heels over head." She had the true sense of woman's rights—the right to be healthy and enjoy health in physical recreation.

Children should be well fed, and these neurotic children, as a rule, are poor eaters, both in quality and quantity. They should be confined to plain, nourishing food, and then forced to eat it. People often tell us they cannot get their children to eat, and are surprised when you insist upon their being made to eat; but I have had the pleasure of seeing several of these rapidly improve under judicious feeding. One case I will briefly describe: She was a delicate, nervous little girl, who was thin and anemic, unable to sleep at night because of nervousness. She wanted no breakfast, unless it were a cracker or a cookie, scarcely any dinner excepting dessert, but at supper she generally did eat a little better. She was absolutely forced to eat a large dish of oatmeal and milk, without sugar, each morning; a dinner of bread and butter, potatoes and eggs or meat, and drink a glass of milk, no dessert; for supper, she had a dish of oatmeal and milk, bread and butter and some kind of sauce, generally stewed fruit, either dried or fresh. Besides this diet she was allowed plenty of fruit, and was sent early to bed. She has been on this stuffing treatment for nearly two years, and is now strong, does her school work easily and well, never has headache, sleeps well, except that she gets up early in the morning. This child's poor health was attributed to going to school, as is done in a number of cases, and the children are kept out of school. I think there is no question but that their school work is too much for them in their physical condition, but too often this condition is caused more by the improper home life than the school life.

Neurotic children suffer greatly from competitive examinations at school, and from the extra work that is often imposed upon them in preparing for school entertainments—especially in preparing for public recitations. I knew one young girl of 13 or 14 years of age who was standing her school work well till she was selected for an important part in some special exercise. She was impressed with the importance of her part, and from the careful drilling she got she became frightened and nervous, and finally ended by going into a hysterical faint after one of these drills. It was a year and a half before she was able to go to school again, and she had during that time vertigo and other hysterical manifestations. At the end of this period she started to school, but she has never been allowed to do any but regular work, which she has accomplished with ease.

I have not referred to the medicinal treatment, for it has been quite thoroughly impressed on my mind that drugs are seldom indicated, and their value is questionable. The price of healthy children is the same as that of Liberty—Eternal Vigilance—and unless parents and physicians are willing to pay that price, we can not expect to avoid hysteria and many of its relatives.

ORIFICIAL IRRITATION IN RELATION TO NEURAL DISTURBANCES.

Presented to the Section on Neurology and Medical Jurisprudence, at the Forty-ninth Annual Meeting of the American Medical Association, held at Denver, Colo., June 7-10, 1898.

BY GEO. V. I. BROWN, M.D., D.D.S., C.M.

DEAN AND PROFESSOR OF ORAL SURGERY, DENTAL DEPARTMENT MILWAUKEE MEDICAL COLLEGE.
MILWAUKEE, WIS.

It would seem, upon consideration, that many operations brought forward by the gynecologists during the last few years for relief of so-called reflex pains and other nervous disorders, particularly neuralgia and migrainoid headaches, have not given the relief that was promised for them. The same is undoubtedly true to a considerable extent in regard to the rectal surgeon's efforts, and notwithstanding the fact that deviating septa and hypertrophied turbinates have freely given off their offending portions, there still remain many who suffer constantly without a hope of relief; and in directing your attention into a new channel with regard to etiology and treatment of these conditions, I would urge the fact that, with a high nervous organization and free distribution of nerve-supply quite as complex as that of the other orifices referred to, the normal mouth has thirty-two teeth as additional predisposing factors, each with its own separate nerve-fibers from the wide-spreading carrier of sensation—the trigeminus. In addition to these connections each root is surrounded by a highly vascular peridental membrane, also in close connection with the blood- and nerve-supply, making it not unreasonable to suppose that all these taken together would offer a very frequent source of irritation. Considering also the fact that through caries the pulps of these teeth and the nerve elements they contain are often exposed directly to external irritants, the etiological factors, particularly with affections within the range of the direct nerve-supply through which they carry the sensation, ought in a large number of cases to be found to be within the oral cavity. That this is often true without being suspected, through misleading symptoms, the trouble being treated as having its origin in some other of the orifices, will be shown, I think, by the cases described later.

Literature upon the subject of this and kindred disorders is very vague. We find neurasthenia, anemia and numerous other emias, held in a general way to be responsible, and the recommended treatment beyond certain limitations, also becomes very general, depending chiefly upon general constitutional care, which is all very well so far as it goes, but as a rule the patients suffer on just the same. It is evident to those sufficiently interested to be more than casual observers that in close study of the symptomatology of those affections, with a view to discovering certain definite causes of irritation and the mode of procedure best calculated in each particular case to remove the etiological factors, lies hope of relief. During the last few years my efforts have been directed to the study of certain peculiar forms of trifacial irritation, particularly one form of which little or nothing in the way of description seems to be extant in the literature of the subject.

Without attempting to go into a general or even a partially detailed description of the various possible irritations from within the oral cavity, I desire to treat of what I have found to be a very common associate of neural disorders. It is a noticeable fact that in the