white corpuscles 19,500 to 3000, a fall in arterial pressure from 15.9 to 12.9, a relaxation of the refractometric index, a very clear modification of coagulation and the presence of albumin in the urine. It is during the course of the hemoclastic shock or some moments after that one may see some dyspnea appear, elevation of temperature and the presence upon the entire body of the large urticarian plaques itching extremely. We renewed this experiment several times and each time according to the nature of the effort demanded we saw appear a hemoclastic shock and a urticarian crisis.

Two most interesting facts were established further with this patient, on one hand the presence of an autoprécipitation of serum at the moment of the shock, on the other the condition of colloidal instability in which the serum of this patient was found so that a single intravenous injection of sodium chlorid was able in her case to determine a slight hemoclastic crisis and to permit a urticaria to follow. It is this method of therapy which we tried with this patient with MM. Widal and Abrami and which determined in her entire organism a series of small shocks of desensitization for avoiding the great shock caused by fatigue or effort. It is the first time that one could see the fatigue alone without any alimentation and without the introduction of any foreign elements produce in the organism a hemoclastic crisis followed by urticaria. It is a matter therefore, as in the attack of paroxysmal hemoglobinuria under the influence of cold, of a true autocolloidoclasia, following the expression of Widal, in opposition to the heterocolloidoclasia produced in the organism by the introduction of foreign albumin. [Author's abstract.]

Solomons, B. HERPES AS A TYPE OF VICARIOUS MENSTRUATION. [Dublin Journal of Medical Science, May, 1921, Ser. IV, No. 15.]

This patient, a woman of twenty-seven years of age, had a sore on her left cheek every month. It resembled a catarrhal herpes and remained for about a week. When it is present she gets the sensations which some women feel previous to menstruation. This herpetic eruption began when she was eighteen and has appeared monthly ever since. Absence of the uterus, tubes and ovaries was disclosed by gynecological examination.

Wood. ERYTHREDEMA. [Med. Jour, Austria, Feb. 19, 1921.]

According to this observer this condition is not uncommon. He has observed it in forty patients and he instances its occurrence in fifty-one cases as reported to him by a medical confrère. The symptoms are characteristic. The child holds the head bent down; usually it is whining and fretful. Some patients do not seem able to rest, scratching at their feet or pulling at their hair or ears, frequently making them bleed. If placed on the floor or in their perambulators they will bend their heads forward almost down to their feet. They do not smile, and they resent any attempt

to amuse them. In some cases the red swollen appearance of the hands is an early symptom. These patients become worn out for want of sleep and in absolute distress from the intolerable irritation of the skin of body, hands and feet. Sometimes they are vicious, scratching and biting at their mother or nurse.

Painter, C. F., and Bean, H. C. HEREDITARY EDEMA. [Boston Medical and Sur. Journal, May, 1921, CLXXXIV, No. 19.]

These authors class this syndrome as an endocrinopathy. If such cases are congenital, then there must be an inherited insufficiency or hypersecretion of some or all the endocrine glands. They do not analyze the cases to their ultimate sources apparently.

Lereboullet, P. ORGANO THERAPY IN SCLERODERMIA. [Bull. d. l. Soc. Med. des Hôpitaux, July, 1921, XLV, No. 26.]

In the extreme case of sclerodermia described, at the age of seventeen the face seemed to be a wax mask; the cheeks looked as if glued to the skeleton. The neck was hard and rigid, and it was impossible to take up a fold in the skin on the chest. The hands were bluish and the fingers immovably pressed against the palms. The upper part of the arms seemed the only regions that escaped the severe sclerodermia. Under treatment three years later with thyroid, pituitary and suprarenal extract simultaneously—all of which seemed to be called for by certain symptoms—plus phosphoric acid and arsenic tonics, marked improvement was realized even in two months, and by the age of twenty-four the sclerodermic condition and infantilism had entirely disappeared, the general aspect of the young man being now normal. [J. A. M. A.]

Weil, E., and Plichet. HYPERTRICHOSIS AND DIABETES. [Presse Méd., 29, 1921, 4.]

Hirsutism in women may be part of a pluriglandular syndrome, in the course of which there might develop glycosuria. The author's case occurred in a woman thirty-eight with well-developed beard, who succumbed to an intense saccharine diabetes. The patient also had pulmonary tuberculosis, the stage and degree not being given. Autopsy showed no tumor of the adrenal and the only glandular lesion was sclerosis of the ovaries. The totality of endocrine manifestations was the male distribution of hair in part (beard), virilism (no details), obesity, diabetes. The relationship is obscure and one can only compare this with similar cases. Tuffier and Guémès have reported cases of hirsutism in which the women showed respectively mild glycosuria and well marked saccharine diabetes. The latter readily yielded to diet and probably should really be regarded as glycosuria. Two other cases of hirsutics in women (patients of Laignel-Lavastine) were examined carefully for sugar in the urine, but although none was ever found, tests for