REVIEW OF AMERICAN CHEMICAL RESEARCH.

VOL. IX. No. 1.

WILLIAM A. NOYES, Editor.

REVIEWERS :

M. T. Bogert,	E. M. Chamot,	B. S. Cushman,
Benton Dales,	L. M. Dennis,	A. H. Gill,
H. M. Goodwin,	W. F. Hillebrand,	L. P. Kinnicutt,
H. W. Lawson,	H. N. McCoy,	A. A. Noyes,
J. W. Richards,	S. P. Sadtler,	J. O. Schlotterbeck,
W. H. Seaman,	F. P. Underhill.	

INORGANIC CHEMISTRY.

The Radioactivity of Thorium Compounds. I. An Investigation of the Radioactive Emanation. BY E. RUTHERFORD AND FREDERICK SODDY. *J. Chem. Soc. (London)*, **81**, 321-350.—A method of measuring, with considerable accuracy, the intensity of the effect of radioactive substances was devised. It is based upon the well-known property of such radiations of "ionizing" gases and thus enabling them to transport electrical charges. The magnitude of this effect was measured by means of a quadrant electrometer. This method is said to give much more accurate results than the ordinary photographic method. Thoria produces two kinds of radiations. One is of the ordinary sort and travels in straight lines. The other seems to be due to a radioactive gas. The latter emanation can be carried by currents of various gases through bent tubes, through plugs of cotton, through wash-bottles containing sulphuric acid or alkali without being retained or absorbed or having its activity diminished. To measure the radioactivity of this emanation, dry air or other gas was passed over a definite weight of thoria, then through a cotton plug and finally between electrodes connected on the one side with a battery giving 50 to 100 volts and on the other with the quadrant electrometer. The activity of the "ionized" air or other gas diminishes rather rapidly. The rate of "decay" was determined by causing the gas to pass slowly between three successive pairs of electrodes and noting the decrease of the effect. The radiations of different substances show different rates of decay, but the rate was uniform for all kinds of thorium compounds. It was found that samples of thoria from widely different sources had about the same activity. The emanating power of 10-gram portions of a certain lot of thoria was used as

the unit of comparison. The amount of emanation, that is, the effect as shown by the electrometer, was found to be directly proportional to the weight of the substance. Thoria loses its emanating power partially by heating. A sample having but 7 per cent. of its original activity was obtained by heating strongly. Loss of activity is not due to loss of moisture. The "straight line" radiation of thoria was not diminished by ignition. De-emanated thoria may be rendered active by converting it into chloride and then into hydroxide. The hydroxide was more active than the original thoria before de emanation. After four days it was $1\frac{1}{2}$ times, and after twelve days it was $2\frac{1}{2}$ times as active as the original sample. The activity of thorium hydroxide reaches a maximum only after a week or two. Samples of thorium compounds have been separated into small, very active portions and larger slightly active portions. These separations were made by fractional precipitation of solutions of the salts or by washing crude thoria repeatedly with water. The evaporation of the wash-water from 290 grams of thoria yielded 6.4 mg. of residue which was 1800 times as active as an equal weight of the original thoria. In this extremely active residue no other substance than thoria could be detected by chemical analysis.

H. N. McCov.

On Some Hydroxylamine Compounds. BY MAXWELL ADAMS. *Am. Chem. J.*, **28**, 198-219.—The methods of preparation and estimation of hydroxylamine are discussed, and a number of new compounds described. The method of preparation of Divers and Haga was improved in several ways. Sulphur dioxide was obtained from the commercial liquid substance. In the process of saturating the carbonate-nitrite solution with sulphur dioxide, decomposition was entirely prevented by using a small excess of sodium carbonate over two molecules of the latter substance to one of sodium nitrite. A method of readily separating the hydroxylamine sulphates from sodium sulphate is given, together with one for working up the mother-liquors by means of the phosphate. Tables of solubilities of the sulphate and the phosphate are given. The bromide, $\text{NH}_2\text{OH} \cdot \text{HBr}$, made from barium bromide and hydroxylamine sulphate, forms large acicular crystals, very soluble in water. Stable, if pure. $(\text{NH}_2\text{OH})_2 \cdot \text{HBr}$, plate-like crystals, very soluble in water. $\text{NH}_2\text{OH} \cdot \text{HI}$, from hydroxylamine and hydriodic acid, in ligroin solution, crystalline precipitate, unstable either dry or in water solution. $\text{HgSO}_4 \cdot (\text{NH}_2\text{OH})_2 \cdot \text{H}_2\text{O}$, crystals, rather unstable. $\text{HgCl}_2 \cdot (\text{NH}_2\text{OH})_2$, long acicular crystals from absolute alcohol. Decomposed by water, giving mercurous chloride. $\text{HgCl}_2 \cdot (\text{NH}_2\text{OH})_4 \cdot 2\text{HCl}$ crystals, from methyl alcohol. $\text{HgCl}_2 \cdot (\text{NH}_2\text{OH})_4 \cdot \text{HCl}$ crystals, soluble in water, with subsequent decomposition. HgCl_2 .

$\text{NH}_2\text{OH} \cdot \text{HCl}$, crystals, extremely soluble in water and alcohol. $2\text{HgCl}_2 \cdot 5(\text{NH}_2\text{OH}) \cdot \text{HCl}$, prepared in many ways. Flat hexagonal crystals, soluble in water, giving slowly mercurous chloride. $(\text{NH}_2\text{OH})_2 \cdot \text{HBr}$, crystals. $\text{HgBr}_2 \cdot (\text{NH}_2\text{OH})_4 \cdot 2\text{HBr}$, quadratic plates. The water solution decomposes. No compound of hydroxylamine with mercuric iodide could be obtained, owing to the rapid reduction of the latter to mercurous iodide. $\text{CdBr}_2 \cdot (\text{NH}_2\text{OH})_2$, fine crystalline precipitate from alcohol. Hot water gives a basic salt. $\text{CdI}_2 \cdot (\text{NH}_2\text{OH})_8$, large, colorless, acicular crystals.

H. N. MCCOY.

Address of the President of the British Association for the Advancement of Science. BY JAMES DEWAR. *Science*, 16, 533-551, 567-579, 621-631. (Delivered at Belfast, Sept. 10, 1902).—The address deals first with some of the important factors in the progress of scientific development in the last four or five decades in Great Britain and the parts played by some British scientists. The main portion of the address is devoted to a discussion of modern low temperature research, and the bearing of this work on the investigation of the upper regions of the atmosphere, auroras, comets and some related topics.

H. N. MCCOY.

The Precipitation of Ammonium Vanadate by Ammonium Chloride. BY F. A. GOOCH AND R. D. GILBERT. *Am. J. Sci.*, 14, 205-210.—The accuracy of Gibbs' method of determining vanadium by precipitation of ammonium vanadate by ammonium chloride having been questioned by several observers, the authors have repeated Gibbs' work. When Gibbs' exact conditions are fulfilled the precipitation is practically complete and the method yields good results.

H. N. MCCOY.

The Decomposition of Mercurous Chloride by Dissolved Chlorides; A Contribution to the Study of Concentrated Solutions. BY T. W. RICHARDS AND E. H. ARCHIBALD. *Proc. Am. Acad. Arts and Sci.*, 37, 347-361.—Mercurous chloride is acted upon by solutions of hydrochloric acid, sodium chloride, calcium chloride and barium chloride. Some calomel goes into solution and is apparently changed thereby into the mercuric form, while free mercury is deposited. The amount of mercury contained in the solution increases with the concentration of the soluble chloride. With equivalent solutions, up to about five times normal, hydrochloric acid and sodium chloride have about equal tendencies to effect the reaction, barium chloride has less tendency, calcium chloride still less, and cadmium chloride no appreciable tendency. The extent of the reaction, in solutions not too concentrated, is approximately proportional to the square of the concentration of the chlorine ion. This relation, taken in

connection with a number of other considerations, points to the existence of a highly ionized complex HgCl_4'' in the solution and thus confirms the work of LeBlanc and Noyes. The suggestion is made that this reaction may be of use as a means of determining the concentration of the chlorine ion. Corresponding reactions are much less marked with sulphates but much more so with bromides and iodides. Caution is recommended in the determination of mercury as mercurous chloride and also in the medicinal use of other chlorides in connection with calomel.

H. N. McCoy.

An Improved Drying Oven. BY EDMUND S. SMITH. *Sci. Am.*, 87, 186.—The oven is $9\frac{1}{2} \times 7\frac{1}{2} \times 6\frac{1}{2}$ inches. It is made of No. 28 galvanized iron and lined with $\frac{1}{4}$ -inch asbestos board. Heat is supplied by two 20 candle-power 110 volt lamps fixed in the lower part of the bath. One lamp alone gives a temperature of 70°C ; both give 110°C . Detailed drawings accompany the article.

H. N. McCoy.

The Recorded History of the Members of the Argon Group. BY HENRY P. TALBOT. *Technology Quart.*, 15, 195-228.—The article is an address delivered last February before the North-eastern Section of the American Chemical Society. It is a comprehensive review of the most important work done on the argon group. References to the literature are given.

H. N. McCoy.

An Acetylene Generator. BY H. A. RANDS. *Phys. Rev.*, 15, 178.—A simple form of generator, constructed from two flasks and two aspirator bottles, and designed for the production for laboratory use, of small quantities of the pure gas is described. A drawing of the apparatus is given.

H. N. McCoy.

Review of Some of the Recent Literature of the Periodic Law. BY R. H. BRADBURY. *J. Frank. Inst.*, 154, 283-290.—The article discusses chiefly the position of hydrogen and the members of the argon group in the periodic system.

H. N. McCoy.

On Some Double and Triple Thiocyanates. BY H. L. WELLS. *Am. Chem. J.*, 28, 245-284.—The more important results of this investigation are summarized as follows: (1) the description of numerous double thiocyanates (22 of them being new), among which the salts $6\text{KSCN} \cdot \text{Pb}(\text{SCN})_2$ and $\text{CsSCN} \cdot 4\text{TiSCN}$ present novel types, while the alkali metal salts with strontium and calcium show that alkaline earth metals enter into complex negative radicals more readily than has been supposed; (2) the conclusion, from a study of several series of salts, that the thiocyanates gen-

erally form double salts in smaller variety than the halides; and (3) the discovery of the existence of triple thiocyanates of various types, and particularly the preparation of four distinct triple salts from the thiocyanates of caesium, silver and zinc; in all, fourteen triple salts were prepared. EXPERIMENTAL.— $\text{Cs}_3\text{Fe}(\text{SCN})_6 \cdot 2\text{H}_2\text{O}$ (by W. K. Wallbridge): dark opaque tabular crystals with beautiful bronze luster, perfectly stable in the air, exceedingly soluble. Cs-Pb and K-Pb thiocyanates (W. K. Wallbridge and H. L. Wells). $\text{Cs}_3\text{Pb}(\text{SCN})_5$: colorless transparent crystals, apparently monoclinic; it requires the presence of an excess of caesium thiocyanate for its formation and cannot, therefore, be recrystallized from water unless a part of the lead salt be removed by filtration. $\text{KPb}(\text{SCN})_3$: brilliant white crystals, stable in the air, but decomposed by water. $\text{K}_6\text{Pb}(\text{SCN})_8 \cdot 2\text{H}_2\text{O}$: white opaque prisms, stable in the air. This 6:1 salt represents an entirely new type of double salts. *Cs-mercuric thiocyanates* (H. S. Bristol). $\text{Cs}_2\text{Hg}(\text{SCN})_4 \cdot \text{H}_2\text{O}$: large white prisms, somewhat flattened, rather freely soluble in water, especially on warming, and may be recrystallized unchanged, m. p. 168° – 170° . $\text{CsHg}(\text{SCN})_3$: large colorless crystals, often of pyramidal habit and with numerous faces, difficultly soluble even in hot water. $\text{Cs}_4\text{Mn}(\text{SCN})_6$ (C. S. Leavenworth): pale yellow-green crystals, massive or in slender prisms. $\text{CsCu}(\text{SCN})_2$ (R. T. Roberts): brilliant colorless prisms, stable in the air, decomposed by water with separation of cuprous thiocyanate. *Cs-Ag thiocyanates* (H. L. Wells). $\text{Cs}_3\text{Ag}(\text{SCN})_4$: large, transparent, colorless crystals of octahedral aspect, but evidently not isometric, and stable in the air. $\text{Cs}_2\text{Ag}(\text{SCN})_3$: brilliant transparent colorless prisms, stable in the air. $\text{CsAg}(\text{SCN})_2$: prismatic crystals, more flattened and with higher adamantine luster than the last. CsSCN : crystals resemble those of the potassium salt in appearance, and contain no water; they are exceedingly soluble in water and are stable in the air (unless it is very damp); it was prepared from caesium carbonate and ammonium or barium thiocyanates. *K-Ag thiocyanates* (H. F. Merriam). $3\text{KSCN} \cdot \text{AgSCN}$: beautiful large transparent stout prisms which, after drying, gradually become opaque upon standing. $2\text{KSCN} \cdot \text{AgSCN}$: large transparent crystals, stable in the air. $\text{KSCN} \cdot \text{AgSCN}$: thin diamond-shaped plates. *Cs-Ca, Cs-Sr and Cs-Mg thiocyanates* (H. F. Merriam). $2\text{CsSCN} \cdot \text{Ca}(\text{SCN})_2 \cdot 3\text{H}_2\text{O}$: colorless transparent prisms, stable in the air. $\text{Ca}(\text{SCN})_2 \cdot 4\text{H}_2\text{O}$: crystallized calcium thiocyanate carries four molecules of water, instead of three as reported by Meitzendorff (*Pogg. Ann.*, 56, 63 (1842)); it crystallizes in long prisms, very hygroscopic, and becomes opaque on standing over concentrated sulphuric acid from change to $\text{Ca}(\text{SCN})_2 \cdot 2\text{H}_2\text{O}$. $2\text{CsSCN} \cdot \text{Sr}(\text{SCN})_2 \cdot 4\text{H}_2\text{O}$: colorless transparent prismatic crystals with rectangular faces, and more soluble than the calcium

salt. $2\text{CsSCN} \cdot \text{Mg}(\text{SCN})_2 \cdot 2\text{H}_2\text{O}$: clear colorless hygroscopic crystals. This salt differs in type from all the known halides of magnesium. Cs-Zn and $\text{Ag-Zn thiocyanates}$ (H. L. Wells). $\text{Cs}_2\text{Zn}(\text{SCN})_4 \cdot 2\text{H}_2\text{O}$: small colorless crystals, moderately soluble in water, stable in the air. $\text{ZnAg}_2(\text{SCN})_4$: small, brilliant, colorless prismatic crystals, stable in the air, but decomposing when boiled with water. Ba-Ag , Sr-Ag and $\text{Cs-Ag thiocyanates}$ (H. L. Wells and H. F. Merriam). $\text{Ba}(\text{SCN})_2 \cdot 2\text{AgSCN} \cdot 2\text{H}_2\text{O}$: colorless prisms. $\text{Sr}(\text{SCN})_2 \cdot 2\text{AgSCN} \cdot 2\text{H}_2\text{O}$: colorless prisms. $\text{Ca}(\text{SCN})_2 \cdot 2\text{AgSCN} \cdot 2\text{H}_2\text{O}$: plates. The three latter salts are all stable in the air. $\text{CsTl}(\text{SCN})_3$ (H. L. Wells): small fern-like growths of crystals, faint yellow when dry, and stable in the air. This 4 : 1 type is unknown among the double halides with univalent metals. $\text{Cs}_3\text{BaAg}_2(\text{SCN})_7$ (O. G. Hupfel and H. L. Wells): large transparent tetragonal crystals resembling apophyllite, stable in the air, but readily decomposed by water; 100 parts water at 19° dissolves about 92 parts of the triple salt. $\text{Cs}_3\text{BaCu}_2(\text{SCN})_7$ (H. L. Wells): colorless, transparent crystals similar to the last, stable in the air, decomposed by water and cannot be recrystallized from it. $\text{Cs}_3\text{SrAg}_2(\text{SCN})_7$ and $\text{Cs}_3\text{SrCu}_2(\text{SCN})_7$ (H. F. Merriam) closely resemble the corresponding barium salts in every respect. The last four compounds are isomorphous. $\text{Cs}_3\text{CaAg}_2(\text{SCN})_7 \cdot 2\text{H}_2\text{O}$ and $\text{Cs}_3\text{MgAg}_2(\text{SCN})_7 \cdot 2\text{H}_2\text{O}$ (H. F. Merriam) may be recrystallized from water; both form colorless transparent crystals, the magnesium salt being the less soluble of the two, and the calcium salt having about the same solubility as the preceding barium and strontium triple compounds. $\text{Cs}_2\text{MnAg}_2(\text{SCN})_6 \cdot 2\text{H}_2\text{O}$ (C. S. Leavenworth and H. L. Wells): slender needles of faint greenish tint, stable in the air, sparingly soluble. $\text{Cs}_2\text{NiAg}_2(\text{SCN})_6 \cdot 2\text{H}_2\text{O}$ and $\text{Cs}_2\text{NiCu}_2(\text{SCN})_6 \cdot 2\text{H}_2\text{O}$ (R. T. Roberts and H. L. Wells): small crystals, stable in the air, less soluble in water than the corresponding calcium, magnesium and manganese salts containing caesium and silver, and decomposed by much water. The silver salt is sapphire blue, the copper salt a greenish blue. $\text{Cs-Ag-Zn thiocyanates}$ (H. L. Wells). $\text{CsZnAg}(\text{SCN})_4 \cdot \text{H}_2\text{O}$: slender colorless flattened crystals, brilliant when removed from the mother-liquor, but quickly losing water in the air, becoming opaque and brittle. $\text{Cs}_2\text{ZnAg}(\text{SCN})_5$: slender white needles of rectangular section, stable in the air. $\text{CsZn}_2\text{Ag}_3(\text{SCN})_8$: small thin colorless scales, usually of rhombic or rhomboidal outline, stable in the air, slowly decomposed by cold water, more rapidly by hot. Compound crystals of different habit were also observed. $\text{CsZn}_2\text{Ag}_4(\text{SCN})_9$: brilliant flattened colorless crystals, which vary greatly in habit, stable in the air, and slowly decomposed by water. The presence of ammonium thiocyanate appears to be necessary for the formation of this com-

pound. All the above CsZnAg salts are too insoluble to be recrystallized from water without decomposition. $\text{K}_4\text{BaAg}_3(\text{SCN})_8 \cdot \text{H}_2\text{O}$ (H. L. Wells): truncated rectangular pyramidal crystals, brilliant and transparent, stable in the air, very soluble in small amounts of water, but decomposed by much of it with precipitation of silver thiocyanate. Most of the compounds described above were obtained by mixing the different thiocyanates in varying proportions and under different conditions, and then evaporating to crystals.

M. T. BOGERT.

MINERALOGICAL AND GEOLOGICAL CHEMISTRY.

A Quantitative Chemico-Mineralogical Classification and Nomenclature of Igneous Rocks. BY WHITMAN CROSS, JOSEPH P. IDDINGS, LOUIS V. PIRSSON AND HENRY S. WASHINGTON. *Jour. of Geol.*, 10, 555-690.—This most elaborate paper represents the combined labor of the authors, with whom the late Professor George H. Williams was originally associated, for quite a number of years. All attempts to make any of the existing systems meet the demands of modern petrology having failed, an entirely new one has been evolved which has been tested by the calculation of thousands of analyses. "Its concepts of rocks are in a large measure new, and hence, except in a very small degree, it demands a new nomenclature. . . . All igneous rocks are classified on a basis of their *chemical composition*; all rocks having like chemical composition are grouped together. The definition of the chemical composition of a rock and of a unit of classification is expressed in terms of *certain minerals* capable of crystallizing from a magma of a given chemical composition, and the expression is quantitative." It is quite impossible to give, in the limits of an abstract, an adequate idea of the nature of the proposed system and of the arguments offered in its behalf, therefore the reader must be referred to the original, which will later be reprinted in book form by the University of Chicago Press, together with a glossary of new words and their definitions and tables to aid in calculations.

Such a revolutionary step will doubtless meet with strenuous opposition, especially from the more conservative, but it is to be hoped that the spectacle of several leading petrologists united in harmonious advocacy of this radical change may have its effect in securing for the system not only approval but its gradual adoption, and thus end a most vexatious situation, quite as troublesome as that confronting chemistry.

W. F. HILLEBRAND.