"The sun's rays are the ultimate source of almost every motion which takes place upon the surface of the earth. By its heat are produced all winds and those disturbances in the electric equilibrium of the atmosphere, which give rise to the phenomena of terrestrial magnetism. By their vivifying action vegetables are elaborated from inorganic matter, and become in their turn the support of animals and men, and the sources of those great deposits of dynamical efficiency which are laid up for human use in our coal strata. By them the waters of the sea are made to circulate in vapour through the air and irrigate the land, producing springs and rivers. By them are produced all disturbances of the chemical equilibrium of Nature, which by a series of compositions and decompositions give rise to new products and originate a transfer of materials."

In a note in Mr. Herbert Spencer's "First Principles" (2nd Ed., p. 496), which first led me to look at this passage, it is remarked that Herschel "expressly includes all geologic, meteorologic, and vital actions, as also those which we produce by the combustion of coal," in the effects of the solar rays. When, therefore, Prof. Tyndall states that Mayer *revealed* the source of the energies of the vegetable world, it appears to me that Herschel anticipated the revelation twelve years previously. Of course, I apprehend that Mayer's merit consisted in seizing at once a physical principle of immense generality, and in applying it to very different phenomena. Herschel began at the other end; but appears equally to have seen the solar energy underlying these phenomena, though in a general way, and without demonstrating numerical relations.

2. De Saussure is credited unreservedly with the observation of the reducing power of the solar rays in the vegetable economy. But he seems to me, as, indeed, he seemed to himself, to have only crowned a theory which other workers had elaborated. Priestley began by ascertaining that air depurated by animals was purified by plants. Ingenhousz showed, what Priestley candidly confesses he missed, that this effect is due "chiefly, if not only, to the light" of the sun. Senebier found that "fixed air" was the ingredient which plants removed from a vitiated atmosphere, and that this underwent elaboration in the tissues, oxygen being set free as the result.* Lavoisier having previously shown that fixed air was a compound of carbon and oxygen, Senebier's results implied the fixation of carbon by plants. This fixation De Saussure actually demonstrated by feeding a plant with carbon dioxide and water alone, and showing that the carbon in the tissues increased. He further found the unexpected fact (and this is what he added to the matter) that the oxygen evolved by plants does not correspond to that contained in the carbon dioxide absorbed, but that it is smaller in quantity.

De Saussure's researches are a beautiful example of quantitative work, but they would have, I imagine, merit of a different order if Priestley, Ingenhousz, Senebier, and Lavoisier had not broken ground before them.

W. T. THISELTON DYER

Phenomena of Contact

IN NATURE of August 24 I objected to, as misleading, the statement by Mr. Newcomb that "we find ligaments, black drops, and distortions sometimes seen in interior contacts of the limbs of Mercury or Venus with that of the sun, described as if they were regular phenomena of a transit, without any mention of the facts and experiments which indicated that these phenomena are simple products of insufficient optical power and bad definition which disappear in a fair atmosphere with a good telescope well adjusted to focus." I asked for references to the facts and experiments by which the statements are justified.

IN NATURE of September 28 I find Mr. Newcomb's reply, but without the references which I desired. Mr. Newcomb considers that I controvert the two following propositions:—

1. That the irregular phenomena of internal contact of a planet with the sun, variously described as distortions, black drops, ligaments, &c., are not always present, but are only seen sometimes.

2. That when seen they are due to insufficient optical power or bad definition.

If the word "irregular" is cut out, and the word "seen" substituted for "present" in proposition (1) there can be no doubt about its truth. It will be found that all the arguments adduced

by Mr. Newcomb to prove this proposition have no bearing either upon the word "irregular" or "present" in contradistinction to "seen."

It appears to me, therefore, quite unnecessary to allude further to this proposition.

With reference to proposition (2), I believe it to be utterly erroneous. I believe that the phenomena of the fine connecting ligament can only be seen in a fair atmosphere, with a good telescope well adjusted to focus, and with considerable magnifying power. When it is remembered that the fine connecting ligament is confined to within about a second of arc of the sun's limb, I think my statement will at least commend itself to practical observers. Mr. Newcomb appears to regard it as a great difficulty in my view of these phenomena, that some of the observers should see the ligament and some not. I am rather surprised at the persistence with which this point is again and again brought forward in his letter. I thought that it had been answered by anticipation in my letter which appeared in your number of August 24. In all my writings upon the subject I have maintained that the phenomena could only be seen under favourable circumstances and with sufficient power; and in my letter of August 24 will be found this statement, which appears to have been entirely overlooked, at least unanswered, by Mr. Newcomb:—"The optical enlargement by irradiation is a function of the brightness, and can be made insensible by sufficiently diminishing that brightness. Unfortunately, however, when this diminution of brightness is carried to a very great extent, errors in an exactly opposite direction to those of irradiation will come into play, similar, in fact, to the results of Wolf's experiments. The observations of Mercury on the sun's disc in 1868 were made with very different optical means, and some very different methods were adopted for diminishing the sun's glare." In my view those observers who did not see the connecting ligaments failed to see it, either from want of attention to the point as not a contact such as they expected to see, or from the observations having been made under such circumstances that some of the necessary conditions which I have indicated were not satisfied.

The fine connecting ligament is only seen by contrast against the illumination of the sun's disc near the point of contact, and it may well be that some of the observers have pushed the diminution of brightness of the sun's image to such an extent that the contrast was too feeble to attract attention before the apparent contact. To me, and I think to others who will give the matter some consideration, it is clear enough "that an observer with the naked eye, a telescope of low power, would not, in the case of a transit of Venus, see the connecting ligament at all." It is as clear "that without seeing any ligament, the planet, at egress, would appear to touch the limb without distortion, necessarily, earlier than the contact would appear to be established to observers who were watching the transit with good telescopes and with high powers." It appears to me equally clear that, if the brightness of the sun's image be reduced to excess, then the ever-diminishing small portion of the illuminated disc between the sun's edge and the advancing planet, at egress, may be made to disappear, from sheer inability to appreciate so faint a light, before the contact would appear to be established to observers who had not so reduced the brightness of the image. Disturbing causes such as these do exist, and their effects must be recognised. I must apologise for bringing forward such arguments; but, since one observer has published his opinion that the "ligaments, &c.," do not exist in contacts, because he looked at the transit of Mercury with an opera glass and saw nothing of the kind, it would appear necessary to recall attention to a common-sense view of the points at issue.

But whatever may be the opinion of Mr. Newcomb respecting the explanation which I have given of the probable reason why some of the observers have not seen the connecting ligament, he must feel that it will at least be difficult for him to explain away the positive evidence of the numerous observers who profess to have seen the ligament with first-rate telescopes. Some of them, at least, were gentlemen not likely to have forgotten to adjust their eye-pieces to focus, even if such a neglect would have produced the phenomena observed. In cases where the ligament has been seen, it will be found that the earlier lines of contact at egress have been given by observers with the best telescopes and high powers. This is strikingly shown by the Greenwich observations of the transit of Mercury, 1868. It is a result in perfect accord with my views. Very large and systematic discordances will be found to exist between the times of internal contact at the transit of Mercury, 1868, in cases where no connecting ligament was seen at all. This has been passed over in

* Recherches sur l'influence de la lumière solaire pour métamorphoser l'air fixe en air pur par la végétation, 1793.

silence by Mr. Newcomb; but it is important. It would be difficult to select from such groups of observers—the French, for example, who saw no connecting ligament—those who saw “the phenomena exactly as we know they are;” and unless this can be done, I am afraid that Mr. Newcomb’s somewhat unique argument upon this point might be made as easily to prove the converse as the result he deduces from it. All these observations, in my view, are good; but they are not strictly observations of the same phenomena.

Mr. Newcomb rejects at once the force of the evidence of the observers of the transits of Venus, 1761 and 1769, upon the question of the connecting ligament “till we have better evidence than now exists that their object-glasses were such as Clarke or Foucault would call good.” The phenomena connected with the ligament must be far more marked in the case of Venus than in that of Mercury, on account of the large diameter of Venus. To reject therefore by an impossible condition all the evidence in our possession respecting transits of Venus is certainly a bold step; but Mr. Newcomb appears to me to attach far too much importance, so far as irradiation phenomena are concerned, to the improvements effected in modern telescopes. The image of a point of light on the most perfect object-glass which can be conceived is not a point, but a disc, of which the illumination degrades rapidly from the centre, and which is surrounded by concentric rings of light. The law of degradation of the illumination of the central disc has been given by me in the Monthly Notices, November 1865. The result of theory upon these points has been most completely tested by experiment. The existence and regularity of these concentric circular rings is one of the most delicate tests of the perfection of a telescope. Since we have a disc of light corresponding to a point in the most perfect object-glass which can be made, the visible image of the sun formed by such a glass will not terminate with the geometrical image. This result of theory is confirmed by experiment. The optical enlargement found under degrees of illumination similar to those very commonly adopted in observations of the sun is amply sufficient to produce by its destruction near the point of contact the phenomena which so many observers of experience have declared that they have seen. That the optical enlargement is sufficient for the purpose can be seen from the experiments of Dr. Robinson, of Armagh, and from the Greenwich discussions of eclipse observations. This was pointed out in my letter in your number of August 24. With respect to Mr. Newcomb’s remark as to the application of this theory of irradiation to a transit of a planet, viz., “we require to know whether the irradiation of an extremely minute thread of light darkened so as to be barely visible is the same with that of a large disc, I am decidedly of opinion that it is not, and if not, the fact that the sun’s disc is optically enlarged by the telescope or the eye of the observer cannot be directly applied to the phenomena of transit.” I have merely to remark that Mr. Newcomb is undoubtedly right when he asserts that the irradiation from the minute thread of light darkened so as to be barely visible is not the same as that of the large disc. *It is simply because such is the case that the phenomena of the connecting ligament appear.* When the planet is well on the disc, the irradiation around the disc will not be disturbed, but as the planet approaches the edge, the irradiation near the point of contact must eventually be disturbed, and this disturbance, or change, gives rise to the phenomena observed—a black drop, connecting ligament, or whatever name you prefer to give to that apparent cutting out of a piece of the sun’s edge near the point of contact which must take place. After the disturbance of the irradiation has once commenced, the connecting ligament must at egress increase in breadth; but I do not profess to be able to give the law of the changing form with any degree of exactness.

The experiments of Wolf and André were, as I stated in my letter of August 24, made upon a disc presenting no sensible traces of optical enlargement. The results can therefore have no bearing upon the question of irradiation. These results are undoubtedly valuable in themselves, as showing experimentally the tendency of errors of observations of contacts under feeble illumination. They may throw light upon those observations at which no connecting ligament was seen, but they are useless to disprove or prove irradiation effects.

My authority for stating that the observations of Wolf and André were made upon a disc showing no sensible traces of optical enlargement, is contained in the memoir itself. If Mr. Newcomb is pleased to call the phenomena of “telescopic irradiation” a species of bad definition, there can be no objection on my part

to his doing so; but it is not a species of bad definition “which disappears in a fair atmosphere, with a good telescope well adjusted to focus.”

With respect to the ligament not being a celestial reality. The contact is not a celestial reality. My views of the reality of the phenomena are that the reality is neither more nor less than the reality of the phenomena presented at the focus of an object-glass when turned upon a star. The irradiation can in my view be got rid of to the same extent and in the same manner that the central disc corresponding to the star’s image can be got rid of. You can reduce its dimensions by cutting down the illumination, and the disc will become a point, but only as it vanishes.

If I may be allowed to give one word of warning respecting the preparations for the transit of Venus 1874, it is *uniformity*. Make such arrangements as you think best, but once made stick to them even if not absolutely the best. The observations which are to be compared must be made as early as possible under the same optical conditions. The whole success or failure of the work will, in my opinion, turn upon the extent to which this necessary condition is approximated to.

E. J. STONE

Royal Observatory, Cape of Good Hope, Nov. 18, 1871

The Origin of Insects

WITH your kind permission I will answer Dr. Beale’s questions, published in his letter in your issue of December 21, 1871.

Dr. Beale asks me what part of the nervous system of the maggot is present in the fly? My answer is that I have traced the changes of the one directly into the other; and that Weismann has done the same. There is no time in the pupa state when the nervous system is absent; but it is difficult to demonstrate this, as amongst so much molecular matter it is not easily found, and is very easily crushed and destroyed.

Again, Dr. Beale asks me if I have seen any vestige of the mouth organs of the imago in the larva? I reply that the mandibles and maxillæ exist in the egg twelve hours before the young maggot emerges, together with the fore and hind-head segments; that these have all disappeared when the egg hatches; but that the imaginal discs are already formed at that time. Now, I would ask if it bears the slightest aspect of probability that the larval head segments and mandibles, maxillæ, &c., are formed for nothing, and that the imaginal discs are new formations arising contemporaneously with the disappearance of the larval head segments? Dr. Weismann has shown unmistakably that the abdominal segments of the pupa skin are formed from the abdominal segments of the larval skin. Does it appear in the slightest degree probable that the thoracic and head segments have a totally dissimilar origin? I admit that I have not been able to see the imaginal discs in contact with the head segments of the embryo; but I have found the imaginal discs immediately after the egg is hatched, and they are then too much like the embryonic structures alluded to, to have had any other probable origin. The proboscis is formed from cells laid down within these discs; of that there is not the smallest doubt. Dr. Weismann makes the same assertion, and, although I did not know it to be so at the time I wrote my work on the fly, I acknowledge it is so now, and that in my description of the origin of the proboscis I was wrong. In the Lepidoptera, and in some beetles, imaginal discs may be seen to have their origin in the inner layer of the larval skin.

Again, Dr. Beale says:—“Does Mr. Lowne mean to say, for instance, that he or any one else can adduce any reliable observations to prove that the sexual organs are gradually developed, even from the time when the embryo is enclosed within the egg?” I answer, yes. My own observations confirm those of Weismann on this head, and Dr. Beale will find, on looking again at page 112 of my book on the blow-fly, that he has not correctly quoted my statement. I will also refer Dr. Beale to Dr. E. Bessel’s paper, “Studien über die Entwicklung der sexual-Drüsen bei den Lepidopteren,” in the “Zeitschrift für wissenschaftliche Zoologie,” vol. xvii. I believe future observers will find the sexual organs are always so formed, even as they are in the vertebrata. There is another paper, by Siebold I think, on the same subject in the above-quoted periodical.

Lastly, Dr. Beale asks me to explain what I mean by the sentence occurring at page 116 of my book:—“All the tissues of the larva undergo degeneration, and the imaginal tissues are redeveloped,” &c. I apprehend that the redevelopment of all the