Original Articles.

A STUDY OF THE DIFFERENT APPROACHES TO THE HIP-JOINT, WITH SPECIAL REFERENCE TO THE OPERATIONS FOR CURVED TROCHANTERIC OSTEOTOMY AND FOR ARTHRODESIS.

BY E. G. BRACKETT, M.D.,

Surgeon to the Orthopedic Department, Massachusetts General Hospital.

THE hip-joint has perhaps been the last to be freely approached since surgery of the joints has become legitimate, probably because of its naturally strong barriers, which have more successfully guarded the different avenues of approach. The hip-joint presents its special problems because of conditions which differ from those in nearly all other joints, and in these respects:

(1) It is the deepest joint in the body.

(2) It is covered on all sides by more than one layer of muscles, the edges of which so overlap as to prevent any direct entrance without either section or very marked retraction. A most important factor is the fact that many of these muscles are short and cannot be retracted, or separated throughout their whole length, without the danger of injury to their nerve supply, a very probable incident in the attempt to reach a joint so deep.

(3) It is not possible to expose more than a limited area in the neighborhood of the hip-joint without serious damage to its essential coverings, and even when an avenue of approach to the joint is clearly opened, the joint itself is still hidden, and cannot be opened without damage to its intrinsic articular structures.

The factor of importance, then, is to effect the approach as completely as possible by such intermuscular route, that the desired part of the joint may be reached without injury to the muscles, and with the least retraction possible, in order to avoid injury to the smaller nerve branches which supply these muscles at their points of entrance. In most joints there is but little possibility of this injury, the nerve supply being remote, but in the hip-joint the danger is decidedly imminent. It becomes important to choose carefully the entrance of approach to this joint, and to select the one best adapted for the special purpose in each case, for the reason that only a small portion of the joint or its surrounding structures can be exposed by any route. One must have as many routes from which to choose as possible, with an accurate knowledge of the area which each will best expose.

The routes or the methods of approach which are considered in this paper are not new; they are all old and known, but accurate information of surgical aid is difficult to find. Dissections which have been carried on in connection with this subject were made to determine the surgical anatomical guides in the approach to the hip-joint, in order to enable one to find the way with the least damage to the parts while en route.

Two of these, the anterior and the latero-posterior, have been studied in special detail with reference to special operations for which they may be used.

It is possible to choose between many routes, but among the most practical are the following:

Inner route: The incision, made from one inch to the inside of the anterior superior spine directly downwards six inches, discloses the inner border of the sartorius muscle (Fig. 1, C-D) attached to the fascia covering the vessels. The external cutaneous nerve lies near to the outer edge of the fascia; the external saphenous vein lies under the superior layer and to the inner side of the nerve; to the inner side and deeper lies the anterior crural nerve. Dissection downward along the dividing edge of the sartorius and the fascia, and retraction of the sartorius to the outside (Fig. II), opens to the inner side of the upper portion of the rectus, covered by a fold of the same fascia. Incision through this opens to the line dividing the inner edge of the rectus and the outer edge of the iliacus; the vessels and anterior crural nerve are near but to the inner side, the nerve lying in this fascia a little to the inside and below the external cutaneous nerve and with the large vessels immediately in the inside neighborhood. In the lower region of the wound, between the rectus and iliacus, are the small but unimportant circumflex vessels. The floor of the field is made by the iliacus muscle, which lies directly on the capsule of the inner portion of the neck.

From this point, the way may diverge and two choices are open:

(1) The outer — between the iliacus and rectus; the separation of these two opens immediately to the capsule, just at the inner side of the neck at the point of its junction with the head. A portion of the Y ligament of the capsule is in the bottom of the field. To reach the upper anterior edge of the acetabulum, the dissection must be carried high, nearly to the level of Poupart's ligaments.

(2) The inner — retracting the fascia still further to the inside, along the surface of the iliacus, the dividing line of the iliacus and of the psoas tendon is reached. These are easily separated, and between them lies the anterior edge of the head and the anterior rim of the acetabulum.

The first opens up the anterior capsular cavity. The second is too near the capsular attachment, and above the acetabulum rim, to be of much use in the exploration of its cavity. The field is so restricted in either that their use is mainly in an attempt to inspect the anterior portion of the articular head, or of the neck at its junction with the head, which can be done by rotation of the leg, with the capsule relaxed by flexion of the thigh.

In a conservative operation the capsule may be opened and the anterior part of the neck, the edge of the head and the rim of the acetabulum examined, but the restricted field prevents any extended exploration.

Anterior route: The incision, made from just

below the anterior superior spine directly downward five or six inches, exposes the line of division between the sartorius and tensor fascia femoris (Fig. I, A-B). At the upper portion, the fibers of the sartorius are nearly vertical, and parallel to the fibers of the tensor fascia femoris, but about three inches below its origin, the outer edge of the muscle takes a rather sharp bend inward. The two muscles are easily separated, and this separation must be carried high, nearly to the point of their origin, where the two muscles are closely blended, having their origins much in common. Retracting the sartorius to the inside, the upper portion of the rectus is seen lying in the interval between the two (Fig. III). Retracting the sartorius and rectus to the inside, and lifting at the same time the outer edge of the iliacus (Fig. IV), exposes the outer portion of the capsular cavity. Through this opening the anterior surface of the neck and the edge of the head may be brought into view, particularly when the thigh is flexed. Retracting the sartorius and rectus to the inside and the tensor fascia femoris to the outside, exposes the anterior surface of the trochanter and outer portion of the neck (Fig. V). In the lower portion of this area are found the small circumflex vessels imbedded in the intermuscular fat. On the inner side, lying on the neck, is the outer edge of the iliacus, passing to the lesser trochanter; in the lower and outer portion of this area are the upper fibers of the vastus externus, extending to the middle of the trochanter in their attachment to the femur. The muscles on the outer side are easily retracted so as to expose the outer edge of the trochanter; and on the inner side, the edge of the iliacus is lifted and retracted so as to bring into view the lesser trochanter. This field, when retraction is made to the outside, gives free access to the anterior surface of the femur from the outer border of the middle of the trochanter to the junction of the neck and shaft, just above the lesser trochanter (Fig. V).

Lateral route: The lateral incision (Fig. VI, A-B-C) is made from the anterior superior spine, obliquely downward and outward, to the middle of the outer side of the trochanter, and is then turned downward two inches, in the line of the femur. In the center of the field is the dividing line between the tensor fascia femoris and gluteus medius (Fig. VII) covered by the upper portion of the fascia lata, which, at this point, is dense, so that the line of muscle division cannot be seen through the fascia, and is often not easy to find even after opening the fascia. Two anatomical facts are here of aid: (1) The upper portion of the tensor fascia femoris is entirely free from the fascia, while the fibers of the gluteus medius are attached to the under surface of the fascia. (2) The gluteus medius is a coarser muscle than the tensor fascia femoris, and, therefore, the difference in grain is a distinguishing mark. The upper portion of the tensor fascia femoris is clearly defined and band-like; the lower part is ill defined and begins to be attached to and merged into the fascia lata; its lower edge overlaps

the anterior edge of the gluteus medius, the fibers of which run in a more oblique direction downward. The lower portion of the gluteus medius begins to get fascial, merging into the fascia of its attachment to the trochanter; therefore, the easiest place so find the dividing line is midway between the anterior superior spine and the trochanter.

Underlying the tensor fascia femoris, seen when looking toward the joint from the lateral aspect, is the upper part of the rectus in its musculo-tendonous portion (Fig. VIII), which lies between the inner edge of the tensor fascia femoris above and the neck of the femur below. Carrying the separation of these muscles high, the fibers of the horizontal extension of the tendon of the rectus are seen, just above the upper and anterior edge of the acetabulum, and are a guide to the joint. Elevation of the upper portion of the rectus with the tensor fascia femoris discloses the outer edge of the iliacus, in its passage over the neck to the lesser trochanter.

The area exposed is the upper part of the trochanter and the neck; inward to the rim of the anterior and upper edge of the acetabulum; downward to the lesser trochanter; and upward to the level of the tip of the greater trochanter. The portion of the capsule, and the upper side of the neck and the portion of the acetabular rim corresponding to it, cannot be reached by this route with sufficient freedom for any extensive surgical procedure. The capsule in the exposed region is in view, and by the flexion of the thigh is relaxed and the cavity opened with ease. The head of the femur can be rotated so as to expose a portion of the cartilaginous head, enough to determine its character, but the field of operative procedure is limited to this smaller area, except by extensive myotomy.

Latero-posterior route: The first part of the incision and dissection is identical with the lateral incision (Fig. VI, A-B-C). Especial care must be taken to end the vertical portion well up on the outside of the trochanter (1½ inches from the tip) and well down towards its posterior border. At the point where the oblique portion joins the vertical, just over the trochanter, a right angle incision is made directly backward (Fig. VI, B-D), two to three inches, down to the fascial portion of the gluteus maximus. After separating the tensor fascia femoris and the gluteus medius, the line of separation is extended downward along the line of the original incision, through the fascia lata, to the femur, freeing the attachments of the muscles (vastus externus) from the outer and upper surfaces of the femur. The fascial expansion of the gluteus maximus is then cut through, along the line of the posterior part of the original incision, and the outer part of the trochanter exposed. This gluteal flap (Fig. IX, A-B) is turned backward, and the outer and upper surfaces of the trochanter fully exposed, this being necessary for the later wiring. The outer and upper surface of the trochanter is then chiseled off by cutting directly backward with a narrow osteotome, on a curved line (Fig. IX,

FIG. I. Showing the general relation of the joint to the overlying muscles. A and B=the line of incision for the anterior route. C and D=the inner route.

FIG. II. Second step of the inner route, which is continued by (1) between the rectus and iliacus; or (2) between the iliacus and tendon of the psoas.

FIG. III. Second step in anterior route. Separation of tensor fascia femoris and sartorius shows rectus, with edge of iliacus, overlying the trochanter and outer portion of neck.

FIG. IV. Showing separation of muscles for opening field for operation on trochanter and outer neck. The dotted line indicates the position of the line for curved Gant.

The drawings for this article have been made by Miss A. J. Ripley; the anatomical plates mainly from photographs of dissections; the others from tracings of x-ray negatives.

FIG. V. Last step in anterior route in opening capsule.

FIG. VI. Showing line for lateral and latero-posterior routes of incision with the relative position of underlying head and neck of femur. A, B and C=line of lateral incision. B and D=posterior part of latero-posterior incision.

FIG. VII. Lateral route—first stage. Fascia turned backward showing position of line of division of tensor fascia femoris and gluteus medius. In the lateral incision the posterior cut is not made.

FIG. VIII. Lateral incision—second stage. Gluteus medius with fascia attached to lower portion retracted backward. Tensor fascia femoris retracted forward, showing tendinous portion of rectus lying on iliacus, the outer edge of which is seen.

FIG. IX. Latero-posterior incision — third stage. The fascial flap of the gluteus maximus has been cut in the line of the posterior incision and turned back (A and B), which exposes the bone, ready for the removal of the lip of the trochanter.

FIG. X. Latero-posterior incision — last stage. Tip of trochanter removed and turned backward and upward (A) with the attachment of the gluteus medius and minimus and pyriformis; (B) outer surface of trochanter, from which (A) is removed. Anterior muscles retracted. The upper part of the neck with its capsule is in the center of field. Capsule opened longitudinally.

FIG. XI. Showing edge-to-edge apposition in subcutaneous osteotomy.

FIG. XII. Showing splintering in subcutaneous osteotomy and slipping by of lower fragment. (Case I.)

FIG. XIII. Showing splintering in subcutaneous osteotomy. The fracture occurred during the osteotomy and the fragment was a long spiral, the anterior surface of the upper fragment overhanging. The osteotomy was entered about two inches above the lower point of the overhanging fragment. (Case II.)

FIG. XIV. Showing moderate degree of slipping.

FIG. XV. Showing slipping by of fragment with loss of length.

FIG. XVI. Showing result of old osteotomy (five years previous). Good functional result, but alignment bad and some shortening resulting.

FIG. XVII. Showing slipping of lower fragment and bad alignment. The osteotomy is too low, but the deformity of the trochanter is a misleading landmark.

FIG. XVIII. Showing line of curved Gant with lower fragment turning securely in hollow of upper. Gives alignment of weight bearing through the joint. The curved bone incision may be made lower if there is reason for keeping farther from the joint.

Fig. XIX. Curved osteotomy intertrochanteric. This case represents a type not suited for osteotomy. Ankylosis is not complete, but motion is not enough to be of practical value. Arthrodesis of the joint would have been preferable.

Fig. XX. Typical case for osteotomy. X-ray taken ten weeks after curved osteotomy, showing line of incision. Alignment is through the joint. Slipping of fragments is impossible.

Fig. XXI. Marked hypertrophic arthritis. Joint practically stiff and in deformity. Would be a typical case for arthrodesis if other joints were not involved. (Case III.)

Fig. XXII. Monarticular arthritis after trauma. Leg practically useless for weight bearing. Type of case for arthrodesis. (Case V.)

Fig. XXIII. Showing result several months after arthrodesis. Trochanter had been removed and wired in place. Union of joint and trochanter both firm. (Case IV.)

Fig. XXIV. Showing type of old hip disease for arthrodesis. Ankylosis is not complete and joint subject to frequent strain. Trochanter impinges on ilium. (See also Fig. XIX.)

FIG. XXVII. Showing old infectious arthritis (in childhood) with incomplete ankylosis; marked overgrowth and enlargement of neck. Typical case for arthrodesis. (Case VII.)

FIG. XXX. Case of Fig. XXIX after operation. Trochanter removed and wired in place. One-half motion with good weight-bearing leg.

FIG. XXVI. Case of Fig. XXV after operation. Upper part of trochanter removed in order to reach joint freely.

FIG. XXIX. Old dislocation (Case XVI) with injury to head. Arthrodesis not necessary. (Fig. XXX.)

FIG. XXV. Showing typical case for arthrodesis in old hip disease without complete ankylosis. (Case XII.)

FIG. XXVIII. Infectious arthritis with leg in marked flexion and adduction. Ankylosis not complete. Typical case for arthrodesis. (Case XIII.)

small figure) beginning on the outer surface of the trochanter, $1\frac{1}{2}$ inches below the tip, and cutting inward for one quarter of an inch, then curving upward to the fossa at the junction of the upper part of the neck and the trochanter. This removes the outer portion and tip of the trochanter, and with it the attachments of the gluteus medius, gluteus minimus and piriformis. Care should be taken in the removal of this piece not to encroach on the neck, or the bone will be weakened at this angle. The portion of the bone removed, with the muscles attached, is deflected backward and upwards (Fig. X, A) enough to uncover the upper part of the femoral neck, and the muscular covering of the anterior fibers of the gluteus medius and minimus are easily separated from the region above the acetabular rim. The incision through the trochanter just borders on or opens the outer edge of the capsule on the upper surface of the neck, and incision of this opens directly into the cavity of the joint. The capsule may be split along its upper surface, parallel to the neck, and near to its acetabular insertion cut transversely on each side, which opens the view of the edge of the head and the rim of the upper half of the acetabulum. This route gives the most extensive view of the hip and its surrounding structures, and brings into view, within the capsular cavity, the upper part of the neck, the portion of the head outside of the acetabular cavity, and the upper half of the rim of the acetabulum and cotyloid ligament. It is the best approach for any of the more extensive explorations and operations on the joint, particularly for the arthrodeses. By rotation of the femur, the extensive opening allows the largest amount of the articular surface of the head to be brought into view, and thus in operation on the articular structures the opening gives a freer access than any of the other avenues of approach.

Posterior route: The incision is made over the upper part of the buttock, in a line extending from the junction of the middle and posterior third of the ilium to the middle of the trochanter. The subcutaneous fat must be well separated from the underlying fascia, and the field then brings into view the anterior edge of the gluteus maximus, blending into the fascia lata, which along its upper portion is closely adherent to the anterior border of the gluteus maximus, and does not give a sharp line of demarcation. The gluteus medius lies beneath that anterior portion of the fascia which is exposed, but its posterior edge underlies the gluteus maximus. To reach this border, the gluteus maximus must be strongly retracted backward, or a few of the anterior fascial fibers of attachment cut and deflected backward. The posterior edge of the gluteus medius and minimus practically overlie each other; the tendon of the piriformis lies in contact with the posterior edge of the minimus, closely adherent to it. It is not necessary to disturb their relation to each other. Retraction forward of the posterior edge of the minimus discloses the capsule of the joint, which may be opened along the edge of the piriformis tendon. This divides

the capsule slightly more horizontally than the direction of its fibers, and opens the capsular cavity along the posterior and upper part of the neck and head. This route should seldom be taken for any procedure, excepting excision of the joint proper, or exploration of the posterior capsular cavity. It is the natural avenue for reaching the gluteal bursa of this region, which it is often necessary to reach in conditions of its primary involvement; which conditions closely resemble the early stages of hip disease.

The adaptability of two of these routes to special operative procedures has been especially studied in this series: (1) The anterior route for a modified Gant (curved Gant); and (2) the latero-posterior, for the arthrodesis of the hip-joint (Albee's operation).

(1) The anterior route for the curved Gant (curved osteotomy for correction of deformity at the hip-joint). The original method, and the one much used at present, subcutaneous subtrochanteric osteotomy, is not an operation of precision, and how incompletely this operation has met the needs is seen by the number of modifications which have been attempted. It was a necessity in the days of aseptic surgery, and it is interesting to see how this operation has intruded itself into this period of frank surgical methods. In addition to the lack of precision in surgical technique, the operation is unmechanical. In no other method of treating fractures would one be content to replace fragments in an edge-to-edge apposition, as in this form of osteotomy (Fig. XI), and with the forcible abduction in the correction, the chance of slipping by of the lower fragment is one of not only great probability, but of frequent occurrence, as the x-ray now shows. Moreover, the division at this site does not give us the proper alignment of weight bearing through the joint, a factor of particular importance, with the joint already weakened by disease, and when the division is made below the point of selection, as so frequently occurs, the distortion between the line of the shaft and the line of weight becomes greater. Whether or not this operation should be done in children may be an open question, but in adults (and it is an operation which is most essentially applicable to adult life) the conditions are so different as to require quite a different consideration. In adults the bone lies deeper (usually); the bone is much larger and more difficult of section; the contractions from long-existing deformity are greater and more difficult to overcome; and, what is most important, the bone is more brittle and prone to splinter. This last factor is, in itself, of sufficient importance to demand a method of operation of greater precision.

The danger from splintering is illustrated by two cases, — both adults, with firm ankylosis in adduction and flexion deformity. On both the subcutaneous intertrochanteric osteotomy was done. One broke (Fig. XII) after the usual amount of chiseling, and without more than the ordinary amount of force to complete the fracture. The other (Fig. XIII) broke during the cutting

of the bone. Both of the patients complained of an unusual amount of pain following the operation, and examination by the x-ray showed a transverse fracture, with a splintering of the anterior edge of the upper fragment, making practically a spiral fracture of this half of the shaft, and leaving an overhanging shelf with an oblique under surface, on which the lower fragment slid by. In one, the splintering extended three inches down the shaft of the bone; in the other, the splintering was much shorter; in each, the lower fragment rode upward and under the oblique surface of the upper. Both cases required subsequent operation for replacement and wiring together of the fragments. These two instances, which are extreme, illustrate the danger in the usual osteotomy of adult bone, and the occurrence of which, in a minor degree, frequently happens. The study of old cases by the x-ray reveals a decided amount of displacement, usually inward, very often associated with a loss in length, and in many instances this occurs where a supposedly good result had been obtained (Figs. XIV, XV, XVI, XVII).

Many operations have been advocated to avoid the disadvantages of the old method, but the majority of them have the disadvantage of sacrificing a fraction of the length. The operation planned and here described, and which has been used in the series here reported, most nearly resembles that of Mr. Robert Jones, which is most mechanically perfect, and does not give the objectionable shortening. The chief aim was to determine an open method, to obviate the disadvantages of the bad mechanics, and the danger of slipping by of the fragments. This may be designated as a Curved Gant by the open method.

For this Curved Gant the anterior incision is used to open to the anterior surface of the trochanter, and the bone is exposed from the outer side of the greater trochanter to the inner side of the neck, and its junction with the shaft (Fig. IV). It is necessary here to free this surface, which is easily done by lifting and retracting inward the iliacus muscle as far as the lesser trochanter. A blunt dissector is then placed vertically downward on the inner side of the bone, at the point of junction of the lesser trochanter with the neck, and here left in position. As a rule, it will not be necessary to disturb the attachment fibers of the vastus, which are found in the lower portion of the field. With a very narrow osteotome a curved incision is then made (Fig. IV, A-B) with the convexity upwards and inwards, beginning on the outer side of the trochanter, and ending at the point of junction of the neck and lesser trochanter, where the blunt dissector is in place. The section is made vertically downward (the patient lying on the back) from the anterior through the posterior surface of the bone. When the leg is abducted, in correcting the abduction deformity, the convex end of the lower fragment can be seen to turn in hollow of the upper fragment, with no possible danger of subsequent slipping by (Fig. XVIII).

The place on the outer side of the trochanter where the incision of the osteotomy is begun, depends on the amount of abduction to be overcome. If the degree of adduction be slight, the outer end of the curved osteotomy line is lower, about opposite the point of exit. If the degree of adduction be greater, the osteotomy line should be higher on the trochanter. If there be a large degree of flexion the vertical (anteroposterior) line of osteotomy may be deflected slightly backward, so as to give a slight overhang of the anterior edge of the upper fragment.

The advantages of the method are:

- (1) It insures the impossibility of slipping by of the fragment.
- (2) It allows of an adjustment of the line of weight bearing and the line of the shaft through the joint, so that they exactly coincide.
- (3) It allows (by the open method) the operator to see the exact point of the osteotomy and the position of the fragments during and after the correction of the position.

Since the object of choosing this operation over that of the ordinary Gant is that of obtaining a sure and true alignment, the only interest in cases is in this particular. This true alignment is perhaps best shown by a comparison of cases which represent the two methods. It is not claimed that all cases of subcutaneous linear osteotomy show the deformity here represented, but it is certainly a very frequent occurrence, and, in fact, in looking over the x-rays of the cases after operation, it is unusual to find cases which do not show deformity. The illustrations of this deformity were taken from the general service, and many of the observations were collected from cases which came under observation long after operation in various places, and by surgeons skilled in this technique, so that it is fair to claim these results as representative. The lack of displacement and the straight alignment of the weight-bearing line, extending through the femur to the hip-joint, is noticeable in the modified (Curved Gant) operation (Figs. XIX, XX).

(2) The latero-posterior route for arthrodesis of the hip-joint (Albee's operation): There is no doubt that this gives the best approach for the operation of arthrodesis (as described by Albee), for it opens a lateral view of the parts involved in this operation which is almost necessary in order to insure a mechanical operation. By the exposure of the lateral aspect of the joint structures, it allows exploration and removal of the overgrowths, which are usually found about the acetabulum and about the edge of the head, which sometimes very materially interferes with the good apposition of the resected parts. It also allows a clear view of the parts in shaping the resected surfaces, which are made to fit together in the reduction of the deformity.

The application of this operation is much wider than was at first supposed, and a very distinct advance was given by Dr. Albee in his method of producing the ankylosis of the hip-joint. The types of cases for which this latero-posterior incision is applicable for arthrodesis are:

(1) Monarticular hypertrophic arthritis, particularly in working people, who need a straight weight-bearing leg.

(2) Deformity in adults of old hip-joint disease of childhood, in which ankylosis is not complete.

(3) Deformity with incomplete ankylosis following infectious arthritis.

(4) Traumatic conditions and old fractures of the neck, which require an open method of operation.

(5) Open operation for congenital hip dislocation.

(1) Hypertrophic arthritis (Figs. XXI, XXII, XXIII): The application in the cases of hypertrophic arthritis has been fully described by Albee in his earlier article. It is for those patients unable to walk but a few rods without pain, and which amounts to disability. The condition is one which ordinarily progresses through a painful stage to a more or less imperfect ankylosis, in the course of a few years. The operation gives an ankylosis, substitutes a strong weight-bearing leg, and cuts short the period of incapacity during the natural ankylosing process. In the experience of the writer, the choice of these cases should be made with care. The cases of patients too far advanced in years, of too great weight, and those showing similar change in other weight-bearing joints, should be selected with a great deal of caution. But such conditions are not necessarily prohibitory. The ideal cases are the younger, of light or moderate weight, monarticular cases of traumatic origin. Healing in these hypertrophic joints is apparently slower than in normal bone, and retention for a longer time than in ordinary bone healing is advisable. As Albee has urged, abduction in the retentive plaster is imperative, as some of this lateral position is always lost.

(2) Deformity after old hip-joint disease, with imperfect ankylosis: One particular application of this operation is for the correction of deformity of old hip disease, on those cases not suitable for osteotomy. It is obvious that such an operation would not be attempted except at a period remote from the cure of the original process; but in adults, who have recovered from the disease of childhood, the procedure is apparently safe. Many of the cases present the deformities usual after hip disease, but with imperfect ankylosis, allowing a slight degree of motion, not enough to be of use or convenience, but enough to be of menace to the patient. Occasional stress frequently gives adhesion strain, with occasional lighting up from these injuries, so that correction of the deformity is only a part of the indication for operation. The distortion of the remaining portion of the head and neck varies very greatly in these cases, and one type commonly found results from the absorption of the neck, the trochanter remaining normal. In this the upper part of the trochanter lies nearly in contact with the ilium, and with any attempt at abduction of the leg serves to pry on the adhesions of the remnants of the joint (Figs. XXIV, XXV). If firm ankylosis has resulted from the original

disease, this has no influence on the function of the limb, but with motion remaining, however slight, the effect of the abduction pry is very different. This operation has the advantage that it corrects the deformity in the most mechanical way, and at the same time, by producing an ankylosis, relieves a vulnerable joint of its constant source of menace (Fig. XXVI).

(3) Deformity with incomplete ankylosis after acute infectious arthritis of hip (Figs. XXVII, XXVIII). This class is perhaps the most favorable for this operation, and in these cases it is safe to operate much sooner after the original process has subsided than in the cases of tubercular origin. The bones are not fragile, as in recent tubercular process, but, on the other hand, are quite as hard, or even harder, than ordinary bone. It very seldom if ever happens that these acute infectious processes in adults result in a true ankylosis, certainly not until after long periods, and the saving of time to these patients by earlier operation is an important factor.

(4 and 5) Traumatic conditions and dislocation (Figs. XXIX, XXX): In dealing with these, especially with dislocation, it may not be necessary to remove the top of the trochanter, for it may not be desired to reach the articular surfaces. If, long after, retention is necessary, as is usually the case, its removal has no disadvantage, as the union of the fragments will take place in this time, and it is far less destructive than to sacrifice muscles by cutting, or by the strong retraction which is otherwise necessary in reaching the deep structures of the joint.

A brief résumé of the cases operated upon for arthrodesis is given to show the general type for which relief is attempted. With these are a few (Cases XVI, XVII, XVIII) in which the ankylosing operation was not done. In each the incapacity of the patient was such that he sought some radical measure for relief, and the character of the operation was left open to the operator, to be decided after the joint structures were exposed. In these the patient was given the benefit of any doubt, and the joint preserved. In none did there seem to be any reason to regret the removal of the trochanter (when done), and the greater view afforded made the decision in regard to the choice of procedure much the more sure. In the earlier series the anterior route was used, and later the lateral, but neither gave satisfactory approach without too great violence to the covering muscles. With the latero-posterior route, and removal of the trochanter, these objections are overcome, and this route has since been used exclusively for this type of operation. The cases upon which an ankylosing operation has been done, are put in a double plaster spica, on the affected side, including the foot, and on the well side reaching half way to the knee. The body portion on the side opposite to the affected leg is carried well up on the side of the chest, to prevent the patient from swinging the body into the line of the affected leg, and thus lose a portion of the abduction. On the opposite side of the trunk the plaster may be left low. The short

plaster on the well leg is removed after two or three weeks. The long spica is worn eight weeks, and may then be replaced by a short spica, reaching to the knee, and this lesser retention is continued four weeks longer. The consolidation in these cases apparently takes longer than in fractures of normal bone, and abduction may be lost during any period while consolidation is still not complete.

I wish to express my thanks to Dr. John Warren, of the Harvard Medical School, for the courtesy in providing the material and opportunity for dissections made in connection with this study.

CASE I. Age, —. Old hip disease; recovery with ankylosis in flexion and marked adduction. Gant from subtrochanteric osteotomy by subcutaneous method. Bone chiseled two thirds through, and complete fracture obtained with slight force. Plaster spica with leg in abduction.

After recovery from ether, patient complained continually of pain on any attempt at movement in bed. Examination by x-ray showed bone to be broken below point of osteotomy with long splintering anterior and inner portion of shaft; lower fragment had slipped upwards underneath the overhanging fragment of bone. Reposition required open incision with reshaping of end of bone and wiring.

Convalescence delayed and union slow on account of small portions of bone in apposition. Final result good as to union, but considerable shortening resulted.

CASE II. Age, thirty-four. Hip disease during childhood. Has been treated for lameness by occasional plaster spicas for past two years without relief.

Examination: Right hip — permanent flexion, 45°; adduction, 35°; inversion; no knock-knee; marked sensitiveness with attempt at motion and with walking.

March 11, 1909, operation, intertrochanteric osteotomy of right femur; bone broke during osteotomy. Plaster spica with leg in abduction.

April 21, 1909, patient has had pain with attempts at movement in the bed, and x-ray shows long splintering of bone with inner and anterior portion of upper fragment overriding the lower, and which has slipped by.

April 29, 1909, operation, anterior incision inside of sartorius; fragments found to be overlapping, and spicule of anterior and upper end of upper fragment lying over the lower, which had slipped upward. No fresh surfaces of bone were opposing for union. The two ends were brought into position, surfaces freshened, and the bones wired together. Plaster spica, which was worn five months. At this time union was firm and leg in good position, but length had been lost.

CASE III. Age, forty-two. Hypertrophic arthritis of left hip; symptoms dating back three to four years; disabling symptoms for nine months; practically complete disability for three months. Motion in hip limited to 15° of motion in flexion. Deformity in flexion and adduction; x-ray shows large exostoses of head; overlapping of acetabulum and absorption of cartilage.

Operation, Sept. 11, 1908, by lateral incision over trochanter. Head found to be enlarged, roughened by large exostoses, articular surface devoid of cartilage. Arthrodesis by Albee method. Plaster spica with leg in abduction.

Final result: Good weight-bearing leg. The case was not in all respects best suited for the operation. There were slight hypertrophic changes in the right

knee, and the general condition of the patient was not good.

CASE IV. Age, fifty-one. Hypertrophic arthritis of right hip, apparently following rheumatic fever eleven years ago.

Examination: Patient very stout, weighing 240 lb. Marked limitation of motion in flexion; other motions practically nil; sensitiveness and spasm on motion; permanent flexion and adduction. Has had practical disability of right hip for two to three years. Right knee slightly affected by same condition.

Oct. 9, 1908, operation, lateral incision; joint opened between gluteus medius and tensor fascia femoris; marked hypertrophy found about the upper edge of acetabulum, and no cartilage on joint surfaces. Arthrodesis by Albee method. Plaster spica.

On account of size of patient it had been impossible to keep leg in plaster and bony union was not obtained. Second operation, March 25, 1909, operation by anterior incision between sartorius and tensor fascia femoris. Fresh surface removed from acetabulum and head of femur and inner side of head curetted down to rough bone. Plaster spica. Patient had uncomfortable convalescence on account of great weight.

June 14, 1909, plaster removed; able to walk with chair, with support or with crutches. Leg shows practically no motion; difficult to determine whether there is actual ankylosis on account of size of patient. Knee gives some trouble.

Final result: Distinct improvement over original condition, but there is very considerable difficulty in getting about. The knee is in part accountable, but the hip is far from painless. The result is not satisfactory.

CASE V. Age, thirty-nine. February, 1908, traumatic dislocation of left hip; reduced following day, with good recovery. Well for one year, then began to have pain in left knee and hip, and disability on walking and weight bearing.

Examination: Practical disability of left hip; motion in flexion limited to about 20°; marked spasm; other motions in other directions nil; permanent flexion 20°; abduction.

March 21, 1910, operation by the anterior incision. Capsule found markedly thickened and tough; a little fluid in joint; marked lipping of acetabulum, and distinct erosion of cartilage of head of femur; exostoses about the head of femur. Arthrodesis by Albee method. Plaster spica with leg in abduction.

Final result: Four months after operation patient had good weight-bearing leg, but not entirely free from sensitiveness; patient has not been seen since this date.

CASE VI. Age, thirty-one. "Rheumatism" thirteen years ago. Neisser infection seven years ago; frequent tonsillitis. Applied to out-patient department September, 1907, with pain in knee and limp. Examination showed atrophy of right leg; crepitus in both knees; motions of hip limited; slight deformity in position of hip. X-ray shows hypertrophic process of right hip.

Returned February, 1910, with increase of symptoms. Plaster spica applied with some relief.

June, 1910, deformity of abduction and flexion increased. Motion in flexion, 20°; abduction, adduction and rotation nil.

June 2, 1910, operation, anterior incision. Large amount of exostoses found. Arthrodesis by Albee method. Plaster spica with leg in abduction. At time of discharge from hospital leg was firm, but patient has not been seen since and final result cannot be given.

CASE VII. Age, forty-five. Monarticular hypertrophic hip; several years onset; two to three years of practical

disability. Patient extremely heavy. Leg in position of permanent flexion and adduction, and marked limitation of motion in all directions, particularly rotation.

December, 1910, operation, latero-posterior incision; large overhanging lip of acetabulum; cartilage practically disappeared from articular surfaces. Arthrodesis by Albee method. Plaster spica with leg in slight flexion and abduction.

Convalescence of patient delayed on account of sepsis of superficial wound, but recovered with good position and firm union.

Six months afterwards, patient had good weight-bearing limb and walked with only slight limp.

CASE VIII. Age, twenty-eight. Treated eighteen years ago for hip disease. Gives history of frequent exacerbations of pain, limp and sensitiveness during past ten years. Last winter had a fall with severe and persistent sensitiveness.

Examination: Motion in flexion 10° with pain and spasm. Deformity in flexion, 30° ; in adduction, 20° . Shortening two inches. Trochanter above Nélaton's line. X-rays show old acetabular and femoral disease; marked destruction of head; shortening of neck.

Nov. 17, 1910, operation by latero-posterior incision with removal of trochanter. Arthrodesis by Albee method. In one portion of the anterior portion of neck, in the field of bone incision, a small area apparently an enclosed focus. This area entirely removed, curetted and washed with carbolic and alcohol. Plaster spica with leg in abduction. Considerable pain after operation. Patient ran a slight temperature for about three weeks. Temperature then became and stayed normal. Plaster worn fourteen weeks.

Result: Ankylosis in good position; leg strong for weight bearing.

CASE IX. Age, twenty. Two attacks of arthritis of hip with partial recovery. Cause of arthritis not known. Practical disability since last attack six months ago.

Examination: Left hip shows practically no motion and with leg in good position. Right leg shows few degrees of motion with spasm.

November, 1910, operation by latero-posterior incision. Neck found slightly enlarged; articular surface entirely devoid of cartilage; considerable roughening about edge of acetabulum and rim of the head of the femur. Arthrodesis by Albee method. Plaster spica with leg in slight flexion and abduction.

Six months afterwards leg firm, painless. Patient later died of intercurrent causes.

CASE X. Age, twenty-four. Osteomyelitis several years previously, during childhood, followed by sinuses for three years. Later, involvement of hip. At time of entrance $1\frac{1}{4}$ inches shortening; very few degrees of motion; permanent flexion 40° and marked adduction.

Operation, April 11, 1910, latero-posterior incision. Arthrodesis by Albee method. Plaster spica with leg in abduction.

Four months later ankylosis firm and leg strong in weight bearing. Some discomfort from an old relaxed knock-knee.

CASE XI. Age, forty-three, infectious arthritis, polyarticular, with recovery from all except hip.

Examination showed permanent flexion 30° ; adduction 20° ; no motion; $\frac{1}{2}$ -inch shortening.

Operation, March, 1911, latero-posterior incision. Ankylosis found not to be complete. Arthrodesis by Albee method. Plaster spica with leg in abduction.

Four months later ankylosis firm and leg strong in weight bearing.

CASE XII. Age, twenty-six. In childhood was treated for hip disease with abscess. For several

years practical ankylosis of hip. Patient also has ankylosis of knee, on opposite side, from arthritic condition at time of childhood. X-ray shows very marked overgrowths of bone about acetabulum and enlargement of head and neck. Condition considered to be that of infectious arthritis at time of childhood.

March, 1911, operation by latero-posterior incision. Arthrodesis by Albee method. Plaster spica with leg in slight flexion and abduction.

Plaster removed at end of eight weeks. Leg apparently firm. Four months afterwards leg firm with no symptoms and strong in weight bearing.

CASE XIII. Age, forty-eight. Severe tonsillitis in spring of 1911; ill for three months in bed and several abscesses in throat. After recovery leg was found to be flexed and without motion. Seen in October, 1911, at which time leg was flexed in 45° ; 15 to 20° of adduction; no motion. X-ray showed distinct destruction of articular surfaces.

November, 1911, operation, latero-posterior incision; joint surfaces found destroyed; no ankylosis. Arthrodesis by Albee method. Plaster spica with leg in slight flexion and abduction.

CASE XIV. Age, fifty-four. Ten months ago severe septic thumb, resulting in partial amputation. During this time, acute septic arthritis of hip, resulting in deformity of 90° of flexion and 45° of adduction. Patient when first seen in too poor general condition for operation. Leg brought into improved position.

November, 1911, operation, lateroposterior incision. Considerable induration of tissues on outer side about the fascia, joint being found entirely destroyed but not ankylosed. Arthrodesis by Albee method. Plaster spica with leg in slight flexion and abduction. Case still under treatment.

CASE XV. Age, fifteen. History of traumatic dislocation from a fall at age of four. Mechanical reduction of dislocation attempted; the head was felt to slip into place and reduction apparently occurred. After removal of the plaster several weeks later, the leg could be brought into the position of complete extension, with 20° of abduction, but full extension in the normal line was not possible, apparently on account of contraction of the fascia lata. By incision of the fascia, the leg could be brought nearer into complete extension, but some abduction remained.

April, 1910, leg in position of abduction and flexion; motion at the hip practically nil; unable to use the leg in walking.

April 14, 1910, hip-joint exposed by the lateral incision; an old fracture was found across the upper part of neck, and extending to the articular surface; the head was rudimentary and the cartilage much destroyed. Arthrodesis, Albee method. Plaster spica applied with leg in abduction.

Final result: Complete ankylosis and useful leg for weight bearing.

CASE XVI. Age, forty-five. Dislocation of hip, with other injuries, two years ago. Hip reduced under ether on second day. In bed for several weeks on account of other injuries. Hip gave no trouble for several months. Later increasing disability, and finally, at time of entrance, practically complete disability on account of stiffness and pain in walking. X-ray shows line of apparent cleavage on a portion of the articular surface of head, with some lipping of acetabulum.

Operation by latero-posterior incision. A few overhanging fragments removed from acetabulum. Thin triangular piece of head found loose; other portions of head seemed to be covered with apparently normal cartilage. Loose portion of bone removed; capsule closed. No plaster.

Four months later leg held in good position and shows motion in flexion to 90°; able to walk without pain. Patient has not been seen since, and final result cannot be given.

CASE XVII. Age, forty-five. Increasing pain and disability in right hip for several months. Unable now to work. Limited and painful motion of hip; slight permanent flexion; atrophy. X-ray shows hypertrophic changes about acetabulum and edge of head of femur.

Operation, July 17, 1911, latero-posterior incision. Exostosis removed from about edge of acetabulum. Articular surfaces not eroded. Capsule closed. Dressing without plaster.

Six months later motion free in 90° flexion; abduction and rotation one half. Nov. 4, 1911, walks with scarcely perceptible limp.

CASE XVIII. Age, —. For ten years increasing stiffness and sensitiveness of both hips. Unable to walk for a year. X-rays show very decided exostoses about acetabulum and edge of head, and flattening of articular surface of head.

Operation, June, 1911, lateral incision; trochanter not removed; large amount of exostoses removed. Cartilage on articular surface thin, but no erosion to be seen. Capsule closed. Dressing without plaster.

Four months after motion increased to practically twice that before operation. Sensitiveness markedly diminished. Walks short distances with comfort. Has requested operation on other hip.

PREVENTION OF INFANT MORTALITY BY BREAST FEEDING.*

BY WILLIAM H. DAVIS, M.D.,

Vital Statistician of Boston Board of Health.

MUCH has been said from time to time in favor of breast feeding of infants; and statements have been made that 80 or 85% of infant deaths occur among bottle-fed babies.

But such statements have lacked force because it has not been apparent how many mothers nurse their babies and how many bring them up on the bottle.

If 80% of infant deaths occur among bottle-fed babies, and if it should be found that 80% of all babies are bottle fed, then there would be no case at all against bottle feeding.

In order to find out the percentage of babies that are bottle fed, the Boston Board of Health recently sent about 900 letters to mothers who had births recorded within the past year. Care was taken to select proportionate numbers of babies of various age periods, namely, two weeks to one month, one to three months, three to six months, six to nine months, and nine to twelve months, and to select proportionate numbers of babies from the various wards and from the various mother nativities, — the number of births in 1910 in the various wards of various mother nativities being taken as the basis for these proportions. Aside from obtaining the proportionate numbers of babies in the way just indicated, care was taken not to select these babies in any way but to take them at random from the

* Read at annual meeting of Massachusetts Association of Boards of Health, Boston, Jan. 25, 1912.

birth returns furnished by the Registry Department. The circular sent to each mother was as follows:

"Dear Madam, — To assist the Board of Health in its campaign to save the babies of Boston, will you please answer the following questions regarding your child and return this slip at once in the enclosed envelope.

"Is this child breast fed? (Answer yes or no.)

"How long was it breast fed?

"Is this child bottle fed? (Answer yes or no.)

"How long has it been bottle fed?

"If bottle fed, what food has been used?"

Replies were received promptly from over half of the mothers addressed, and an additional 200 replies were obtained through the Board of Health nurses, who went to the homes for the information.

Table I gives the results of this investigation, — the number of replies received for babies of various ages and mother nativities and the percentages of breast and bottle fed babies.

The number of replies was 736, of which 533, or 72.4%, were breast fed and 203, or 27.6%, were bottle fed. But the information thus obtained is only for living children. To determine what proportion of all babies are bottle or breast fed, it is necessary to add a proportionate number of the deaths of 1911. These necessary additions are given in Table II. Adding the figures in each square of Table II to the figures in the corresponding square of Table I, totals may be obtained which show the relative proportions of breast- and bottle-fed babies in Boston. The corrected percentages shown in Table II were thus obtained. At various places in the table there appear higher percentages of breast feeding for the older babies than for the younger. This, of course, is an error due to too few replies and would be rectified if answers from all the mothers of 1911 could be tabulated. But the evidence furnished by these 736 replies is very striking and may be used for present calculations.

Italy leads, with 83% breast fed; Russia and Poland come next, with 79% breast fed; Ireland and the United States follow, with 73% and 59% respectively; while Canada makes the poorest showing, with only 51% breast fed. The replies from mothers born in England, Scotland and Wales, Germany and Scandinavia, were too few to warrant any estimate for the babies of these mother nativities.

Table III gives for 1911 the total mortality of infants by age, by mother nativity and by feeding (for infants over two weeks of age).

Table IV gives for 1911 the total mortality of infants by month, by age and by feeding (for infants over two weeks of age).

Notice in Table IV that there is only a slight increase of deaths among breast-fed babies during the summer months. July, August and September, with 39 and 40 such deaths each month, are lower than February, March and April with 49, 41 and 38 deaths respectively. The number