acute inflammation through the whole articulation, and that inflammation becomes diffused through all the textures round about. It did so in this man. When the state of affairs became manifest, he himself begged for the amputation of his foot, and I was not long in consenting, for it seemed certain that though the foot might have been preserved for him, and though it might have served for a man of ease and leisure, it would never do for a man who had to work for his living. One must have left a deep scar at the bottom of his foot on which he could not have borne pressure in standing or walking; and one could not change the position of his foot, for he could not bear the pressure of an instrument while the metatarsus had suppuration over it. Therefore, on these grounds, we have decided on the amputation of his foot. I shall amputate it according to the method called after Mr. Syme—that is to say, removing it at the ankle-joint, and including the os calcis. You may ask why I am going to perform an operation at so great a distance from the seat of disease, which does not extend very far above the metatarso-phalangeal joint of his toe. It is a principle in all operations upon the foot, that you should take care that the scar is not afterwards exposed to the pressure and friction to which it is liable when the foot is used in walking. Now, this man has a contraction of his tendo Achillis—a contraction, it seems, not curable by any means that one can apply to the front of the foot. He would be left, after an amputation anywhere in the front of his foot, with a scar—or, at least, with the stump of the foot—directed, not straight forwards, but obliquely downwards. The stump, if for instance Chopart's operation was done, would be directed obliquely towards the ground, as the foot is now. The man would be put into a condition no better than he has been in hitherto. If he were able to walk on the stump at all, corns and bursæ would form, and, after a little time longer, ulcers, and in all probability ulceration of the scar, quite unmanageable and unfit for healing. So that I cannot doubt—and in this opinion I have the concurrence of all my colleagues—that the best plan will be to remove the whole of the man's foot at the ankle-joint, including the os calcis, and then to get for him a round stump, on which, as he walks, he will come plump down, and that upon a piece of integument naturally fitted for bearing the support of his body and the friction of the ground.

The case, then, may serve to illustrate to you these two or three points of great importance. First, in regard to the position the foot must be put in, in the treatment of all cases of diseased ankle-joint, diseased tarsus, diseased bones of the foot—namely, with the foot at right angles to the axis of the leg. Then the series of events that occurred: corns and bunions not carefully treated; the liability of the bunions to suppurate, and to open into the cavity of an adjacent joint; then the symptoms indicating the necessity of amputation; and then, lastly, the rule that in amputations you must look most carefully that the surface of the foot which you leave is one that will fairly and thoroughly bear pressure and friction.

ON THE TREATMENT OF ASTHMA BY BELLADONNA.

By HYDE SALTER, M.D., F.R.S.,

FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS, SENIOR PHYSICIAN
TO CHARING-CROSS HOSPITAL.

I do not know any sedative that has more "improved its position" of late years than belladonna. It is certainly given much more extensively, and is much better thought of, than it was twenty years ago. And I think the one is the result of the other: it is better thought of *because* it is more extensively given. There are cycles of therapeutics as well as cycles of disease; and many valuable remedies pass into a temporary desuetude by a common neglect, and then, from some accident, emerge into repute and usefulness. And this is not only true with regard to the medical world in general; it is equally so with regard to the practice of individuals. We are all of us apt to get into a groove in pre-

scribing; and not only get into the habit of employing certain remedies, but get *out* of the habit of employing others. I must confess to this fault myself, and feel at a loss to explain on other grounds how it was I went on so long without giving belladonna a fair and thorough trial in asthma. For the last two years, however, I have used it extensively, and am so satisfied with the results that I think it worth while to publish a few of my cases, in illustration of its value and of my method of employing it.

If I were to express what appears to me to be the peculiar excellence of belladonna as a sedative in asthma, I should say it consisted in its power of diminishing reflex irritability—a power which it appears to me to possess in a degree greater, in proportion to its other sedative effects, than any other sedative.

My method of giving it will be best seen in the following cases. I do not know that the tincture has any advantage over other preparations, only it happens to be that which I have tried. I should think that the liquor atropiæ might, from its uniformity of strength, be even preferable.

CASE 1.—R. G—, a lady aged twenty-five, who has been liable to hay fever for eleven years and to asthma for seven, came to me on May 18th, 1868, for the cure of her hay asthma. She had been suffering for nearly a month, and had not had a single night's rest. She was not troubled with her asthma by day, but only with her hay-fever symptoms. As soon, however, as she went to bed on came her asthma, and lasted regularly all night. She was worn out for want of rest, and felt fagged and miserable. I ordered her ten minims of the tincture of belladonna three times a day, telling her to increase the quantity, day by day, till the characteristic effects of the drug were produced. She gradually raised the dose from the ten minims to twenty. She felt better the first night after taking it. She took it for nine days, and had no asthma whatever. She then, thinking herself cured, and feeling her head very uncomfortable, omitted it for one day, and the following night got no sleep from asthma. She then resumed it, and has had no asthma since. She now regularly takes twenty drops three times a day. Her head is slightly affected, and her eyes decidedly so—misty, and she cannot read; but there is not a trace of asthma. She has never found anything else prevent the attacks. The belladonna seems to have no influence whatever over the other hay-fever symptoms; the sneezing, running at the eyes, &c., go on as before.

CASE 2.—Sarah P—, aged forty-six, has had asthma ever since she was nineteen years of age—that is, for twenty-seven years. For the last five months she has been worse than ever, and for some time past has had asthma regularly every night. It begins between two and three o'clock, and after that she gets no rest. I ordered her to take the tincture of belladonna every night on going to bed—to begin at ten minims, and gradually to increase the dose till her head and sight became affected. She came to me on the 29th of last July. On the 14th of August I find the following entry in my note-book:—"This patient came to me a fortnight ago yesterday. She has almost completely lost her symptoms ever since, having had uninterrupted good nights. She has gradually reached thirty drops of the tincture of belladonna, and now takes it every night. She looks wonderfully better, and is able to take long walks." From that time I have not heard anything of her.

CASE 3.—Col. R—, aged forty, has had asthma for sixteen years. For the last few years it has been gradually getting worse, and now he has it almost every night, commencing at from one to two o'clock A.M. He has tried various things without any success—Indian hemp, chlorodyne, ether, the inhalation of chloroform, dry cupping, coffee, iodide of potassium, abstinence, &c. I saw him on Sept. 16th, and ordered him to try the belladonna at night in the usual way. On the 23rd I find the following note:—"He has gradually got the belladonna up to forty minims without any appreciable inconvenience. The nights have been very good. The two last mornings there has been no oppression whatever, nor any sitting up in bed; he has slept right through the night. He walked this morning, directly after breakfast, from his house at Blackheath to the station—a mile—in sixteen minutes, and the same pace up from Charing-cross station here." I have not seen him since.

CASE 4.—Wm. P—, aged thirty-eight, has been asthmatic seventeen years. His attacks come on every morning on rising from bed, but often trouble him throughout the day as well. There is defective respiration at the right base, violent paroxysmal cough, and a good deal of expectoration. This patient came to me on Oct. 13th. A week after I made the following note:—"Since coming to me this patient has taken the belladonna every night—twenty, twenty-five, and then thirty drops, to which last dose he has stuck. The result is that he is quite a different man. He has slept all through the night, dressed himself without any trouble or difficulty, and has walked half over London as quickly and well as ever in his life. When he arrived in town his breath was so bad that he had the greatest difficulty in getting from the Vauxhall station to the church near the bridge. He spits hardly anything. There is now a complete absence of all musical wheezing, and of any adventitious sound whatever: respiratory murmur quite re-established at right base, and indeed perfectly natural everywhere. And this change has taken place in one week."

For the following case, in which liquor atropiæ was used, I am indebted to my friend Mr. Hodson, of Bishop's Stortford.

CASE 5.—"John L—, aged fifty-five, by occupation a gardener, of active and temperate habits, suffered, from the summer of 1864, from frequent attacks of shortness of breath, followed by expectoration. When he first came under my observation (Sept. 2nd) he presented the appearance of a confirmed asthmatic. The respiratory sounds, whenever I saw him, were always more or less characteristic of asthma. At night he generally became worse, so much so that he often found the greatest difficulty in getting up one flight of stairs to his bedroom. I gave him at first iodide of potassium (five-grain doses), then combined it with decoction of senega, afterwards with squills and compound tincture of camphor. He also took lobelia, &c., and burned nitre-paper. With the exception of the last, which always afforded some relief, the frequency and severity of the attacks were uninfluenced by the treatment; indeed he became so bad that his life was a burden to him. I was watching his distress one day, and deploring the inefficiency of treatment, and thinking how the constricted state of his bronchial tubes might be overcome, when the idea of trying a drug that so quickly dilated the pupil occurred to me, and I at once determined to bring him fully under the influence of atropia. He began, Oct. 26th, to take one-sixtieth of a grain in solution every six hours. A few doses produced marked relief, but I pushed the atropia until he took at each dose one-fortieth of a grain. He then became fully under its influence, with greatly impaired vision, entire absence of saliva, and most troublesome strangury. From this time the breathing improved, but he found it necessary to take the medicine two or three times a day up to the middle of the following February. In the spring I sent him for a few weeks for change of air, and upon his return he resumed his full work as a gentleman's gardener, which he has continued without interruption up to the present time (Dec. 1868). When he went away I gave him a supply of his atropia, in case an attack should come on, but he told me yesterday that he has never opened the bottle. During the last three years I have found atropia, in all cases of genuine asthma, quite as useful as in L—'s case, but I have never pushed it with anyone else as I did with him."

I believe one reason why belladonna has not had a greater reputation as a remedy for asthma is that it has not been given in large enough doses. I think that, like lobelia, it must be given in doses sufficiently large to produce its physiological effects, otherwise we have no right to say that it has been fairly tried, or to conclude that it has been a failure if it has not achieved a cure. I think to give ten minims three times a day in some mixture is simply worthless. I have seen now numberless cases in which both belladonna and lobelia have been consigned to the limbo of failures, when a fresh trial of them, on the plan of gradually increasing the doses till an ultimatum has been reached, has proved them to be perfectly successful remedies. Sometimes, but very rarely, belladonna will relieve asthma when given short of a physiological dose; sometimes, but still rarely, it fails to give any relief even when pushed to its full physiological effects; the common thing is for it to fail till so pushed, and then to succeed. When I find belladonna has only been taken in the ordinary small doses, without

any increment, I always regard it as not having been tried at all.

The advantages of administering it in the way I have described are:—

1. That, giving it at night, you bring the full force of the drug to bear upon the disease at the time at which it is most liable to come on, and thus, if you are successful, tide your patient over the critical time.

2. By gradually feeling your way up to the required dose, you are able ultimately to reach without fear a dose which you would be unwilling to prescribe without such a tentative approach.

3. In those cases in which the therapeutical dose is reached before the physiological—that is, in which the asthma yields before the sight or head is appreciably affected,—it enables you to stop short as soon as relief is obtained, and thus spare your patient any of the disagreeable effects of the drug.

4. By giving it only once in the twenty-four hours, you are able to give a larger dose than you would be able to do if oftener repeated.

5. By confining the dose to bedtime, the patient's days are, in spite of a large dose, passed in comfort; for, as the morning advances, the dulness of head, confusion of sight, and drought of mouth pass away.

6. You are thus enabled to find out what is the dose for the individual—a very important point. People differ very much in their tolerance of belladonna. Some of my patients have been unable to take more than twenty minims once in the twenty-four hours without very unpleasant symptoms; while I have known others able to take a drachm three times in the same interval without any inconvenience. And as they differ in their tolerance of the drug, so do they differ in the dose at which their asthma will yield. The only way to ascertain what that dose is, is to make each case a separate experiment, and this can only be done in the way I describe.

7. By giving the remedy three or four hours before the attack is likely to come on, the treatment becomes *prophylactic*. If by taking a dose every night for thirty nights the attacks have been for that time prevented, the patient has ceased to be an asthmatic for a month. This is a very different thing from having had thirty attacks in the same time which have been cut short by the remedy. In all "habitual" diseases, in which the recurrence keeps up the tendency, prophylactic treatment has, in relation to final cure, a pre-eminence it does not possess in diseases in which habit has no place. For such diseases it is *the* treatment. It does more than spare your patient an attack of his malady; it breaks, *pro tanto*, that chain of sequences which is the very life of the morbid tendency.

TABLES ILLUSTRATING THE EFFECTS OF REMEDIES UPON UNCOMPLICATED ACUTE RHEUMATISM.

By W. H. DICKINSON, M.D. CANTAB., F.R.C.P.,

SENIOR ASSISTANT PHYSICIAN TO ST. GEORGE'S HOSPITAL, AND TO THE HOSPITAL FOR SICK CHILDREN.

(Continued from p. 118.)

THE following tables are in continuation of those published last week, giving the progress of acute rheumatism as modified by different methods of treatment. It may be well to repeat that those cases only are included which presented on admission no evidence of heart complication.

The first of the tables already published represented the treatment of eight cases by venesection.

The second table exhibited six cases under mercurial treatment.

The third table consisted of thirteen cases treated by specific and other remedies not alkaline.

The fourth table presented seven cases under the simple saline treatment.

The heart was affected in four of the first group of cases; in two instances in the second group; in five cases in the third group; and in two cases in the fourth group.