dilatation of the cervix uteri; but adhere to the opinion expressed in my work seventeen years ago, that careful dilatation by prepared sponges is the safest and easiest plan, quite as effectual as the division, and has the advantage of being so simple that it can be carried out by any well-informed practitioner. As either this plan of treatment or simple linear division will do all that is required to re-establish the uterine passages when closed, I cannot accept for that purpose the more severe and serious operation now proposed, save as a very exceptional measure.

Grosvenor-street, June, 1865.

REPORT OF A CASE OF ANEURISM OF THE ABDOMINAL AORTA; DISEASE OF THE HEART (AORTIC REGURGITATION).

BY WM. HENRY DAY, M.D.

I. B., aged thirty-two, by occupation a coachman, consulted me on Sept. 13th, 1864, for "pain in the abdomen and back." He drove to my house, and appeared to be ailing very little. I ordered him some stomachic medicine and a mild aperient pill at bedtime. Next day I was requested to see him, as he had passed a most painful night, and was unable to rise from his bed. He referred his pain to the centre of the body as yesterday, but was unaware of any pulsation in the epigastrium till I reminded him of it. Four years ago he fancied he got a "rick." Ten years ago he had cough and shortness of breath, supposed to be phthisical; never had syphilis or rheumatism; was married three months ago.

On examination a large pulsating tumour was detected in the epigastric region, to the left of the median line; the pulsation was distinct and strong, and a short rough systolic murmur was heard over it. When the pain came on he turned faint, and the skin was bedewed with perspiration. He had never passed blood per anum or by vomiting. The area of the heart's dulness was increased on percussion, and there was a loud aortic murmur with the first sound; there was a louder murmur with the second sound, growing fainter towards the apex of the heart. This sound was variable in quality, sometimes blowing, whizzing, or sawing. The pulse regular, and somewhat strong, but compressible, averaging from 64 to 70 per minute; respirations 16. I ordered ten leeches to be applied to the tumour, and a hot fomentation afterwards. A full dose of chlorodyne was given to relieve pain, and to be repeated according to circumstances.

Sept. 19th.—He is better, and the pulsation less. The bowels are rather costive. To take one teaspoonful of confection of senna every night.

He improved for a time under rest and a scanty diet, but as I gave an unfavourable opinion of his case, he went to St. Bartholomew's Hospital, where he was admitted on Sept. 29th. Dr. Kirkes confirmed my diagnosis. The patient died on December 23rd at his own home. Mr. Maberley, of the above hospital, has most kindly furnished me with the following report:—

On admission he was fairly nourished; posture, countenance, lips, teeth, and gums were natural. Respirations 16; pulse 60, of moderate volume, and rather jerking; skin moderately soft and moist, but cold extremities; no eruption; no thirst; no vomiting; bowels relaxed from medicine; urine yellow, specific gravity 1020, slight effervescence with NO_5 ; abdomen soft; sleep disturbed by pain across iliac region; the mental faculties are clear; pulsation at epigastrium; breathing very feeble.

Auscultation.—Cardiac impulse normal. Double murmur at base; diastolic loudest and most prolonged; both murmurs audible at apex, but less distinctly; short rough systolic murmur over pulsation in epigastrium; no diastolic murmur audible there; no aneurismal murmur heard at back.

Oct. 2nd.—Pulsation at the epigastrium much less marked, though still audible through the stethoscope; cardiac sounds as before; pulse 76, and still jerking. The next day he was discharged at his own request.

15th.—The patient was readmitted yesterday. He was weaker than when he left; pulse jerking, with back stroke, and occasionally intermittent. Pulsation very distinct in the abdomen, and traceable all along the curve of the femoral artery and posterior tibial. Appetite good; slept well after tincture of opium.

18th.—Pulse 72, with back stroke.

21st.—Pulse 92, jerking. Pulsation at the epigastrium very strong, and murmurs loud.

24th.—Pulse 102, incompressible and regurgitant; pupils dilated; diastolic murmur most audible at the apex; both murmurs, however, heard at the base and apex; aneurismal systolic murmur also audible at the epigastrium. Discharged at his own request.

27th.—Visited him at his own house, and found him about the same. (Dr. Kirkes expressed his pleasure at my continuing to see him.)

Nov. 3rd.—Half-past four P.M.: Was summoned to him on account of unfavourable symptoms, but found he had recovered within the last two hours. He had suffered from coldness of the legs, and seemed semi-unconscious of things around him; but when I saw him his pulse was strong as usual (84). The pain in the right iliac region had left him; but he was troubled with "shooting pains" in the cardiac region.

6th.—Much as usual; cannot get sleep without an opiate draught; pulsation at the epigastrium loud and strong; heart's impulse strong.

13th.—Sitting up near the fire, and feeling much pain in the epigastrium and right lumbar region; countenance troubled; face slightly flushed; pulse 84, full and jerking. Cardiac impulse and epigastric pulsation much increased. A mustard poultice to be applied to the epigastrium and loins.

20th.—Sitting up in bed, and looking thinner and somewhat oppressed; pupils dilated. The pulsation in the epigastrium now seems to extend along the whole course of the abdominal aorta, and is more perceptible below the umbilicus than at the epigastrium. Heart's action considerable. Diastolic murmur loudest and most prolonged; at the base it becomes almost musical. Systolic murmur much less distinct. He complains of cold sweats, and has almost constant pain in some part of the abdomen. Has lost flesh, and is much weaker. He is irritable in temper, and very fidgety. Pulsation of the axillary artery under the clavicle, near the acromion on both sides, distinctly felt with the finger, and heard through the stethoscope very loudly. Double endocardiac murmur as before. Mitral diastolic "sawing" murmur loudest at the apex; at the base the diastolic loudest, and more musical in character. He complains of numbness in his legs, and in his arm if he lifts it; also pain in the groin, right side, and epigastrium.

27th.—Sitting up in bed, and cannot lie down from strong pulsation in the aneurism. Has had no sleep for many nights (increase the dose of opium to thirty minims, and the same quantity in a poultice). No pulsation is now perceptible in the axillary artery.

Dec. 18th.—A peculiar thrill observed about the centre of the aneurism. The aorta seems now to lie more on the left than the right side. The pulsation in the epigastrium, though still evident, is not so much so as below; there would seem to be two distinct pulsations. Pulse 80, of moderate volume, hard and hammering, though quieter than before. Heart's sounds less distinct; systolic murmur (sawing) loudest and prolonged; second sound scarcely perceptible; murmurs loudest (over aortic valves) at base. Pulsation in femoral artery distinct. Has been troubled much with diarrhoea during the week (two or three days); afterwards costive, and relieved by injections.

23rd.—Suffering much with occasional cold sweats. Fainting came on in the evening, and after ten minutes' partial insensibility he uttered the words "Oh! oh!" and expired.

24th.—On Saturday, next day, Mr. Haynes, F.R.C.S., of Crouch End, and Mr. Maberley, made a post-mortem examination. Mr. Maberley thus writes: "We found a pound or more of coagulated blood under the omentum and between the folds of the mesentery, which blood was traced to the aneurism. The lower part of the aorta was quite healthy. The aneurism seemed to hang free like a bag about the size of a medium orange at the epigastrium. The finger passed readily into the aorta upwards and downwards; the sac had burst, the lower part having become so thin that the finger went readily through it. The lungs were non-crepitant, whitish-grey, and sodden. The heart was large and fatty."

Newmarket, 1865.