his family that it occurred prior to 1829; they know this by circumstances impressing it indelibly upon the memory. He sent a model explaining the application to Paris by Dr. Hall, brother of the late David A. Hall, and received a letter from the French Academy of Medicine, congratulating him and thanking him for his valuable contribution. This letter has been mislaid only within the last few years. I may add that Dr. Jos. Barrows remembers its application at the almshouse in this city before 1828, when he was a student there."

Dr. James C. Hall, under date of Feb. 15, 1878, writes: "My evidence as to the application of adhesive strips as a means of extension in the treatment of fractures by Dr. McWilliams must be circumstantial. I came to Washington in October, 1828. After graduating (1827) I was for one year a resident physician and surgeon in the Philadelphia Almshouse, now Blockley Hospital. The visiting surgeons and clinical lecturers were Profs. Gibson, Horner, Rhea, Barton and Harlan. Very many and all kinds of fractures were treated in the House, and I am positive that I never heard McWilliams' mode either mentioned or practiced during my term of service. . . . My recollection is almost positive that my first knowledge of Dr. McWilliams' use of the strips as a means of extension was derived from an article which the Doctor published in a newspaper here, I thought it was the *National Intelligencer*, but Dr. Toner has searched for it and not found it there. This has not weakened my conviction that Dr. McWilliams published in some paper an account of his method of treating fractures of the femur. . . ."

Dr. Joseph Barrows, under date of June 6, 1878, writes: "Excuse my apparent neglect in not answering until now your letter in reference to the use of adhesive plaster in fractures of the lower extremities. In the year 1827, I was one of the resident students in the Washington Asylum or Almshouse, of which Dr. Alexander McWilliams was the attending physician. Dr. McWilliams at that time exhibited and used adhesive plaster for the purpose above indicated."

In the *Archives Générales de Médecine*, Book ii, p. 247, 1836, I find: Société Royale de Médecine, Paris, séance du 21 Janv: "Rapport de M. Larray sur un appareil à extension continue pour les fractures des extrémités inférieures; par le Docteur Williams de Washington.—Cet appareil est une copie à peu près entière de l'appareil de Boyer. La seule différence consiste dans une attelle interne qui se fixe à la traverse sur laquelle la vis s'appuie. M. Larray blâme la méthode en général et tous les appareils qui s'y rattachent. Il n'adopte pas d'avantage le double plan incliné, et préfère toujours son appareil inamovible."

Dr. B. Troutman, of Philadelphia, Pa., at my request called upon Prof. Gross, asking him for some definite data of Dr. Swift's application of this method of extension and he writes under date of June 27, 1879: "I called on Dr. Gross several times, but he always happened to be out of town, but last week I called again and saw him. He told me that Dr. Swift employed adhesive plaster in complicated fractures first in 1829."

The record of the Almshouse shows in the morning report of July 18, 1827: "Admitted P. Boady, 40 years. Report of July 21, 1827: Fracture femur."

This is evidently one of the cases to which Dr. Joseph Barrows refers in his letter.

In no work on surgery of this date—1827 to 1836—have I found any reference to this method of extension, so it is very evident that there had been no published development of the idea until Dr. McWilliams sent his model, a doll baby with the apparatus applied, to Paris. Dr. J. F. May was present at the presentation and verbally testified to the favorable reception accorded it.

To my satisfaction, I have established the following facts: 1, Dr. Gooch was the first to employ adhesive plaster as a means of extension in fractures; 2, the idea was allowed to pass into oblivion; 3, Dr. Alexander McWilliams, of Washington City, was entirely ignorant of the work of Dr. Gooch; 4, Dr. McWilliams developed and put in practice the idea in 1827; 5, the claim of Dr. Swift is too vague to carry weight with it; and 6, the published report of the Société Royale de Médecine is the first published account of the method after its revival by Dr. McWilliams.

TREATMENT OF OPEN OR COMPOUND FRACTURES.

Read in the Section on Surgery and Anatomy, at the Forty-fifth Annual Meeting of the American Medical Association, held at San Francisco, June 5-8, 1894.

BY HENRY H. MUDD, M.D.

ST. LOUIS, MO.

An open wound communicating with a fractured bone is a serious complication. Every effort is directed to its closure by sound healing. Its importance and the difficulty of its management is determined largely by the amount of injury inflicted upon the soft tissues about the fracture, and by the extent of the comminution of the bone. This open wound is usually determined by the violence which inflicts the fracture, but may result from ulceration. An oblique fracture where the skin has been torn by the sharp angling of the bone, with or without impact against an external object, offers a good opportunity to convert a compound into a simple fracture.

Fractures by direct violence which is not crushing, even if the bone is comminuted, present often good opportunity for the immediate closure of the wound. The soft parts are not necessarily much injured by such a fracture. This is especially true where the bone is superficial.

Extensive lacerations and contusions of the soft parts with a small or a large wound of the skin preclude an effort at immediate closure of the skin wound. They demand drainage. Even in such conditions a compound fracture may occasionally be promptly converted into a simple one by the agglutination of structures contiguous to the fracture, though the skin opening may remain free for drainage.

Absolute cleanliness is the first requisite in an attempt to convert a compound into a simple fracture. A razor, soap and water, are essentials, and must be well used in order to cleanse the skin. The purity of the water used in washing the wound is of first importance. The removal of extraneous matter from the wound may be accomplished by a wash of hot water, but where grease, cloth, and dirt of various kinds are adherent to the raw surfaces, the fingers, forceps, etherial soap, turpentine, and sublimated washes may fail to accomplish perfect cleanliness,

though the wound may be rendered aseptic and heal readily. A fairly perfect adjustment of the fragments is a requisite of early healing. Maintenance of the adjustment by a well adapted splint which does not provoke engorgement and pain by undue pressure, is a most important item in the treatment. Many compound fractures where the local conditions would permit primary union of the wound, fail to progress favorably on account of the injudicious application of an apparatus, which is good in itself but is imperfectly applied.

The external wound may be hermetically sealed by suture and collodion, if it is small and the soft parts are not extensively bruised. Where the skin wound is extensive, that portion of it which overlies the fractured bone should be closed by suture, if a point favorably situated for drainage can be found somewhat removed from immediate contact with the bone.

If bone is protruding and has been contaminated by clothing, by a dirty skin, or by external objects, it must be thoroughly cleansed by careful picking and sponging, and then washed with an antiseptic solution, in order that aseptic conditions may be obtained. If the protruding fragment is not essential to solid union, the difficulty of cleansing its surface and the doubtful vitality of sharp long points of bone, which are denuded of periosteum and bruised by violence, justify its removal by bone nippers or saw. In the more extensive injuries, where direct violence has crushed the soft tissues, comminuted bone, and driven foreign matter into the wound, the cleansing of the parts and the removal of dead tissue will very often require an enlargement of the wound, or an incision to give free access to the injured parts. Fascia and muscle if irretrievably damaged must be cut away with the scissors. Loose fragments of bone are to be removed, and a complete knowledge of the extent of the injury to the bone and to contiguous parts, including joints must be obtained. Blood clots should be washed out, and after thorough cleanliness has been obtained, efficient drainage must be provided. The original skin wound may be closed and drainage be secured by special incision, or through an opening made for the inspection and cleansing of the wound. Every effort is to be made to avoid suppuration, as it will destroy still more tissue. It should be remembered that it is difficult to maintain a perfect aseptis about a wound which is slow in healing, hence even where the skin opening can not be closed, it is important to so arrange bone and soft tissues as to favor the agglutination of soft tissues about the line of fracture, in order that it may be shut off from the external wound at the earliest possible moment.

Fragments of bone, or the sharp ends of the fractured surface must not be permitted by malposition, to exercise undue pressure upon the soft parts. Angling at site of fracture must be corrected, and if the tendency to shortening is marked and not easily controlled, as it is not infrequently, where the ankle joint is involved in the fracture, the shortening of the limb by excision of the fractured end is proper and is a conservative measure. The adjustment of the compound and comminuted bone may not be perfect, but it must be practically so good that undue pressure on the soft parts will not produce ulceration, and must be such as to give a good outline and a useful extremity.

Extension and lateral support are the two chief

agencies in maintaining apposition of fragments. Wiring of bone and the use of ivory pegs are occasionally necessary and can be used to advantage in isolated cases. They are, however, rarely necessary, and are an obstacle to rapid and solid union, except where they overcome the greater danger of persistent and obstinate displacement.

A solid union will often result without serious constitutional disturbance, even where the fracture remains an open one. The exposed bone will develop granulations, and if a part of it is necrosed, separation by granulation along the line of junction of the dead and living bone will occur, even while new bone is forming at other areas in the line of fracture.

Cleanliness, an easy and comfortable position for the fractured member, free drainage, and a good constitutional condition will generally insure solid union, though it may be long delayed. Union of bone in a compound fracture is much slower than in simple fractures, or than it is in the compound fractures which are promptly converted into simple ones.

Delayed union should be tolerantly dealt with; a persistent maintenance of the conditions which favor the nutrition of the injured leg will very generally be rewarded by a good recovery.

A thorough appreciation of the extent of the injury to the different structures of the extremity, must be obtained in cases where the judgment wavers between an effort to save the limb and an amputation. Amputation should be immediate,—that is it should be promptly enforced, as soon as the shock has passed, or postponed until the acute inflammation has subsided. A so-called secondary operation is safer than an immediate one.

The perfected technique of aseptic surgery enables us to save many extremities, where suppuration with its attendant damages must have supervened and destroyed the limb, before the details of antiseptic and aseptic methods were well understood. No less important is the selection of a splint which permits cleanliness, and at the same time insures the rest and the undisturbed position of the fracture, which is so essential to the rapid repair of the soft parts, and the union of the broken bone. The suspension splint of Dr. John T. Hodgen with its extension and free motion is a most important adjuvant in the treatment of compound fractures of the leg and thigh.

AFTERNOON SESSION, JUNE 6.

Dr. Graham, Chairman of the Nominating Committee, presented a verbal report, recommending for Chairman Dr. Joseph Ransohoff, of Ohio, and for Secretary, Dr. R. H. Sayre, of New York. The gentlemen nominated were elected by a unanimous vote.

Dr. LEVI C. LANE, San Francisco, read a paper on "Concussion of the Brain."

A paper by Dr. OSCAR H. ALLIS, Philadelphia, on "Treatment of Fractures of the Lower End of the Humerus," was read by the Chairman, Dr. Roberts, its author being absent.

Dr. P. S. CONNER's paper on "Treatment of Fractures of the Lower End of the Radius" was read by title.

Dr. BEDFORD BROWN's paper on "Treatment of Fractures of the Neck of the Femur" was read by title; also the paper of Dr. LLEWELLYN ELIOT, Washington, D. C., on "Treatment of Fractures of the Shaft of the Femur;" also the papers of Drs. H. H. MUDD, St. Louis, Mo., and JOHN B. HAMILTON, of Chicago.

DISCUSSION.

The discussion was opened by Dr. R. H. SAYRE, who said— I would just like to put myself on record as differing in some particulars with that paper which has just been read, and which, unless some remarks are made about it, would

probably go down as the unanimous consensus of opinion of the Section. It seems to me that there were a good many cases in which fractures of the lower end of the humerus are better treated in a flexed than in an extended position, and I think that the reason possibly that we get the various bad results which Dr. Allis has mentioned, of loss of function on account of the destruction of the normal angle between the humerus and the ulna, is on account of their using, as he says, arbitrarily manufactured splints. I think you must adapt your splint to the patient, and if you take an arbitrary rectangular splint and tie a broken arm to it, you will probably get a worse result than if you let the arm hang in a straight position and put a board on the front of it. But it seems to me that neither of those methods is as rational or scientific as placing the arm in a position of right angles, and then adapting the splint to the particular shaped arm that you have to deal with; and not only that you should come down to the wrist, but that you should come down and include the wrist in the splint. I turn the hand with the palm upward, and as the splint is being applied for this purpose I prefer plaster-of-paris rubbed into the meshes when it is put on, rather than to put on a bandage and pepper it over the surface with a plastic substance, as Dr. Allis suggests. And while you are putting on this you put the arm in a position so that the forearm is in a normal position, and then when you get a final recovery you will not have lost the carrying power, you will not have destroyed the relation of the external and internal condyles to one another, and you will have an arm which if you should have the mischance to get ankylosis, which is very improbable unless it is a very bad case, is in a more useful condition than if the arm is straight. I think that except in cases of exceptional severity, fractures of this kind should be so treated that the man is able to go about his business, and if he is a professional man he can attend to a good deal of work and not suffer this loss.

DR. L. C. LANE, San Francisco—As Dr. Sayre has said he wishes to put himself on record, not entirely upon the side of Dr. Allis, that is in favor of the flexed position, I wish, on the contrary to be on record as in favor, wholly, unconditionally, and forever in favor, of treating in the straight position. I have treated over one hundred cases of fracture of the elbow of different kinds, and have never had a bad result when I treated in the straight position; we have a great deal more muscularity in the front than behind. In the fracture at the elbow, the joint is usually torn, or some of its muscles are torn, and if you put the arm in this position (illustrating) these torn muscles here about the blood vessels, here about the nerves, heal, and with the cicatricial contraction of the vicinity, fix and hold those blood vessels so that there is danger. When the arm is kept in the straight position the several irregular surfaces or hollows, that is, the concavities or convexities one fills the other; but when you bend it in this way (illustrating) you leave hollow spaces, into which cicatricial or neoplastic growth will come in and interfere greatly with your passive motion. The only point on which I disagree with Dr. Allis is in regard to passive motion. He would defer it for some three or four weeks. I would have it a great deal earlier. I would say in regard to Dr. Allis, he is really the author of this method of treating in the straight position. That credit is given to him in some of the foreign authorities. You will find it in Billroth's "Surgery." He has the credit of recommending that position, and he has fought for it. When he began he was almost the only man in the United States in favor of it. As I say, I agree wholly with the plan of Allis, except I would commence passive motion earlier than he does.

DR. BERGSTEIN, Nevada—It is almost unnecessary for me to add anything now that Prof. Lane has robbed me of my thunder. And it is not surprising, considering that he taught me. But I can add this, that while dressing the arm in a flexed position may be more scientific, the proof of the pudding is in the eating thereof, and for twenty odd years I have treated a large number of cases of fracture, extending into the elbow joint, in the straight position without a single failure or a loss of a particle of the usefulness of the member. I have generally made my splints from that magnificent material known as a shingle, and have padded it with cotton. I have begun passive motion in three weeks, and have never seen bad results follow it.

DR. KURTZ, California—I only add a few words to what has been said already, and that is this: Whether we get ankylosis, or partial ankylosis from the extended position, or from the flexed position, it is immaterial; if you get that far, you will have great difficulty to remedy the evil by

simple passive motion. It has been my practice for many years to add to passive motion, massage. I think that no fracture should be treated without massage. But fracture in the neighborhood of the joint, and fractures of the elbow most particularly, leading so often not only to ankylosis but to suits for malpractice, I would suggest to you never to omit to practice massage, say from two weeks after the occurrence. It can be just as easily practiced as the passive motions themselves.

DR. C. A. WHEATON, St. Paul, Minn.—I would like to occupy the attention of the Section for only a moment to express my gratification at the position taken by Dr. Sayre. The title of the paper, it seems to me, is decidedly ambiguous. It speaks of fractures at the lower end of the humerus. The gun-stock deformities of fracture at the elbow joint, as a rule, result from fracture of the internal condyle. If we have a fracture of the lower part of the humerus, involving the elbow joint and we attempt to put that fracture in position it seems to me that there is a very strong liability of the overriding of those fragments and the impairment of the ultimate usefulness of that joint. I do not know Dr. Sayre's position exactly, but it seems to me that the anterior angular splint, such as I was taught to use, applied to the anterior surface of the forearm, bringing the fulcrum of the lever at the apex of this splint at the elbow joint, tends to mold these fragments into bad position. Such a fracture can not be molded into position and held there, with the expectation of getting a useful joint, unless it is put in an anterior angular splint, such as we were taught to use in our earlier days; consequently it is my belief that these fractures ought to be treated in the rectangular position.

DR. HORTON, Texas—I have but one or two remarks to make with reference to the discussion and the position of the gentlemen on the two sides of this question. In the first place we should recall the principles laid down in all questions of fracture; first muscular relaxation, afterward the position of the broken fragments. I understand that that underlies all questions of treatment. So far as we can judge of the questions which have been presented to us in the management of practice, it would be about this: That they rely upon the splint, whatever may be used, a straight one or an angular one, for the accomplishment of the entire cure. I do not so comprehend it. This is a principle which I want to impress upon our minds; that unless there is relaxation of the muscle and adjustment of broken surfaces, I care not what the treatment may be, you will certainly not secure a good result. And I ask you to think of that proposition a moment. After that has been accomplished I care not, so far as I am concerned what principle you may use; you may use a straight splint, and I have also used the angular splint as well. I have had a reasonable experience of forty-five years, and I have never had a bad result on the lines of that practice when I secured the principles laid down.

DR. THOMAS, Pennsylvania—If the fracture is above the condyle, a transverse fracture, I think a right angle splint is probably as good a splint as we can use, but if there is a fracture of one or the other condyle, as Dr. Allis says in his paper, and we put on the right angle splint, by removing the bandage and attaching the arm to that splint and the forearm, and pulling up the forearm, you take up the fractured condyle with it, and so long as you keep the forearm in the rectangular position it looks very pretty, but when you change it you find it can not be pulled up. Now you may talk about the relaxing process of the muscles, but in a fracture of one or the other condyles by placing it in such a position the condyle, itself, keeps the fragments in position. My experience is that my cures are almost perfect in that manner, while in the rectangular cases they are not.

The paper of Dr. S. E. MILLIKEN, New York, on "Tendon Grafting for the Deformities following Infantile Paralysis" was read by title, the author being absent.

Evidence of Appearances will not Discredit a Post-Mortem.—The fact of a number of persons meeting another from time to time, and observing nothing in his appearance to indicate ill health, the Supreme Court of Indiana holds, in the case of *Sharpe v. Commercial Travelers' Mutual Accident Association*, decided April 25, 1894, raises no conflict upon the evidence of a post-mortem examination, for example, as to the existence of fatty degeneration of the heart and brain.