Nor can we say that perhaps this man would develop dysentery later, for, although but a single specimen stool was examined from the majority of these cases, all the men have been under observation for a period of nine months and none of them has developed dysentery.

Several of the men have been examined at intervals of several months to see if the *Entamoeba coli* persisted in the stool for any length of time. In Case 15 of the Americans and in Cases 13, 31 and 49 of the natives amebas were always found at each examination. These cases were selected for continuous observation, for the reason that the parasite was present in abundance and easily found when search was made. In this connection it may be remarked that Craig,⁵ in refuting the view of Musgrave and Clegg that individuals presenting *Entamoeba coli* in health have never been observed for a long enough time to be positive as to whether or not dysentery might result, states that he has observed such individuals for a period of six months and has found these parasites constantly present.

CONCLUSION.

The *Entamoeba coli* can not be considered as the cause of chronic dysentery, since it is found as a continuous infection in about 50 per cent. of all normal individuals in many localities. It follows that no diagnosis of amebic dysentery should be made by the microscopist unless he has found the *Entamoeba dysenteriae* as distinguished from the *Entamoeba coli*.

The *Ankylostomum duodenale* is the most important of the intestinal worms found. This parasite seems to be very widely distributed among the natives in this locality, although I have not seen a single case of typical ankylostomiasis during a period of observation lasting ten months. This is apparently because the parasites were not present in large numbers in any individual. In the majority of the natives examined two or three ankylostoma eggs could be found in a single slide after a careful examination, but they are present in much larger numbers when the parasites are abundant enough to produce anemia.

Generally the development of the embryo in the intestine does not progress beyond segmentation of the egg, but it is worth recording that in one specimen, that of Native 14, examined immediately after its passage, numerous free swimming ciliated rhabditiform embryos were seen. This stage is not usually attained until one or two days after leaving the intestine. Ova in various stages of development, including advanced periods of segmentation, and eggs containing moving embryos on the point of hatching were also found.

A point of interest in diagnosis may here be made. No case should be diagnosed as ankylostomiasis and so treated merely because a few of the eggs of this parasite are found. To do this seems to be a tendency of most practitioners who come from a region where the ankylostomum is unknown to a locality where it is frequent. It is safe to say that no case should be diagnosed as ankylostomiasis unless presenting the characteristic symptoms of this condition, combined with a large number of eggs in the stool. In this connection I would like to quote Manson,⁶ who says: "It is not in every instance in which the ankylostomum is present that consequences so serious ensue. There may be dozens of ankylostoma in the intestine without any appreciable anemia, or, indeed, symptoms of any description what-

ever. Grave symptoms are the exception. One must be careful, therefore, to avoid concluding that the ankylostomum is the cause of every pathologic condition with which it may chance to concur."

The other intestinal parasites may be dismissed briefly as of little or no pathologic significance. The one case of tapeworm occurred in native No. 14, who was a Filipino and not a Mohammedan, and therefore was not prevented by his religion from indulging in pork. He was entirely unaware of its presence, and had not missed a day from duty as the result of five parasites that he harbored.

This wide dissemination of intestinal parasites among the natives is, no doubt, facilitated by the universal custom of the Moros, who are Mohammedans, of defecating in the river, and the fact that, excepting the few Americans living here, it is also a universal custom to drink this river water unboiled. The number of natives infected with one or another of the parasites considered above as compared with the number of Americans infected probably furnishes some index of their relative liability to other water-born intestinal affections. Thus out of fifty natives examined there were only two in whom no parasite of any kind was found, while out of the same number of Americans twenty-one had no intestinal parasites, making the relative liability ten times as great among natives as among Americans. Or, considering the ratio of the total number of infections in the natives to the total number in the Americans, we have 428 to 45—nearly three to one. We may say, therefore, that the native soldier is between three and ten times more liable to water-born intestinal diseases than the American soldier. The native soldier in this locality uses plain river water, while the American is furnished distilled and boiled water, with occasional relapses to the unboiled water during "hikes" and in Chinese restaurants.

While, to be sure, there are other methods of disseminating intestinal diseases besides the drinking water, surely the advantages of boiled or distilled water are thus made sufficiently plain.

A CASE OF RIGHT SUBCLAVIAN ANEURISM— ATTEMPTED ENDOANEURISMORRHAPHY (MATAS).

CURE FOLLOWING SIMULTANEOUS LIGATION OF COMMON
CAROTID AND FIRST PORTION OF SUBCLAVIAN.*

HOWARD LILIENTHAL, M.D.

Attending Surgeon to Mount Sinai Hospital.

NEW YORK.

A study of the classical paper by Matas¹ describing his operation for the cure of aneurism impressed me profoundly. The operation aims at the obliteration of the sac with comparatively little disturbance of the circulation of the part. After securing preliminary hemostasis by means of the elastic bandage, the aneurism is opened in the general course of the affected vessel and all of its internal orifices are closed by suture. The sac itself is then obliterated by rows or layers of sutures. In the case of sacculated aneurisms this procedure may be carried out without interfering with the lumen of the vessel itself. In fusiform aneurisms, however, it may be necessary to obliterate the affected vessel together with the sac, although in some cases the vascular lumen may be

5. American Medicine, vol. ix, No. 23, June 10, 1905, p. 939.

6. "Tropical Diseases," "Ankylostomiasis."

* Read before the Section on Surgery of the New York Academy of Medicine.

1. Matas: Annals of Surg., February, 1903.

re-established by suturing around a rubber tube, the ends of which have been placed in the two main openings of the sac (afferent and efferent). The tube is to be removed before the final sutures have been tied. His discussion concerning the possibilities of the method in dealing with aneurisms of the larger visceral arterial trunks not situated favorably for the employment of the elastic constrictor was very suggestive, and I determined to give the method a trial should the opportunity ever present itself.

The present paper contains the report of a case in which Matas' operation was attempted and, although it records a failure, the lack of success should not be ascribed to the method, but rather to the fact that the steps were not fully carried out. The apparently complete recovery of my patient after the employment of the older operation of proximal ligation is gratifying, but it will not deter me from making another attempt to follow out the principles of Matas if the occasion should arise.

History.—The patient was a Russian, 40 years old, and a tailor by occupation. He had served as a soldier in the infantry of his country. For the past ten years he had suffered not infrequently from what he called rheumatism. He stated emphatically that he had never been infected with gonorrhea or syphilis, and careful physical examination strengthened my belief in his truthfulness. For about a year there had been a stabbing pain in the precordial region, which had been pretty persistent but not progressive. In December, 1904, the precordial pain began to radiate to the right shoulder and down the right arm. Soon after this a localized swelling was noted in the right supraclavicular region and there was a general enlargement of the right arm and hand, especially noticeable in the fingers. The patient also complained of a prickling sensation in the right hand. He entered one of my wards at Mount Sinai Hospital, Feb. 4, 1905.

Examination.—The man's general physical condition and nutrition were good. The superficial veins over the right supraclavicular and infraclavicular regions were markedly dilated, running over the top of the shoulder. Dullness and rather high-pitched respiration were noted in the right infraclavicular region. Above the middle of the right clavicle there was a hemispherical, strongly-pulsating tumor, a little smaller than half a lawn-tennis ball. The pulsation was distinctly expansile in character and was accompanied by a marked systolic thrill. With the stethoscope a short systolic and a short diastolic murmur could be heard. The overlying skin was not discolored, nor was it adherent to the mass. The clavicle, too, seemed to be uninvolved. The right shoulder and the entire arm and hand were slightly cyanotic and were covered with numerous minute bluish vessels. The right arm and hand were swollen so that they were considerably larger than the left, but there was little, if any, pitting on pressure. The fingers of the right hand were markedly clubbed (Fig. 1). Pulsation at the right radial was somewhat delayed and was much weaker than at the left. There was no evidence of arteriosclerosis. The temporals were visible but were not abnormally tortuous.

Operation.—Four days after admission, February 8, the patient was anesthetized with chloroform and an incision was made from the outer fourth of the clavicle to the sternoclavicular articulation and thence for four inches along the anterior border of the sternocleidomastoid muscle. The skin flap was dissected off, the insertion of the sternocleidomastoid was divided and the inner half of the clavicle was subperiosteally resected. A temporary ligature of chromicized catgut was thrown around the common carotid artery, but not tightened, and the vessel was then followed down to its origin from the innominate. By working carefully and keeping close to the vessel, no important nerve or vein was injured. Exposure of the innominate was excellent and ligation of this vessel would have been easy. A temporary ligature of chromicized gut was now thrown around the first portion of the sub-

clavian about one-half inch from its origin, and was held by a clamp at a sufficient tension to obliterate pulsation in the aneurism. Pressure over the aneurism caused easy and complete emptying, but on release of pressure there was a gradual refilling of the sac. From the rate of this reflux, which was still further diminished by tightening the carotid ligature, I believed that it might be possible to open the sac and secure the mouths of the tributary and branching vessels, so as to treat the case by Matas' method. I accordingly pierced the most prominent part of the tumor with the point of the scalpel and cut toward the shoulder for about three-quarters of an inch. To my disappointment there was a steady, non-pulsating fountain of blood, which spurted vertically for about four inches. The insertion of my finger for purposes of exploration at once checked the hemorrhage. The only opening in the sac which I could detect was a large one, which occupied a position toward the shoulder of the patient and admitted the tip of the finger. There were no coagula. The lining wall of the aneurism was everywhere smooth. There was no indication of a tributary opening toward its cardiac side. Evidence

Fig. 1.—Photograph taken four weeks after operation. Note difference in size of the two hands, the line of the cicatrix and the asymmetry due to the transplantation of the right sternocleidomastoid muscle. (This photograph was kindly made for me by Dr. *Branower of the House Staff of Mt. Sinai Hospital.)

of irregularity indicating atheromatous degeneration or calcification was wanting. The anatomy of the aneurism, thus shown to be of the sacculated variety, may be understood from the accompanying diagram (Fig. 2).

The patient's condition, which had been excellent, now gave some indication of shock, and I determined to risk nothing further, but at once sutured the vascular wound with catgut and a hemostatic needle. The first portion of the subclavian was then ligated with double No. 3 chromicized gut after the method of Ballance and Edmunds. The carotid was also permanently ligated. The divided sternomastoid muscle was attached to the remainder of the clavicle and the wound sutured with drainage. The usual dry gauze aseptic dressing was applied and the hand and arm were enveloped in cotton and bandages.

Pain following the operation was slight, but for several days there was considerable reaction of absorption and some tender-

ness over the aneurismal tumor. The temperature rose as high as 102 F., but gradually subsided and the patient's condition never gave reason for alarm. He was kept in bed for twenty-two days and then was allowed to be about in a wheeling chair. The circulation in the right upper extremity was never poor, but there was naturally some local cyanosis and the radial pulse remained absent up to the time of his discharge, March 22, 1905, when the patient was presented before the New York Surgical Society. The cyanosis had by that time almost disappeared. Pulsation in the temporal artery returned faintly thirty-eight days after the operation.

Postoperative History.—The patient was seen about three months after the operation and the cure of the aneurism seemed to be complete. The tumor had disappeared, except a small, hard mass, and there was no pulsation in the subclavian region nor in the radial. The right hand had remained larger than the left, but was apparently progressing toward the normal. On testing the circulation by pressing and releasing the fingernails it was noted that the subungual blush returned almost, if not quite, as promptly as in the left hand. There was still slight disability of the shoulder from the resection of the clavicle, but the bone had in great part reformed.

Fig. 2.—Diagram of the aneurism.

In reviewing this case a few of the more interesting points may be discussed. First, the cause of the trouble. In the total absence of anything pointing to general arterial disease we may perhaps conclude that the most probable causal factor may have been an injury to the healthy vessel by a direct blow, possibly with one of the angles of the rifle stock during military exercise. The aneurism, however, had not the anatomic characteristics of the traumatic variety.

Secondly, we may discuss the modern methods of treating these large aneurisms when they do not occur in an extremity. To be sure, the result here was so excellent that one might be forgiven for following the precedent another time, and yet to leave a large blood clot in a vessel to be taken care of by Nature is not the best surgical practice. Not only is there apt to be serious reaction during absorption, but there is a distinct liability to infection of the clot. The method of Antylus, which consists in proximal and distal ligation with incision of the aneurism and turning out clots and with final packing of the cavity is one which has always been

considered extremely dangerous. Proximal and distal ligation without opening the aneurism has been practiced and at times with success,² while extirpation (method of Purmann) has been well thought of by many surgeons. These procedures, however, cut off important blood supply and we must not forget the consequent danger of gangrene of the extremity.

If it were possible to deal with these cases after the method of Matas, which would cure the disease by obliterating the aneurism without interfering with the lumen of the vessel, there would be a distinct gain. Thus far, to my knowledge, this operation has been done only when the elastic tourniquet could be employed. My failure was due to the fact that a sufficient number of tributary vessels had not been cut off. Ligation of the thyroid axis, the internal mammary and even the vertebral would not have been difficult nor directly dangerous. A temporary ligature of the axillary would also probably have been required to keep the sac empty enough to permit suture of its orifice (Fig. 2). Matas, however, calls attention to the danger of dissecting out the aneurismal sac, since its walls are nourished, not by the contained blood, but by small vessels in the surrounding structures, and necrosis of the sac is apt to follow its clean dissection. Bearing this warning in mind, I did not wish to prejudice the vitality of the sac by a dissection sufficiently free to ligate all the vessels mentioned. Having explored the interior of the aneurism, I was aware that, in addition to what had already been done, possibly a temporary ligature around the axillary alone would control the current, but by this time a considerable quantity of blood had been lost and I felt the need of some haste. Were I called to operate in a similar case, I would certainly control the axillary before incising the aneurism.

I believe that the Matas method would have proved perfectly feasible in this instance, because I am sure that neither the thyroid axis, the internal mammary nor the vertebral was connected with the sac (Fig. 2) and that, therefore, a free dissection of the aneurism with the consequent danger of necrosis would not have been necessary. The collateral circulation into the axillary must have supplied the blood which filled the aneurism after the subclavian and carotid had been tied.

Whether or not the incision and digital (rubber glove) exploration of the interior of the sac with its subsequent suture had anything to do with the prompt clotting and the speed of cure is an interesting question.

The entire subject of aneurism of the vessels higher than the axillary is too large to be treated here. The literature of aneurism of the subclavian is rather voluminous. Dr. Eben Alexander, Jr., of the house staff of Mt. Sinai Hospital, kindly secured for me notes on eighty-nine cases, most of which were operated on before 1890. Many of these aneurisms were of the traumatic variety and the operative procedures were various. The mortality rate in the entire number from all causes was 52.8 per cent. It is, in reality, probably much higher, since failures are unfortunately less likely to be reported than successes. It seems, however, from the figures of Jacobsthal,³ in an article on the operative treatment of subclavian aneurisms, that there has been a distinct lowering of the mortality rate since 1890, probably due to improvement in technic. Thus, before 1890, 55 out of 75 cases succumbed, showing a mortality of 73 per

2. Dr. J. A. Blake presented a beautiful case of aneurism of the subclavian treated by this method. See report of meeting of New York Surgical Society, Nov. 22, 1905.

3. H. Jacobsthal: *Deut. Zeit. f. Chir.*, vol. lxxviii, p. 239.

cent., while since 1890 only 4 out of 25 cases were fatal, or 16 per cent. This author believes in extirpation as the proper operative procedure and reports 20.8 per cent. cures: 5 in 24.

Confining this discussion to those operations in which there has been simultaneous ligation of the subclavian or innominate and the common carotid arteries proximal to the aneurism, I have been able to find the reports of 13 or, including my own, 14 cases, with 7 deaths, or 50 per cent. mortality. Sheen⁴ has reported 12 of these cases, by various operators, including one of his own, in which there was simultaneous proximal ligation of the innominate and carotid. The thirteenth is an old case by Liston.⁵ One of Sheen's fatal cases, however, was a secondary operation, ligation having been performed for hemorrhage following extirpation of a subclavian aneurism—operator, B. G. A. Moynihan. Omitting this, we have a mortality in 13 cases of 46.1 per cent. The interesting cases collected by Wyeth⁶ illustrating the results of distal ligation for aneurisms near the heart do not concern us here.

The following are my conclusions and suggestions of technic for performing the Matas operation in sacculated aneurism of the subclavian artery:

From an examination of the literature one may readily see that the greatest danger following the ligation of the large arterial trunks is that of hemorrhage, either because of erosion of the vascular walls by the ligation, or on account of infection and ulceration into the lumen of the ligated vessel. Speaking on theoretical grounds, hemorrhage is all but impossible after endoaneurismorrhaphy (Matas). By performing this operation instead of ligation or even extirpation we shall probably greatly improve the statistics. It is simple, and, although best suited to the sacculated forms, may be perfectly feasible in certain cases of the fusiform variety. In a sacculated aneurism of the right subclavian, such as the one here reported, the technic suggested is the following: The incision should be along the anterior border of the sternocleidomastoid muscle from a point about at the level of the cricoid cartilage to the interclavicular notch, thence along the clavicle to its outer fourth. The sternocleidomastoid may then be divided at its insertion and the common carotid artery exposed and followed to its origin in order to find the subclavian. One should remember that anomalies are sometimes found here. Temporary ligatures should now be placed on the carotid and subclavian and the aneurism carefully observed, both as to pulsation and as to the refilling of the sac after it has been emptied by manipulation. If pulsation persists, or if there is a rapid refilling, a temporary ligation of the axillary is necessary. There should be no free dissection of the sac and no exposure of the smaller subclavian branches, such as the vertebral, thyroid axis, internal mammary or superior intercostal. The aneurism may now be opened by an incision small enough to be plugged by the exploring finger and the interior of the sac carefully examined to ascertain the number of large openings.

The character of the hemorrhage must be noted. It will probably not be severe and may be found to proceed from a single orifice of moderate size. There will probably not be pulsation. Endoaneurismorrhaphy may now be performed as described by Matas,¹ and on

its completion the ligatures may be removed, first that of the carotid, second the axillary, and third the subclavian. If everything is in order the wound may be closed without drainage.

In dealing with aneurism of the left subclavian the steps must be changed somewhat to suit the anatomy.

NOTE.—Soon after this paper was sent to THE JOURNAL for publication, at a meeting of the Southern Surgical and Gynecological Association, reported in THE JOURNAL A. M. A., Jan. 6, 1906, page 66, Dr. F. W. Parham, New Orleans, stated that he had cured a case of left subclavian aneurism by suture of one orifice within the sac after the failure of proximal ligation.

Special Article

THE PHARMACOPEIA AND THE PHYSICIAN.

CHAPTER XIII.

ASTRINGENTS—Continued.

METALLIC ASTRINGENTS.

The insoluble salts of bismuth are admirably adapted for the protection of the intestinal mucosa from irritations of various kinds. While as much as thirty times the average single dose of bismuth salts has been given in the course of a day, by mouth, without any distinct evidence of untoward results, absorption and poisoning have followed the external use of bismuth salts as a dusting powder when used over a large surface.

Owing to the extremely poisonous nature of bismuth when absorbed, the soluble salts seem to us to be entirely superfluous, particularly as their solubility precludes the very object for which bismuth compounds are chiefly used—that is, for the mechanical coating of the surface with a protective layer of insoluble powder. It must be admitted, of course, that even the soluble salts of bismuth are astringent, but we have many safer astringents which are harmless that might well be used in their stead.

The activity of the nostrum-makers has resulted in placing on the market a number of preparations that are intended to combine the antiseptic properties of the benzene derivatives with the astringent properties of bismuth, but until we know more of the possibilities and the limitations of internal antiseptics it is preferable to use the simple astringents and to use our antiseptics separately.

Bismuth subnitrate, or subcarbonate, when triturated with water, should form a creamy mixture, and the physician should not permit the use of that which readily separates as a heavy powder, leaving a nearly clear stratum of liquid above it. The covering power of bismuth is in direct proportion to the creaminess of the mixture.¹

BISMUTHI SUBNITRAS.—U. S.—An insoluble white powder, without odor or taste.

BISMUTHI SUBCARBONAS.—U. S.—Closely resembles the subnitrate.

Average dose (of either salt): 0.5 gm. (8 grains).

BISMUTHI SUBSALICYLAS.—U. S.—This preparation closely resembles the subnitrate in physical properties and is intended to combine the antiseptic properties of the salicylic acid with the astringency of bismuth salts.

1. A typical prescription for gastric irritation, or ulcer is as follows:

R.	Bismuthi subnitratis	grs.	cl	10
	Mucilaginis acacie	3xv		50
	Aque cinnamomi q. s.	5ii		100

M. Sig.: Shake well and take one or two teaspoonfuls three or four times a day.

In cases in which marked irritability persists, the addition of 0.01 gm. (1/6 grain) of cocaine hydrochlorid to each dose, as a local anesthetic, may be advantageous, but we do not advise it as a routine procedure. For diarrhea one or two drops of the deodorized tincture of opium may be included for each dose, or, if preferred, 1 c.c. (15 minims) of paregoric, the official camphorated tincture of opium. For this purpose, it is usually preferable to give one dose every hour for three or four doses then every two to four hours as needed. Bismuth subcarbonate may be used instead of the subnitrate and tincture of kino, tincture of krameria or the compound tincture of gambir (catechu) may be added in the treatment of diarrhea.

4. Wm. Sheen: *Annals of Surg.*, July, 1905, p. 1; also November, 1905, p. 802.

5. Quoted by Wyeth in "Essays in Surgical Anatomy and Surgery," Wm. Wood, 1879.

6. Wyeth, J. A.: "Text-Book on Surgery," Appleton, 1888.