is also true, in a lesser degree, in other places. The beneficial effects of circular constriction with bandage and cast, has been over-estimated, while its dangers and disadvantages have been too little commented on, if they have been fully appreciated. Open dressings facilitate inspection, dressing, bathing, ventilation, massage, and any indicated treatment, the importance of which needs no emphasis. The attachment for immobilizing the hip-joint is applied in fractures of the neck of the femur, and also used in hip-joint diseases, and is on the same principle as that now in vogue. When the key is removed and the caps replaced, the splint is locked, and the adjustment can not be interfered with. The pneumatic pads are covered with satin-finished waterproof leather, and can be cleaned with warm water and soap when necessary.

The splints are easily worn under the regular clothing, and are so well constructed that they will last practically a life time.

They have been tested by the writer in some severe cases, with perfect results; among them, one case of compound, comminuted, multiple fracture of both legs. The bones in the right leg penetrated the skin in five different places, and in the left leg, in three. Several loose fragments of bone had to be removed. The usual dressings were tried, but had to be abandoned on account of the excruciating pain and stasis. Amputation of both legs was considered necessary by attending physicians, but the patient would not submit to it, and further counsel was called, resulting in the application of pneumatic splints to both legs. The pain ceased at once, on application. The absence of circular constriction allowed complete restoration of the circulation, and permitted of proper open dressings for the wounds. The splints were only worn thirty-three days; ankle and knee motion was in no way impaired; the union and result was perfect.

I mention another case, on account of the attending circumstances, one of simple fracture of the tibia. The patient was erecting a house at the time, and hauled lumber seven miles daily, without discomfort and with perfect result.

Before bringing the appliances to the notice of the profession, they were submitted to some of our most eminent surgeons, whose unqualified endorsement has demonstrated the unparalleled excellence of the treatment.

POSSIBILITIES OF LIQUID AIR TO THE PHYSICIAN.*

A. CAMPBELL WHITE, M.D.
NEW YORK CITY.

It is now over a year since I began the use of liquid air in the hospitals of New York, and less than a year since I published in the *Medical Record* a preliminary report of the results obtained by its use up to that time; and I am led to-day to accept your invitation to present this subject to your Section, no less by the pleasant recollections of the last time I appeared before you, than by the many communications I have received since publishing my report, from all parts of the world, demonstrating the lack of knowledge of the public as well as physicians as to what liquid air really is. I have received requests from physicians from nearly all parts of the United States, from Mexico, from Germany and even from New

Zealand, asking that a small package be sent and all expense would be paid at the other end. I have been asked whether it were given hypodermatically, and whether it would not be good taken in small doses internally for abscess of the liver. One physician, who has represented many drug houses, wanted to know if it could not be put up in small packages and retailed at the drug stores and whether he could not act as my Southern agent. Nearly all physicians say, in discussing the subject, "Oh, we have tried extreme cold before and know it is no good." Possibly there are some of you here this afternoon who have never seen liquid air, and if so, I shall demonstrate that you really never knew what cold was before. I intend at least to show you what liquid air really is, and have brought with me, through the courtesy of the Tripler Liquid Air Company, a generous supply in order that I may show you its physical properties and demonstrate some of its applications, as well as show you some of the appliances which more directly concern us.

After doing this, I desire to briefly tell you of some of the things we have accomplished with it at the Vanderbilt clinic of Columbia University, and elsewhere; and then call attention to some of the things which may be done in the treatment of internal diseases. [Dr. White gave several demonstrations with liquid air.]

In my former paper I told of the results of our attempt to kill germs with liquid air. With Dr. William H. Parks, we tested the effects of liquid air on typhoid, anthrax and diphtheria bacilli. In order first to test the effect of extreme cold on these germs without bringing them in direct contact with the liquid, we put into a number of capillary tubes pure cultures of virulent anthrax, diphtheria and typhoid bacilli. These capillary tubes were marked for identification, sealed at both ends, and dropped into a glass of liquid air. The liquid was renewed at intervals, so that the tubes were at all times completely submerged. Some were removed in thirty minutes, others in forty-five minutes, sixty minutes, and ninety minutes. After they were removed from the liquid and the sealed ends were broken, they were dropped into separate culture-tubes containing fresh sterilized bouillon, and these in turn marked for identification. These bouillon tubes were placed in incubators kept at a temperature of 38 C., where they remained for over forty-eight hours, when they were all examined and found to contain pure cultures of anthrax, typhoid and diphtheria bacilli according as they had been inoculated. In order to test whether liquid air itself in addition to its temperature would have any effect on germ life, a culture-medium containing numerous colonies of anthrax bacilli was dropped into a glass of liquid air and allowed to remain over an hour. When removed, a fresh sterile culture-medium was inoculated with the one exposed to liquid air and placed in the incubator as before. The same result was obtained as when the capillary tubes were exposed.

Dr. J. H. Huddleston tried the effect of liquid air on vaccinia virus, and found that fifteen minutes' or less exposure to the liquid air had no effect whatever on the power of the virus to produce vaccination.

Longer exposures to liquid air have been tried with the same result. Dr. Park has recently made further experiments, making the exposure over two hours and exposing all kinds of germ life in form of pure virulent cultures, from hay infusion to anthrax bacilli. With this exposure, taken all together, from the weakest germs to the strongest, it kills from 98 to 50 per cent. Of the typhoid germ it kills about 75 per cent. in two

* Presented to the Section on Diseases of Children, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.

hours' exposure, the staphylococcus about 50 per cent., and the diphtheria typhoid, about 65 per cent. This exposure does not completely destroy the virulence of any pure culture, but of course the activity or virulence of all these germs is suspended for a long time after exposure to the liquid air, and they are only brought back to activity by the most careful handling and under the most suitable conditions. Professor Dewar, of London, is reported to have given bacteria exposure to liquid air for two weeks, with practically the same results.

Now the question is, knowing these facts, what is the effect of liquid air upon germs existing in the living tissues of the body. We know that the human organism is the natural enemy of the bacillus. We have reason to believe that the human system develops an antitoxin when invaded by the ptomain-producing bacteria, which under most conditions is successful in withstanding this invasion. Is it not possible that the temporary suspension of the vitality of these germs, due to the application of liquid air, gives nature the opportunity to completely destroy them, particularly when we remember that the local circulation is much increased shortly after the application.

Buchner,¹ in an article on the natural defenses of bacteria, speaks of the results obtained by us in the use of liquid air, and attributes all our success to the subsequent local hyperemia produced.

CLINICAL USES OF LIQUID AIR.

Liquid air, applied to the healthy living tissues, causes a temporary local anemia accompanied by anesthesia, or a slough, depending upon the amount used.

The results obtained by an excessive application are best explained by the experiment performed on the rabbit, before we had used the liquid air on the human body. About three-quarters of the ear of a rabbit was completely frozen by placing it in liquid air for some minutes. The ear was frozen so stiff that a small piece of the tip was broken off with the fingers without pain or hemorrhage. In less than ten minutes' time the circulation was entirely re-established with hemorrhage at the point where the piece had been broken off; the frozen part had regained its normal temperature—possibly a little higher—and color. The ear remained perfectly normal for about five days, when it began to swell somewhat, and by the end of two weeks about one-half of the frozen part had shriveled and dropped off. The remainder of the ear regained its normal condition. The part to which the liquid air is applied—with the spray—immediately loses all the blood, becomes perfectly colorless and is accompanied in the application by no pain excepting a slight tingling sensation which is only felt at the very beginning of the application. If only a slight amount is used the circulation almost immediately returns, the part becomes congested and frequently elevated, looking like a wheal or urticaria; this subsides soon and may or may not for a while itch to a slight extent. The anesthesia lasts not only while the anemia exists, but for some time after, owing to a temporary paralysis of the sensory nerve terminals. A freezing sufficient to produce anesthesia for ten minutes can be given without causing any more destruction of tissue than a slight desquamation of the superficial epithelium, which in no way interferes with primary union.

I recently did an operation under the influence of local anesthesia produced by liquid air, for the purpose of relieving a deformity of the hand resulting from a

poorly-treated burn of childhood. The hand was freely incised, and the fascia cut here and there, relieving whatever adhesions existed. The operation caused no pain, the result was perfect, and primary union was in no way interfered with. I could mention any number of cases equally successful where liquid air has been used as an anesthetic.

I think you can readily appreciate in how many cases this anesthesia would be preferable to any other, for example, opening of an abscess, where there is always an element of some danger in the use of cocaine; incising a paronychia; and of what great value it would be in a cardiocentesis, paracentesis thoracis or abdominis, where the vital functions are already at so low an ebb as to preclude the use of anesthetics.

In this connection I will just mention the almost invariable success resulting from the application of liquid air in the same way over the spinal end of the nerve in neuralgias and herpes zoster. The pain ceases almost immediately; rarely is more than one application necessary, and the only bad result has been some ulcerations at the point of application in some patients of advanced years, where the vitality of the skin was exceptionally low. These ulcerations readily heal with proper treatment, and are, as a rule, a pleasant substitute for the pain of neuralgia and herpes zoster.

As a cold application to allay or abort an acute inflammation, its use is ideal, and should be applied intermittently with the roller. The same would apply to an acute adenitis—not neglecting to treat the cause.

In the treatment of a boil I would use the spray, and, if just beginning, I would completely freeze it, possibly pricking it while frozen here and there with a needle, in order to relieve any subsequent congestion. If pus to any extent had formed I would freeze and make small incisions. It will always abort a boil if taken in time, although of course not preventing others forming.

From my experience in the local treatment of carbuncle with liquid air I am satisfied from every point of view that this is by far the best form of treatment. It is less painful to the patient than any other form of treatment. Only one application is necessary; within twelve hours of the first application, the pain entirely ceases, not to return again, and at the end of a few days only a small ulcer is left—representing the tissue between the openings of the carbuncle—which readily repairs with usual dressings. In the treatment of the carbuncle the spray is used, first projecting it into the openings and using the air quite freely, then quite thoroughly freezing the external surface, which must be well cleansed of discharge resulting from sending air inside of the carbuncle before freezing. After freezing, the carbuncle should be dressed with a dry absorbent dressing, so that the discharge, which will be abundant and accompanied with considerable bleeding, can be readily absorbed. The reaction from the freezing takes place in about twenty minutes, and it is to this extreme hyperemia that I attribute the success of liquid air in the treatment of this affection more particularly.

Before using liquid air in lupus we anticipated success, and our expectations have been realized, both in the treatment of the erythematous form and in lupus vulgaris. In the former variety we use the cauterizing instruments frequently, not freezing to any great extent at each application. Excessive desquamation results after each application, until good healthy skin appears, slightly reddened. The applications are then stopped and the redness soon disappears. Lupus vulgaris is treated rather generously with the spray once

1. Münch. Med. Wochschr., Oct. 3, 1899.

or twice. This causes a slight acute inflammation, which results in a healthy ulcer that quickly heals.

As a stimulant in the treatment of chronic ulcers, varicose and non-specific, one application, not to the point of freezing, but intermittently for about five minutes with the spray, is of great utility.

Liquid air as a cauterizer is probably the first thing that would occur to us after having seen and handled it, particularly for the removal of the benign foreign growths. For this purpose we now use liquid air to the exclusion of almost all other methods. For the quick and comparatively painless removal of a nevus, without any resulting scar, it is not equaled by anything we now have at our disposal. I have already shown you the **instruments for its application** in this direction for the destruction of adenoids, polypi, nevi and other hypertrophies.

The study of carcinoma has been universal and popular during the last few years, and this study promises to disclose the true cause of this dreadful and undoubtedly increasing malady. No doubt when its true etiology has been established we will find some treatment which will prove either prophylactic or curative. I think we must all have been impressed who have seen many of these cases, with the utter futility of almost every method of treatment at our disposal; even surgery seems to give in nearly all cases only a temporary relief, to be followed by even more malignant growth. The consideration of cancer is certainly not a subject of discussion for this Section, but I would not be doing justice to my subject did I fail to mention the great good liquid air has done in these cases. Even in cases in which the treatment has failed to cure, which were far beyond operative interference, where the patient has had two or three operations, where the patient has had every "special treatment," quack and otherwise, in the country; even in these cases, the great good done in the relief of pain and the destruction of bad odor, both so characteristic of this disease, places liquid air in efficiency far above any other treatment we have to-day.

In my previous paper I said I could make no claim for liquid air in the treatment of cancer, for my limited experience at that time, though promising, did not warrant my offering any encouragement; I can truly say to-day that I believe that epithelioma, treated early in its existence by liquid air, will always be cured, and that many cases inoperable can also be cured by its application. Prof. George H. Fox, in one of his lectures in the College of Physicians and Surgeons, New York, presented last autumn one of the first cases treated at his clinic by liquid air. This patient, a woman some 60 years of age, had an epithelioma on the right face, involving the outer canthus of the right eye. Professor Fox, in presenting this case, said, "No treatment at our disposal could have destroyed an epithelioma of this character in so short a time and with so slight a scar remaining as liquid air has done in this case." This subject requires a paper devoted entirely to its consideration, and I am sorry I can not present the history of some of these cases in detail, those which have recovered and those less fortunate, for every one bears testimony to the efficiency of liquid air in the treatment of carcinoma.

When we come to consider the use of liquid air in the treatment of non-cutaneous diseases, there is a very broad field open to us and many obstacles present themselves which are difficult to overcome. I think it will at once occur to us all that we ought to be able to use this cold, dry vapor given off by liquid air, for the reduction

of temperatures in fevers in something the same manner as we now give hot air and steam baths to promote activity of the skin. We are now having made an apparatus which we hope will accomplish this result; if so, it will be much more simple than the cold plunge and be much more pleasant and agreeable to the patient, accomplishing at the same time the desired results.

I have also thought that the inhalation of this vapor, which is almost pure oxygen, would be of great value in the treatment of certain diseases of the air-passages, such as hay-fever, acute laryngitis, tubercular laryngitis and possibly pulmonary phthisis. For this purpose I am having made an inhaler which will carry the vapor well back over the glottis before it becomes warm; and, furthermore, this vapor can be medicated, and in this way carry with it some remedy which might possibly add to its efficiency. The results of this treatment are, of course, purely speculative, but I anticipate success in this direction in the treatment of at least some of the diseases mentioned.

I regret that time does not allow me to go more fully into details regarding some of the cases I have treated with liquid air; but if I have given you all a fair idea of what liquid air is, what it looks like and what some of its possibilities are to us and to our patients, I shall feel well repaid for undertaking this demonstration, and I hope the discussion which is to follow will bring out many new points we have not thought of before.

DISCUSSION.

DR. LOUIS FISCHER, New York City—It seems to me that the question of local anesthesia is one of great importance to pediatricists, as for instance in the treatment of scrofulous manifestations, local abscesses, suppurative adenitis, various forms of peritonsillar abscess and the removal of adenoids of the pharynx.

DR. WAIER.—In the treatment of carbuncle, is your object to establish a slough?

DR. A. CAMPBELL WHITE—There are usually four openings around the center, forming, so to speak, the corners of a square. These corners will slough, but not the edges. The good results in carbuncle I believe to be partially due to the fact that these openings exist. The excessive congestion which comes on after the application of the liquid air is relieved by a free discharge through these openings. There is only a temporary suspension of the activity of the bacilli.

DR. L. J. LAUTENBACH, Philadelphia—Did you freeze and operate with the knife upon the case of epithelioma at the side of the eye?

DR. WHITE.—No, we used the liquid air cauterizer and simply closed the eye. When liquid air strikes the skin it is for the moment at a temperature of 312 F. below zero. I should freeze slightly, then wait and repeat the operation perhaps three or four times. The woman referred to had twelve treatments. There was a marked and edematous inflammation of the side of the face. It subsided in about three days. After that the improvement was very decided. This has been true of almost all cases. I have applied liquid air in epithelioma of the tongue. Here there is so much moisture that inflammation is easily excited, so that this agent must be used with great care. In two cases, one in the anterior part, and the other in the posterior part of the tongue, I have used the liquid air. After about three or four applications it was necessary to temporarily stop treatment because of the inflammation induced. Great care must be observed in posterior cases for fear of the possibility of edema of the glottis arising. Almost no sloughing follows the application of liquid air. The destruction is very superficial as we use it. In a case of carcinoma of the face we break up the intercellular substance of the epithelioma by freezing, and destroy the nutrition of the superficial cells. When the patient first returns there is a small ulcer, and we repeat the application. Eventually there is no further return. I would not dare to put a glass instrument into a child's throat

for removing adenoids; I should prefer to make use of one made of thin aluminum. I have treated erysipelas by filling a glass globe with liquid air and rolling it over the surface. For local refrigeration I have made use of a brass roller into which liquid air is introduced, and then the roller is passed over the part.

MOVEMENTS OF INTESTINES.*

ALBERT BERNHEIM, M.D.

PHILADELPHIA.

During treatment of a woman for floating kidney, nervousness and general debility, I administered—in the course of a rest-cure—enemas of cottonseed-oil; one time, about five hours after the injection, the patient vomited, the vomit for a greater part proving to be cottonseed-oil, this incident showing the possibility of a retrograding current of the oil from the rectum to the stomach. These oil enemas were administered for over two weeks and it was only this one time that oil was vomited. I was induced by this fact to make some experiments with injections into the rectum of a negro; these will be mentioned later on; besides I used a black-and-tan dog for experimenting, and, later on, in treating a woman for mechanical gastric insufficiency with a subacid condition, in which vomiting was not infrequent, the nutrient enemas used served as experiments.

It is for the research into the consumption of such nutrient enemas that various investigators experimented on the movement of the intestines, to learn of possible antiperistalsis of the gut.

Celsus (3-5 B. C.) mentioned the efficiency of nutrient enemas; in 1869 Voit and Bauer, in Munich, investigated their value as to absorption in the body, and found that an egg-emulsion, with the addition of a pinch of sodium chlorid, will be easily absorbed by the mucous membrane of the rectum and colon. Eichhorst, two years later, observed the same results. Czerny and Latzenberger, experimenting on men, found that not only albuminates, but also fat in emulsion and paste of starch, would be absorbed, when introduced into the rectum; they could not determine whether the starch is converted into sugar or not before the absorption, and it remained doubtful where the absorption occurred. Leube thought it necessary to peptonize the enema by mixing it with pancreas before injecting it into the rectum. Ewald, making a greater series of experiments, declared the peptonizing process superfluous, finding that an emulsion of egg-albumen will be absorbed even without the addition of sodium chlorid. Huber took an intermediate position between Voit and Bauer and Eichhorst on the one side and Ewald on the other; he recognized the fact of absorption without sodium chlorid, but the absorption was better with an addition of salt. Hemmeter explains the absorption without an addition of sodium chlorid, because this salt is naturally present in the egg. The absorption of the enemas was always said to occur in the lower and lowest parts of the bowels. L. Landois, in his book on physiology, says that the ileocecal valve prevents in general the contents of the colon from re-entering the ileum. When liquid masses are injected gradually into the rectum through a tube, they may get above the ileocecal into the ileum. In 1884, Nothnagel saw antiperistaltic movements of the gut when he touched the exposed intestinal section with a crystal of sodium chlorid or injected a salt solution by means of a Pravaz syringe. Grützner thought the fact of this antiperi-

stalsis a chief point as to the effect of nutritive enemas. His experiments were made on white mice, rabbits, guinea-pigs and cats, which had hungered for twenty-four hours. The injection mixtures consisted of a physiologic salt solution, with animal coal or starch or finely cut horse-hair or sawdust or poppy seeds. Four or six hours after the injection Grützner found the introduced particles in the stomach and in the entire course of the gut, except in the rectum. When, instead of salt solution, distilled water or a solution of .1 per cent. hydrochloric acid or of .6 per cent. potassium chlorid had been used, the introduced particles could not be discovered in the upper parts of the alimentary canal. Grützner at the same time took such precautionary measures that the animals could not have swallowed any excrements. The same positive results were found by him in experiments on man, in whose rectum he had injected starch emulsion.

Christomanos imitated these experiments and found that his researches corresponded to those of Grützner, if he followed the latter's method, but if he subjected the animals to extraordinary measures, such as severing and ligating the esophagus or fastening them down to the vivisection table for up to six hours, or applying the mouth-clamp, the experiments were negative. The latter series of experiments were made under rather pathologic conditions, and therefore can not prove what they seem to show. His experiments were made with carmin, Prussian blue, lycopodium dyed red or blue, and alcan-na-oil, on white mice, rabbits, frogs, pigeons and men. Only two out of ten experiments made on men were positive; in these cases he used either lycopodium or bismuth subnitrate, but lycopodium is often unseen in the microscope, and the bismuth crystals are not always to be recognized as such, even if stained black through the abode in the intestines.

Dr. Swiezynski, in the clinic of Professor Riegel, tried the same kind of experiments on men and dogs. In 12 cases, unstained lycopodium had been introduced 7 times, in 6 instances in a physiologic salt solution; the 6 times were positive; in the seventh case 1 grain of lycopodium; in the eighth and ninth cases, lycopodium stained with nitrate of silver in a salt solution was used, and in both instances grains of lycopodium were found in the stomach. In 2 other cases lycopodium stained with gentian violet was introduced, once with sodium chlorid and with positive result; the other time without addition of salt was negative. In the twelfth case bismuth subnitrate was used; this experiment could not be regarded as absolutely positive. Six other experiments were made with gentian violet lycopodium, 4 times in sodium chlorid solution, and twice without the salt; these were more or less positive.

Dr. Dauber made 17 experiments on white mice, white rats, rabbits and dogs, with indigo or cinnabar; 6 experiments were positive, 11 negative, the positive result occurring in cases where animals were subjected to rather physiologic conditions; the 11 negative ones, I should think, were due to a too severe restraint, whereby the animals were subjected to a pathologic condition. The experiments on animals under physiologic but precautionary measures are, in my opinion, only proving if positive, and not disproving if negative. The chief proof must lie in the human experiment. Hemmeter regards this Grützner antiperistaltic motion as only a very feeble marginal ascending movement, effected by surface contact of the particles with the epithelium, which in turn is moved by the muscularis serosa; further, he says, it is never visible to the eye, and it is not

*Presented to the Section on Physiology and Dietetics, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.