than the central portion. Deep within the tissues are found the acinous glands. The arrangement of the blood vessels is somewhat fan-shaped. J. Hynitzsch has observed villous elevations within the crypts which show a great similarity to the papillomatous proliferation seen in mammary and ovarian cysto-adenoma. The same author has made a study of the cysts occurring in the hyperplastic pharyngeal tonsil and divides them into two varieties corresponding to the two types of epithelium lining the crypts, namely, the columnar and squamous cysts. These columnar epithelial cysts vary in size from a microscopic body to that of a bean, occur frequently and contain a clear ropy mucus, often rich in leucocytes and occasionally red blood corpuscles. In two cases the cysts contained multinuclear elements or giant cells. Their nuclei vary in number, are of irregular arrangement and never situated along the line of the cell, thus essentially differentiating them from giant cells found in tuberculosis. Squamous epithelial cysts are much rarer. These cysts frequently contain horny lamellæ concentrically arranged, the center composed of amorphous granular masses, fat globules and cholesterol crystals. They may be regarded in all instances as due to retention of secretion by closure of the mouths of crypts. The process of involution may be explained as follows: Maturing of the embryonal tissue takes place first in the oldest portion of the growth, which is probably at the base of the tonsil. This process, together with the peculiar fan-shaped arrangement of the blood vessels may be seen in Figure 4, and here, too, we may gain a hint as to the cause of hyperplasia in certain growths. The firm connective tissue in which the blood vessels of the base are embedded prevents diminution of their caliber and this interferes with the normal decrease in the nourishment of other portions of the growth. We thus have a relatively increased blood supply to that portion of the growth supplied by the blood vessels affected. The change of embryonal into adult connective tissue takes place in the interfollicular tissue. Each follicle is enclosed in a bag, as it were, of maturing connective tissue; this gradually contracting tends to cause the follicles to disappear by pressure atrophy.

CONCLUSIONS.

1. The tissue composing adenoids has a wide distribution throughout the body.
2. The epipharyngeal tonsil is an organ normally present in all individuals.
3. The function of the epipharyngeal tonsil is at present unknown. Whether it is simply an evolutionary vestige or has to do with metabolic or other processes is equally uncertain.
4. Its wide distribution among the various members of the animal kingdom, its embryological connections, and its identity with lymphoid structures elsewhere in the body, would suggest that the epipharyngeal tonsil has a function.
5. The histology and the experiments of Massini and Genta would tend to place it in the class of glands having an internal secretion.
6. The relation of tuberculosis, syphilis or scrofulosis to adenoids is not etiologic. Adenoids may be affected with tuberculosis, etc., but only as any other organ may be similarly affected.

Important Scientific Post for a Woman.—Since Nencki's death, his assistant for many years, Miss Sieber-Schumord, has been appointed director of the Institute of Experimental Medicine at St. Petersburg, in his place.

THE ROENTGEN RAYS

IN DIFFERENTIATING BETWEEN OSTEOMYELITIS, OSSEOUS CYST, OSTEOSARCOMA AND OTHER OSSEOUS LESIONS, WITH SKIAGRAPHIC DEMONSTRATIONS.

CARL BECK, M.D.

NEW YORK CITY.

In periostitis, as well as in osteomyelitis, the skiagraphic signs are well marked. Abscesses can not only be localized, but their extent is so well outlined that the technical steps of the operation can be definitely traced in advance. The feeling of security the surgeon has while proceeding under the mentorship of the skiagraph gives a satisfaction unknown in former years, when often the whole femur had to be exposed simply in order to ascertain whether all foci were detected. If the Roentgen

Fig. 1.—Osteomyelitic focus in the middle of the humerus.

rays show but one focus, no other regions of the bone need be attacked.

In such cases a preceding trauma often opens the avenue of infection. The pain, the edema, the fever, and general debility may be sometimes so little marked that differentiation becomes difficult. The skiagraph not only clears this difficulty of diagnosing this disease, the true etiology of which is still so obscure, but it also furnishes a trustworthy guide for the operative technic at the same time.

Osteomyelitis is of a decidedly infectious character, generally due to the invasion of the staphylococcus aureus, which fortunately has a tendency of forming circumscribed foci in the vascular tissues of the bone, viz., the medulla, and sometimes the periosteum. The predilection of osteomyelitis is for the long bones of young

individuals. It is self-evident, therefore, that the early recognition of osteomyelitic foci renders the prognosis of their evacuation extremely favorable.

In the case of a lady of 20 years, the slow onset of the symptoms did not seem to indicate an acute inflammatory process. Pain at the thickened area of the middle of the humerus, being present only temporarily, the fear of a malignant growth was apparently not unjustifiable. The skiagraph at once did away with all anxiety, since it revealed the presence of periostitic proliferations and a circumscribed osteomyelitic focus (Fig. 1). The focus was easily exposed by the chisel, under the mentorship of the skiagraph. That the skiagraph has also spoken the truth by demonstrating the integrity of the remaining portions of the humerus, was shown by the speedy recovery of the patient.

The focus was distinguished by its light shade in the midst of the dark shadow of the cortex. The regularity of the cortical line and the external proliferation distinguished it from osteosarcoma, and the absence of distension from osseous cyst.

Necrosis and other later stages of inflammatory processes can be represented still more distinctly. The size and shape of sequestra can easily be made out. It can,

introduction of a probe did not reveal the presence of rough bone, and we were inclined to suppress our suspicion of the presence of a sequestrum. Our surprise was great when the Roentgen rays revealed the presence of a large splinter exfoliated from the inner portion of the radius, the surface of which was covered by osteophytes (Fig. 2). The direction of the skin incision, a slight oblique, was dictated by the position of the sequestrum as shown by the skiagraph. When the sequestrum was reached it was found to be covered by thick fibrous tissue at the upper surface, while the inner and lower surfaces were exposed. This explains why the introduction of the probe gave no positive information, since it had touched only the fibrous cover and did not come in contact with the rough lateral or posterior surface. Recovery was perfect eleven days after the operation.

Fig. 2.—Sequestrum, exfoliating from radius.

furthermore, be ascertained how they are located in their bony coffin and whether they still adhere or are exfoliated. Under the guidance of the Roentgen rays extraction is very easy.

In the case of a man of 23 years, who had crushed his left little finger, amputation was deferred until septic tenonitis and tenosynovitis had developed. The extensive tissue necrosis in the muscular interstices of the forearm necessitated free and deep incisions, which showed the radius, as well as the ulna, denuded of its periosteum. Amputation was therefore authoritatively advised, but nevertheless the chances for further conservative treatment were taken. Fortunately, the process became confined to the forearm and recovery seemed to make rapid progress. Only a small fistula at the dorsum of the forearm did not close. The repeated

Fig. 3.—Arthritic changes in the enlarged external condyle.

The regeneration of osseous tissue can also be well studied in such cases by the skiagraph. Foci of the same character are sometimes formed in typhoid fever. They must be treated after the same principles.

In diagnosing inflammatory processes in the joints, great difficulties are sometimes offered. As stated before, in the case of acute rheumatism, the integrity of the articular outlines are well marked. The same applies to acute inflammatory processes due to infection. In the latter event the distension of the joint by the serous or purulent effusion may be represented by the skiagraph. In chronic rheumatic processes the articular bone line appears somewhat irregular. In arthritis the contours of the bone epiphyses appear irregular, like indentations on some portions, while others are veiled. The arthritic deposits are recognizable as light shadows

of the deformed epiphyses, as they consist of translucent uric acid salts, while their periphery is distinguished by a dark sphere. In arthritis deformans the osseous proliferations are especially abundant and are well represented by the rays.

Figure 3 illustrates the case of a laborer of 50 years, who sustained an injury of his elbow eleven years ago. He reported that recovery took place after several months and that the elbow had remained stiff ever since. During the last few years inflammatory signs had manifested themselves, which were regarded as rheumatic. No other joints were involved. Since then he also had repeated attacks of pain in the elbow joint. When I examined the patient for the first time I found the elbow very much thickened and fixed at a sharp angle. Pressure below the external condyle caused intense pain. Crepitus, so often found in old arthritic processes, could

mean sometimes of differentiation from other affections, the clinical signs of which resemble it. The walls of an intra-osseous focus appear thickened. Some portions are translucent and their contours irregular. The articular outlines of a tubercular joint have lost their regularity and appear diffuse, cloudy and sometimes shaggy. In later stages, when cheesy foci have formed, for instance, their areas appear translucent. The cortex is sometimes entirely destroyed and leaves the impression of having been scooped out with a gouge. In most cases, especially when the soft tissues are also affected, atrophy is marked. Figure 4 represents the feet of a man of 58 years, who suffered from tuberculosis of the ankle joint for two years. He was treated by the injection of iodoform-glycerin and the administration of guaiacol carbonate. The left leg recovered entirely, leaving considerable deformity and

Fig. 4.—Tuberculosis of both ankle-joints.

not be produced in this instance, as the joint permitted no motion. There were no indications of tuberculosis, syphilis or gonorrhea. The skiagraph revealed the presence of mal-union (sideward displacement) of the coronoid process of the ulna, which probably had given the first impetus for the development of the arthritis deformans, the light shaded changes of which are especially well marked in the external condyle of the humerus. The left condyle shows synostosis with the olecranon. Removal of the projecting fragment by the chisel, separation of the adhesions, and the partial resection of the external condyle, the seat of predilection for the acute attacks, was advised as a therapeutic means.

In tuberculosis of the bones and the joints the Roentgen rays not only give information as to the seat and extent of the tubercular areas, but also offer the only

ankylosis only. The astragalus shows deformity and extensive destruction. The right leg is in an acute inflammatory stage, cheesy pus being present in the ankle joint. When there is calcareous degeneration the foci appear dark-shaded.

In the case of extensive tubercular destruction the eroded and displaced cartilages can be studied. In tubercular coxitis, the spontaneous upward dislocation of the femur and the separation of its head in the acetabulum can also be easily recognized.

The only marked signs of tubercular spondylitis consist sometimes in the presence of an abscess below Poupert's ligament, the nature of which would not be properly interpreted, if the plate did not prove the existence of vertebral changes.

It need not be said that the early detection of a tuber-

cular focus enables the surgeon to do a conservative operation, while at the late stage of extensive destruction such efforts are futile, as it is sadly illustrated by skiagraphic examination.

By realizing that so-called osteosarcoma is the most frequent of morbid growths and that of all tumors sarcoma offers the gravest prognosis, the importance of a thorough diagnosis need not be emphasized. The matrix of osteosarcoma, like that of all osseous growths, is either the periosteum or the medulla, in combination with the tissue originating from their proliferation.

Periosteal sarcoma is of a moderate hardness and contains either round, spindle or polymorphous cells. It attaches itself to the bone laterally, but may, in its further development, encircle it entirely. Periosteal sarcoma may become a real osteosarcoma at a later stage, when osseous trabeculae are formed. The skiagraph of periosteal sarcoma is characteristic, since it shows fine spiculated trabeculae that radiate from the surface. Periosteal sarcoma spreads rapidly and is highly malignant. Whenever the diagnosis of periosteal sarcoma is made, amputation should be insisted upon.

Sarcoma originating from the medulla is called myelogenous and is of a less malignant character. It may

fracture of the carpal end of the radius. But the consistency of the epiphyseal end was soft. The skiagraph failed to show the evidence of bone tissue, only one small remnant being left at the outer aspect of the radius. Resection was advised, but before the patient submitted to it another month had elapsed, during which time the neoplasm had grown to a great extent. The result was reported as being fair, eight months after the operation.

The hard myelogenous variety generally called endosteal or central sarcoma, also shows the ordinary sarcomatous structure. Its distinguishing feature is its fibrous texture and the presence of spindle cells. Some portions contain various tissues; the spindle-cell tissues often contain giant cells. If smaller or larger bone trabeculae are produced, it is called osteosarcoma proper;

Fig. 5.—Osteosarcoma of upper third of humerus.

be classified as soft, hard, alveolar and multiple. The soft myelogenous variety shows the ordinary sarcomatous texture. Its predominating feature is the presence of round cells. It has a decidedly more benign character than the periosteal type. Therefore it justifies a conservative attempt, that is, extensive extirpation. It produces carious destruction of the spongy portion which may cause spontaneous fracture. At a later stage the osseous shell will yield to the spreading sarcomatous tissue. This variety has a predilection for the long bones, especially their ends, and predominates at the lower epiphyses of the femur, tibia, humerus and radius. Skiagraphs of the soft myelogenous variety show the absence of osseous tissue, small fragments of it being sometimes left here and there.

In the case of a woman, age 28 years, who had fallen on her hand when it was in dorsal flexion, the faint outlines of boneshell in a soft myelosarcoma were shown. The swelling resulting from the injury produced the impression that a fracture of the carpal end of the radius was sustained.

Three months after the injury, when the patient came under my observation for the first time, a small deformity was noticed just as it is observed in a badly united

Fig. 6.—Osteosarcoma of lower end of radius.

if there are calcareous deposits, petrifying sarcoma; and if the tissues become vascular, a telangiectatic sarcoma will be formed, so that it may be mistaken for an aneurysm.

In later stages, when there is a regressive metamorphosis, fatty or cystic degeneration may take place. In this case, neoplasms that occur especially in the femur, tibia and inferior maxilla, may attain an enormous size.

The skiagraph of osteosarcoma proper (Fig. 5) shows more osseous tissue than the former variety, but its outlines are very irregular. Osteosarcoma proper usually commences near the epiphysis of a long bone. Skiagraph No. 6 shows the irregular sarcomatous proliferation of the lower one-third of the radius in a woman of 40 years. Treatment was not resorted to until two months afterward, when the skiagraph (Fig. 7) re-

vealed the destruction of the lower third of the radius and of a large portion of the carpus. Resection was performed; the result was perfect, as was illustrated by another skiagraph taken more than four years after the operation, which showed the regeneration of the osseous tissue.

The alveolar variety is characterized by its alveolar stroma, which contains nests of large cells. They have a predilection for the bones of the skull and the trunk. The multiple variety (also called myeloma) is characterized by the presence of numerous whitish foci, which consist of small round cells. It has the same structure as the lymphoid sarcoma and is nearly exclusively found in very old individuals, for whose skull and trunk they show the same predilection as the former variety.

The skiagraph of the alveolar, as well as the multiple type, shows the foci as light irregular shades. The structure of their type, especially their manner of destroying the pre-existing bone tissues, the thin osseous walls and the trabecular formation, is the standpoint for their skiagraphic study. The intra-osseous tension is responsible for the expansion of the compact osseous layer, which is thus made gradually weaker and at last almost

osteomyelitis, the cortex shows nearly normal outlines (Fig 1).

The skiagraphic expression of syphilis is also characteristic. In the congenital form, large ossified areas are recognized in the epiphyses that would appear translucent in their normal cartilaginous condition. On the other hand, light areas are noted in the diaphyses as an expression of insufficient calcareous deposition. The synostosis between the cartilaginous epiphyses and the diaphyseal end appears as a very marked line, indicating the abundance of calcareous salts deposited there. Gummata show regular light-shaded foci. Their disappearance after the administration of iodid of potassium confirms the diagnosis.

Figure 8 shows the evidence of hyperostosis of the clavicle, scapula, ribs and the upper third of the humerus in a boy of 5 years. The pelvis, femur, tibia

Fig. 7.—Destruction of lower third of radius and portion of carpus by osteosarcoma.

Figure 8.

entirely disappears. Thus we see that it is the abnormal and indefinite outline, or even the entire absence of the osseous cells, the cortex especially disappearing, which is more or less characteristic for the various types of osseous sarcoma in contradistinction to other bone diseases.

As to differentiation, it may be said that in aneurysm the bone would show intact. Attention was called by me to the usefulness of the Roentgen rays in a case of femoral aneurysm, which, on account of extremely thick walls, showed no pulsation, so that originally it had been taken for osteosarcoma, an amputation then having been considered. The femur appearing intact, it was evident that there was a disease of the soft tissues.

In chronic osteoperiostitis the walls appear irregular, but the irregularity is one-sided and—exteriorly—there is a globular or spindle-like shape (see Fig. 1). In tuberculosis the shade will be cloudy or shaggy. In

and fibula were similarly affected. The suspicion raised by the skiagraph was verified by the success of the specific therapy.

Osseous cyst showing the same clinical signs as osteosarcoma may easily be confounded with it. But in osseous cyst, while there is the same bulging as in osteosarcoma, the line of the cortex, on account of its thinness, appears narrow, but well marked and regular. The fluid center of osseous cyst renders the skiagraph translucent, the light shade showing the same regularity. The adjacent epiphyses are also normal in osseous cyst. The treatment of their various affections being difficult, the importance of a correct diagnosis is evident.¹

Osteomyelitis, necrosis, tuberculosis, syphilis and osseous cyst demand conservative measures, while sarcoma calls for the most radical treatment. On the other hand, how painful must it be for a surgeon to find that because

1. "Osseous Cyst," Amer. Jour. Med. Sciences, June, 1901.

of his error of diagnosis such radical steps have been taken unnecessarily that, in other words, an extremity may have been amputated where only an osseous cyst existed, which could have been cured by simple incision.

It is, indeed, not very difficult to confound the two diseases, osseous cysts resembling osteosarcoma in its slow painless onset often preceded by an injury; in the gradual bulging of the area involved; and in their preference for youthful age and the long bones. These being characteristic features of osteosarcoma as well as of osseous cyst, it is evident that the differential diagnosis can not be made by considering the history, nor by inspection, nor by palpation.

The fact that the interior of the osseous cyst is filled with opaque bloody serum, and its walls are lined with a smooth coat, while in osteosarcoma solid masses are formed would indicate that an exploratory incision combined with microscopic examination would clear the question of diagnosis.

But the Roentgen rays give us more valuable information than the exploratory incision itself, and for the patient a Roentgen ray exposure is certainly more agreeable than an exploratory operation. Should an operation be decided upon, the microscopic examination can as well be made then.

At the early stage, osseous cysts, be they at the tibia or at the femur, are easily overlooked, the symptoms being insignificant. Sometimes there is a very slight pain that comes and goes. The joints are freely movable; and neither inspection nor palpation reveals any abnormality. After several months have elapsed the circumference of the extremity may appear very slightly enlarged, but it may not be before a fall on the thin shells of the cortex has produced a fracture that the symptoms are fully appreciated.

Other osseous diseases like osteoma, osteomalacia, rachitis and chondroma also offer some skiagraphic peculiarities in proportion to their various textures. Osteoma, of course, shows the shape of the osseous deformity, but there is a normal architectonic structure.

On account of the dissolution of the calcareous salts, osteomalacia is, similar to rachitis, distinguished by the absence of an osseous shade. In contradistinction to osteosarcoma, the whole bone appears translucent. In chondroma there is a regular light-shaded area of lobular shape according to its cartilaginous character.

Metastatic carcinoma in bones can also be represented.

A CASE OF RELAXATION OF THE PUBIC JOINTS DURING PREGNANCY.

JOSEPH B. DE LEE, M.D.

CHICAGO.

Mrs. B., age 39, XII para, four abortions, all deliveries normal. This pregnancy has advanced to the ninth month. Beginning at the sixth month, patient has had difficulty with the limbs and pain about the pelvis. The pains began on the right side over the sacro-iliac joint and have been getting worse there, also extending to the other sacro-iliac joint and to the pubis. Now the pain extends down the right leg, and all motions of the lower trunk and extremities are painful, being referred especially to the pelvic girdle, the pubis and the right side. Difficulty in walking has increased up to the present, locomotion at the present time being laborious and distressing. She can not cross the legs and all movements of the trunk and lower extremities are carried out with difficulty and pain. In sitting down or getting up, she has to use the hands and arms, fixing the shoulder girdle and thus suspending the trunk. She can lie only on the side, and can not turn nor assume the erect posture without assistance. The pains and disability are worse

at night, worse upon beginning to move around, then better after being up a while, but get more unendurable as she becomes tired with exertion. There is no trouble with the bladder or the bowels, and no paresthesia.

Examination: General health fair; abdomen much distended, somewhat pendulous; uterus large; varicose veins in the legs and vulva, which patient says are quite painful. Patient sits on the edge of a chair with legs extended, slightly everted and abducted. There is an apparent paraplegia, but the patient can use all individual muscles; no sensory disturbances; knee jerks slightly exaggerated—as is usual in pregnancy. Abducting the thigh produces great pain at the pubis and at sides of the pelvis, as do nearly all passive movements of the legs. The patient herself says that all the trouble is "found around the pelvis." Pressure over the pubis creates outcry of pain, much less so when the rami are touched. Pressure over the center of the sacrum is not painful, but over the sacro-iliac joints is distinctly so, the demarcation being striking. Squeezing of the pelvis together from the sides is acutely painful, in fact, one thinks of an inflammatory affection of the joints.

Over the pubes a distinct separation of the ends for three-eighths of an inch can be felt, and, when the thigh is manipulated, the ends can be felt to ride on each other.

Vaginally the posterior surface of the pubis is very tender. The interval between the ends of the pubic bones is spanned by a tense elastic membrane, which can be pressed into the joint one-eighth of an inch. There appears to be no fluid in the joint, but this was hard to determine. Moderately extended movement of the thigh showed an up and down riding of the pubic ends of one-fourth of an inch. The tenderness extended down the rami pubis about two inches. There was no tumefaction of the ends of the bones. The sacro-iliac joints can not be palpated owing to the unruliness of the woman.

The history that this woman gives is interesting, but does not throw much light on the causation of the affection. Her mother died of tuberculosis; the father is living. She has had neither rheumatism nor any particular sickness. She menstruated early and regularly. She was married at 18, had a daughter within a year, another child within two years, and was six months pregnant with a third—now a boy of 17—when she "caught cold" and "had inflammation of the womb," being in bed for two weeks. Since that time she has had a persistent pain in the sacrum and the right side, which was especially marked during the numerous pregnancies that have intervened till now. Two or three weeks after delivery the patient has always regained her previous condition. At each succeeding pregnancy the pains and disability have gotten worse. They also begin earlier and the degree has been more marked until the present, when the patient is all but bedridden. She says, however, that in the last week or so there is a slight amelioration in the symptoms. A tight pelvic girdle had been used for a while without the slightest benefit. Counter-irritants and rest have helped, but at times the patient has needed narcotics.

The ancients taught that the pelvic joints always softened during pregnancy and opened during labor, so that the fetus could readily pass through. They ascribed difficult labor sometimes to the too great rigidity of the joints. Later the opposite was believed to be the normal, and those cases where there was relaxation of the joints were called morbid. Recent experiments, undertaken in behalf of the operation of symphysiotomy, show that there is some softening of the joints during pregnancy, and clinical experience confirms this.

The ancients must have reasoned from analogy, having observed the softening of the joints in cows and other animals. The "sinking of the rump" and the difficulty of locomotion of the cow show the veterinarian that the