FOUR CASES OF FRIEDREICH'S ATAXIA WITH A CRITICAL DIGEST OF RECENT LITERATURE ON THE SUBJECT.*

BY J. MACKIE WHYTE, M.A., M.D

Physician, Dundee Royal Infirmary.

SUMMARY:—*A group of three cases of Friedreich's ataxia; an isolated case; tabular comparison; remarks on some of the symptoms, ataxia of voluntary movement, static ataxia, Romberg's sign, nystagmus, choreiform jerks, athetoid movements, deformities of upper and lower extremities, and of spine, condition of sensory system, mental state, reflexes, order in which the symptoms appear; their mode of advance; period of onset; increasing complexity of diagnosis; diagnosis of isolated cases; diagnosis from chorea, from hereditary cerebellar ataxia; diagnosis as affected by indications of mental defect; diagnosis from disseminated sclerosis, from locomotor ataxia, especially in children, from cerebellar disease; pathological anatomy; theoretical views as to pathology of the disease; treatment; literature.*

THE list of family diseases of the nervous system is now fairly lengthy. Higier,⁷⁸ one of the most recent writers on the subject, makes out a list of thirteen groups, as containing the more important representatives, and some of these consist of several sub-groups. It is common to find, as Marie puts it, that each family even of fairly well-defined type such as Friedreich's disease has its own special traits; consequently the grouping of affected families is more or less arbitrary, and in many cases, as yet premature. As we shall see further on, the records of transition forms, connecting one class of disease with another, are multiplying. What is wanted is the continued careful recording of clinical cases of those rarer diseases with, if possible, exhaustive autopsies.

* A Thesis for the Degree of M.D. Edin.

Friedreich's ataxia is a well-marked clinical type; many cases belonging to it have a "family" character, many are "isolated"; only a few are, properly speaking, "hereditary." This paper first gives a clinical account of four cases of Friedreich's ataxia; then follows a discussion of a number of the more interesting points in connexion with the symptoms, diagnosis and pathology, along with a critical review of the recent literature.

A. A GROUP OF THREE CASES OF FRIEDREICH'S ATAXIA IN ONE FAMILY.

The parents of the first three cases of Friedreich's ataxia about to be described were Irish. They lived in Ireland till after the birth of their fourth child. The father, a factory worker, was a heavy drinker, and died at 50 of liver disease. The mother, likewise alcoholic, died of pneumonia. She suffered also from gastric ulcer, obstinate pruritus, and eczema of face and neck. The eldest son informs me that his father used to say he knew of no nervous affection previously in their family. No history of insanity, fits or any other nervous disease can be obtained. The circumstances seem generally to have been very poor, such as are common to thousands of Irish in Dundee and Lochee. The patients belong to a family of nine, accounted for as follows:—

- (1) First child, premature, died at two months.
- (2) HUGH—*ataxic*.
- (3) MARY—*ataxic*.
- (4) Nora, married, died, aged 25, in childbirth; showed no ataxic symptoms; had three children, eldest alive and healthy, the other two died when a few days old.
- (5) Maggie, married, subject to headache since her marriage; two premature children, dead; one alive and healthy.
- (6) John, died aged 13 of typhoid; had shown no ataxic symptoms.
- (7) Agnes, healthy, recently married.
- (8) James, healthy (aged 17).
- (9) WINIFRED, *ataxic*.

Numbers 5 and 8 show nystagmoid jerkings; their knee-jerks are normal. All were suckled by the mother. Mary shared her mother's milk with another child.

Case I.—Winifred M'A., aged 15, a mill-worker.

Patient's History.—She seems to have had good health until March, 1888, when she was sent into the Dundee Infirmary as suffering from typhoid fever. A brother, aged 13, had died at home of this disease. Her illness had begun with rigors and slight diarrhœa, and she had lain at home for three weeks before admission to the Infirmary. She was there found to be underfed, and to be suffering from slight diarrhœa, but beyond this there were no definite signs of illness, and she was soon discharged much improved.

A short time after this her father and eldest brother observed the first symptom of her present illness, namely, some difficulty in walking. Her mother also spoke of the way the girl kicked when asleep. Very gradually a change in her speech developed, so slight that it would probably have escaped notice, had not her brother Hugh been on the look-out for symptoms of the disease from which he and his sister Mary had long suffered. On the sensory side there were no complaints with the exception of some pains in the back of the thighs, occurring in the spring of 1894.

In July of that year I sent her into one of my wards in the Dundee Infirmary, and since then she has been more or less constantly under my observation. During the past two years the girl has frequently stumbled and fallen when running, and this seems to have been always in imperfect light, in the early morning or in the evening. Moreover, she says the staggering is worse when going to her work in the morning, both in summer and winter, rest having apparently a greater influence than fatigue in increasing it.

Present condition (February 21, 1897). Fairly well grown; expression of upper part of face bright and intelligent. The lower jaw usually droops and the mouth is partly open. Cheeks well coloured; nutrition rather below the average. (This is easily explained by her home conditions.) Bones seem well formed; muscular development rather feeble, but not specially so in any locality.

Nervous system.—The following conditions are to be noted on the *motor* side. When *sitting*, slight swaying movements of the upper part of the body occur; the hands can lie perfectly still. When *standing*, patient prefers to keep the feet some distance apart, and can then balance herself steadily enough if her eyes are open. She fixes them on some object in front of her. There is swaying of the trunk and head, made much worse

when the feet are brought close together. If the eyes are then shut, patient at once falls over to one or other side, or forwards. The tendons on the dorsum of the foot are seen to contract (*danse des tendons*), and the toes grasp the floor; the great toe, however, is often over-extended. Balancing on one leg is quite impossible. Her head is always bent forward a little, whether sitting, standing, or walking. Her *gait* is distinctly awkward and ungainly. The steps are of varying length, her feet are sometimes wide apart, sometimes they knock against each other; she cannot keep to a straight line. Her feet are brought to the floor with a slight stamp, through a jerky movement with the character, not of spasm, but of inco-ordination. The staggering character is more marked when walking backwards, and still better if the eyes are closed. She can turn round fairly well, and can stop quickly when told to do so.

All *voluntary movements* can be carried out in a way, but the *ataxia* which is so marked in walking is likewise manifest in other directions. For instance, when patient is told to pick up a small thread off a smooth wooden surface the hand and fingers oscillate a little, and she prefers to rest them on the surface before catching the object; there is an awkwardness about the act. When told not to touch the board, the hovering movement described by Charcot is shown. Apart from this ataxia, however, she can carry out her intentions—can button her clothes behind her back, thread a needle, etc. A full cup is slightly spilt when she carries it, but the oscillations are not increased when she puts it to her mouth. When put through other tests for inco-ordination in the upper extremities with the eyes shut, there is a decided defect. When told to touch the tip of her nose, for instance, her forefinger first reached the upper lip at its lower border. My notes show a distinct increase in the ataxia of the upper extremities since 1894. When lying down she can touch objects with her heel or toes without very marked inco-ordination.

Involuntary movements include some already mentioned—the swaying of the body, the oscillations in carrying out a voluntary movement. But there are also occasional choreiform twitchings in the face, especially drawing up the right side of the mouth, and some jerky movements of the same nature have been observed in the head and fingers. To this class, too, belong the evidently very marked jerkings that occur during sleep. Patient is quite unaware of them, but they must interfere considerably with the repose of her bed companions. They have been continuous since the onset of the illness.

Certain involuntary movements are seen in the eyes when they are strained by being directed to the extreme right or left, a slow nystagmus, about one oscillation per second. When the eyes are fixed on an object above or below the horizontal line, a very fine oscillatory movement can be detected.

When the tongue is protruded, constant slight movements are seen.

Disturbance of speech is to my ear decidedly more marked than it was in 1894. And yet it is not very bad, especially when patient is paying attention to it. Her voice is deep-toned, the words are uttered slowly, there is rarely any slurring; long words or combinations as "post office savings bank," "articulation," can be uttered with a single effort. A stranger might find little difference between her speech and that of other girls of her class, but my own knowledge and her friends' statements show some deterioration to have taken place. She can write legibly, but the writing is tremulous and uneducated; she holds the pen or pencil in a clumsy way, but one must remember that she has hardly written a line since leaving school three or four years ago. I do not think her writing has grown worse since I first knew her.

Sensory Functions.—Patient used to have a giddy feeling at times, but she is now free from that. There are no abnormal sensations whatever and the various forms of cutaneous sensibility are normal.

The muscular sense is affected; patient is a good deal astray in trying to localise the position of arm or leg when blindfolded. (In 1894 the sense of posture in the upper extremities was fairly correct.) She constantly makes mistakes in trying to distinguish between different weights with her legs, but with her arms is fairly correct. Hearing, sight, and taste are normal; deglutition, defæcation and micturition are normal. Patient has not yet menstruated.

The *knee jerks* are completely absent and have been since I saw her first. Superficial reflexes normal.

Condition of Muscles, and Deformities.—There is no sign of flabbiness and no muscular atrophy can be detected. But there is an increasing deformity about the hands. When the right hand is extended, there is a very slight flexion of the wrist, a slight hyper-extension at the metacarpo-phalangeal joints of the fingers, especially the fourth and fifth, and a gentle flexion of the inter-phalangeal joints. The little finger shows a decided bend forwards. The fingers can be well separated from one another, and the wrist moves freely in all directions when the fist is closed, but by no

effort can patient extend the phalanges completely. The thumb is not affected in any way. The same description applies to the left hand, though here the deformities are rather less marked.

This condition of the hands is very distinctly worse since 1894.

The feet show a very high instep and prominent dorsum when patient is resting, not so marked when standing. There is no talipes and no hyper-extension of the great toe beyond the occasional involuntary movement before mentioned.

A slight spinal curvature is present, convex to the left in the middle dorsal region, convex to the right in the upper dorsal and cervical regions. The bend forwards of the head is due to a slight kyphosis.

Grasp is good on both sides, and the muscular power of the legs seems to be normal. Patient is easily tired, but works for the full day, earning seven shillings a week as a shifter.

Electrical reactions are normal, both with faradic and galvanic currents.

Her psychical condition is normal ; intelligence rather above the average.

Case II.—Hugh M'A., aged 33, a mill-worker, began to walk when 18 months old. Between his third and sixth year he suffered from some illness "like water in the head," and was for two years in bed. He went to school when 6, and remained there till he was 12, learning with normal facility.

At the age of 9 he had a bad fall on the back of his head, from a cart, and was rendered nearly unconscious, but there were no bad after-effects. On leaving school he began to work in the mill, and showed some nervous symptoms. Other workers were afraid he would fall among the machinery, and some suspected him of smoking. There do not seem to have been any ataxic symptoms sufficient to attract attention before this.

He soon got into the way of holding his head forwards, with his eyes bent on the ground that he might balance himself better ; he preferred to keep his hands stretched out, the palms being directed forwards when walking.

From the age of 14 he found some difficulty in picking things up. There was not much change in his condition for the next four years, but he found that mill overseers were rather shy of engaging him in case of accidents. Between 17 and 19 he thinks he improved, but after that he became markedly worse, and from the age of 20 walking has been impossible. About that time he

had loss of sensation in his feet, and got a burn on one of his soles through standing on a hot coal. For a long time he has suffered from numbness and feelings of creeping and prickling in his feet and legs, not in the arms. Often about two hours after going to bed he has a sensation of blood rushing up to his head, and his heart seems to stop. His legs sometimes start involuntarily at night. He has had no lightning pains, no girdle pains, no vomiting, no disturbance of bowels, no affection of vision. From about the time at which he lost the power of walking he has had some difficulty in micturition.

On August 1, 1884, he was admitted to the Dundee Royal Infirmary, and though the notes then taken are brief and evidently imperfect, it is worth while reproducing them for comparison with his present state :—"Head hydrocephalic in shape, 22 inches in circumference. Dorsal spine shows right lateral scoliosis of very exaggerated character; sternum raised; left lower ribs touch crest of ilium.

"*Sensory Functions.*—Tendency to coldness in feet; formication in soles; tingling sensations in legs and feet; anæsthesia to touch and pain below level of iliac crests; pupils large, sight good, hearing good.

"*Motor Functions.*—Drop wrist well marked on right side when hand held out, muscles of fore-arm and ball of thumb flabby, diminution of grasping power, no power in lower limbs except slight power of extension of hips and knees.

"*Visceral Reflexes.*—Normal except that patient sometimes takes half-an-hour before he can micturate. No skin or tendon reflexes in lower limbs.

Slight tendency to waste in muscles of legs. Sweats a good deal."

In July, 1894, I examined him minutely, and on comparing the notes taken then with those made now, I find very little alteration.

State on Examination, February, 1897 (see figs. 1, 2 and 3).—An under-sized, fairly nourished man, face somewhat flushed, expression intelligent, features thick, this being especially seen in lips, head large, brachy-cephalic, 59.4 cm. in greatest horizontal circumference, held well forwards. His usual *attitude* is sitting in an arm-chair, with his arms resting on a table in front of him. If he has no table before him he likes to twist his arms round the arms of the chair, to keep himself from sliding off. When thus supported in a sitting position his left shoulder is depressed, and if the *spine* is looked at (see fig. 2) it is

FIG. 1.

Case II.—General helplessness indicated. Patient is supported by man behind curtain.

seen to be curved (convex aspect) to the right, between the first and twelfth dorsal vertebræ. A slight compensatory curve the other way is seen in the lumbar region. The spine also shows a marked convexity backwards, in the upper dorsal regions, and a concavity (lordosis) in the dorso-lumbar. The upper four dorsal and the lower cervical spines are nearly horizontal.

When lying as straight as possible on his back the lower end of the sternum is very prominent, 7 cm. higher than the upper end. The sternum is straight, except the xiphoid, which is bent back. Intercostal spaces are narrow. No prominences at costochondral unions. The hand can be easily passed behind the back on account of the lordosis. The ribs on the left are depressed to the iliac level.

Extremities.—(a) Upper. Muscles moderately well developed and pretty firm. No marked atrophy, though the interossei seem thinner and softer than any other set. No contractures, no fibrillary twitchings; no rigidity; passive movements possible in any direction.

(b) Lower. Muscles fairly firm, but considerably wasted, apparently from disuse. Right calf 25.6 cm., left 27 cm. in circumference; legs lie helplessly in bed, generally flexed at knee; foot shows a permanent deformity, the shape closely resembling that of *pes cavus*, arch extremely well marked on both sides, less on right than left; dorsi-flexion of first phalanges of toes, clawing of other phalanges to some extent. The deformity is less marked when the feet rest on the floor. On first exposing the right foot, a fibrillary tremor was noticed in the muscle on the inner border close to the arch (short flexor of great toe). Limbs very cold, and red or blue below the knees.

Motor Functions.—When in bed the patient prefers to lie on his right side, curled up. Cannot lie on left side longer than half an hour at a time. Has the greatest difficulty in raising himself, or turning round. In trying this he makes enormous efforts, painful to witness, and becomes very much flushed. The voluntary movements of the trunk are thus very jerky; he tosses himself round from one position to another.

Respiratory movements require a distinct effort; when sitting quietly with his table before him, both inspiration and expiration are audible through the room, a distinct pause separating them. The breathing is nasal; with each inspiration the head is jerked up a little. He is said to breathe more heavily and to snore at night. Power in the muscles of the neck seems well preserved; head can be moved in any direction. Occasionally there is a

slight choreiform vibration of the head, especially on making a voluntary movement. Swaying movements of the head and body are constant when sitting. No choreic twitchings of face, or other disturbance of facial movements.

Voluntary movements in the *arms* moderately active, but not so powerful as one would expect from the appearance of the muscles. Grasping power much diminished; free movement at shoulders, elbows and wrist. When told to straighten the fingers, the hand resembles the *main en griffe* of progressive muscular atrophy (see fig. 3). There is flexion at the wrist, over extension at the metacarpo-phalangeal joints, and flexion at the first inter-phalangeal joints. There is dorsi-flexion of the second phalanx of the thumb, and also of the ungual phalanx of the right mid-finger. (This the result of an accident.) The index is less affected than the other fingers, and the condition is more exaggerated on the right side. In extension the fingers are kept slightly apart, but they cannot voluntarily be completely adducted or abducted. The wrist and fingers can be quite easily straightened by an external force. The extensor tendons can be traced up the lower third of the fore-arm, and on the front the tendons of the palmaris longus and other flexors stand out. The palm is hollowed.

Movements of fingers and arms, in trying to pick up or touch objects, are badly co-ordinated. Patient likes to have the arms supported on the table, etc.; when this is done there is not much oscillation. Sense of posture in upper limbs defective.

As for the *lower limbs*, walking is quite impossible. Slight power of flexion and extension at hip. None at knee or ankle, very little at toes.

Sensory Functions.—In the legs there is slight diminution of acuteness in sensibility to touch, but pain is felt normally. No lightning pains; some slight paræsthesia in lower limbs is noted in the history.

Reflexes.—No knee or Achilles jerks, plantar reflex well marked on right, slight on left. Cremasteric and abdominal reflexes marked. Visceral reflexes normal apart from delay in starting micturition.

Sight.—Normal; pupils act readily to light and accommodation; a moderate degree of nystagmus, consisting of irregular jerkings on fixing eyes on an object, to the side.

Hearing.—Slightly dull (this also the case in 1894). Hears watch at 12 in. on right side, at 3 in. on left. Occasional sound of rushing water in his head.

Tongue.—Flabby, marked by teeth, slightly furred on dorsum; when protruded a constant fine tremor pervades its whole extent.

FIG. 2.

Case II.—To show Scoliosis and Kyphosis.

FIG. 3.

Case II.—To show *main en griffe*.

Speech is markedly disturbed. One requires to attend closely to make out all he says, though if his attention is directed to words, he speaks much more clearly. The voice is monotonous, the words are somewhat slurred, not scanned. "Remember" becomes "rememr," "Whether" becomes "wher." "British constitution" becomes "Brish constution." "A dainty dish" becomes "a den dish."

Psychical state is normal. Patient is said to be rather irritable, and to make what the others consider undue demands from his being the eldest of the family. His intelligence is quite unaffected.

Electrical reactions with galvanic current. Extensors of left forearm KCC 7, ACC 12, KOC 8, AOC 9. Flexors of left forearm, KCC 9, ACC 9, KOC 12, AOC 13.

Heart—Sounds normal; position altered, apex being in third left space external to nipple.

Case III.—Mary M'A., aged 31, formerly millworker (figs. 4 and 5). *History*: Patient had inflammation of the bowels, measles and whooping cough in infancy, but there was nothing indicating any ataxia or paralysis until she was fifteen. At this age, in the month of July, she was taken ill one morning when on the way to work, feeling cold and suffering from bleeding at the nose and mouth. She became unconscious that night, and remained so for about six weeks. During the next eighteen months her mind was enfeebled, and she could not walk steadily; she managed, however, to do some work in the mill, and in course of time her mind recovered its former clearness. The staggering, however, increased to such an extent that at the age of twenty-one she was refused further employment. From the beginning of her illness she was subject to cramps and to jerkings in the legs at night.

At the age of nineteen, and again when she was twenty-one, she was a patient in the Dundee Royal Infirmary on account of spinal curvature.

Until she was twenty-six she was able to get about on her legs, pushing a chair in front of her, but during the last few years she has been unable to walk at all. She says that her hands have both been weak—the right more than the left—ever since the acute illness which marks the beginning of her disease. Still she could sew, although her hands were shaky; she has never been able to read or write, her school education having been very imperfect and short in duration. Since July, 1894, she had been almost constantly under my observation, in the Dundee Infirmary and the Parochial Hospital.

Present Condition.—A well-nourished, fresh-coloured woman, with cheerful expression, ruddy complexion and numerous freckles on face and neck. Her hair is light brown, strong and abundant; her nose is twisted from a fall four years ago, and on her right cheek is a scar, due to another accident. Through this scar there is a puckering of the cheek below the inner part of the right eye, when the lids are firmly closed; there is also a slight drawing-up of the right angle of the mouth.

The head is well-shaped, and measures 55.4 cm. in the greatest horizontal circumference; it is kept bent forwards through the spinal curvature, but patient says she was accustomed to hold it so because she was ashamed of her staggering.

Her usual attitude is sitting up in bed, with the back well supported by pillows; the spine shows a marked dorsal curvature, with convexity to the left, extending from the sixth dorsal vertebra to the second lumbar, the deviation at its maximum amounts to $1\frac{1}{2}$ inches; the ribs project backwards to a marked degree on the left side, and are well separated from one another. On the right they are crushed together, and the twelfth rib is on a level with the iliac crest. In the upper dorsal region there is a slight curve with convexity to the right.

Some kyphosis is present in the upper dorsal spine; this can be removed partially, and for a very short time, by an effort of the patient.

The dorsal curvature is somewhat less pronounced when she is raised up by the arm-pits.

She complains a little of soreness, on pressure over the spine in scapular region.

The mammae are well developed, and there is no striking asymmetry when the chest is examined from the front.

Extremities.—Upper extremities show, when at rest, no deformity; the bones are well shaped, the muscles seem smaller than normal on the back of the fore-arm. The lower limbs, on the other hand, show a deformity about the foot, talipes equino-varus being present on both sides. The legs below the knees are more cylindrical than usual, from a want of development of the calf; the antero-lateral group of muscles is deficient in bulk; the heel is drawn up, some transverse wrinkles of the skin being present over the lower end of the tendo-Achillis. The hollow of the pedal arch is very much greater than usual, and this is so even when patient is made to stand. Most of the grooves on the sole run from within, outwards and forwards. The great toe, especially the right, is always over-extended. It can be brought voluntarily

FIG. 4.

Case III.—To show *main en griffe* and deformity of lower limbs.

into line, but not flexed. The other toes show slight over-extension at the metacarpo-phalangeal joints; at the inter-phalangeal joints there is flexion. The dorsum of the foot is prominent.

Motor Functions.—Patient shows distinct weakness in the muscles that raise the *trunk* into a sitting posture. She can raise herself, but awkwardly and slowly. When sitting, there are constant swaying movements of the head and body; there are, besides, frequent jerkings of the head, and also choreic twitchings of the face, especially in the right levator anguli oris. These are not by any means so marked as to cause any confusion with true chorea.

In the *upper limbs*, movements are free at the shoulder, elbow, and wrist joints (with hand closed). When told to extend the finger and hand there is a marked imperfection; the wrist is somewhat flexed forwards, and to the ulnar side, the fingers are over-extended at the metacarpo-phalangeal joints (especially the ring and the little fingers), and flexed at the first inter-phalangeal joints. The terminal phalanges are in line with the middle. The deformity is, in short, the *main en griffe*. The thumbs are unaffected; the fingers can be separated, though less freely than normal.

Ataxia is present in the upper limbs, but patient can button her clothes, or thread a needle, if the elbows are supported. Her movement is like her sister's (Case I.) in picking up small objects, and in carrying out any movement oscillatory jerkings are very noticeable. The ataxia is brought out by the usual tests of bringing the fore-fingers together, or touching the point of her nose, with eyes shut. Hesitation and waving movement are more marked when she is near the point aimed at. The super-added movements are less marked if the eyes are open.

Power of grasping is still fair, though awkward from the extensor wasting. At elbow and shoulder muscular power is well preserved.

Lower Limbs.—Patient can flex and extend at hips and knees, but with lessened power. With her legs extended she can raise the right foot two inches off the bed, the left twelve. Even when supported on each side she cannot walk; her feet are pushed too far forward, so that she is unable to get her balance. Her steps are very short, the right foot is crossed over the dorsum of the left, and thrown two or three inches beyond the place where it ultimately touches the floor. She puts down her feet with a slight stamp, first the toes and then the heels. There is only

very slight voluntary movement at the ankle, and that is mainly extension. The toes allow of somewhat further voluntary movement. The ankle can be passively moved to a much greater extent, but dorsi-flexion of the foot is much interfered with, through contracture of the tendo-Achillis. The heel can be placed on the opposite knee, but slowly.

Sensory Functions.—Occasional headaches, no giddiness. There is some slight delay in conduction of impressions from the lower extremities, but sensibility in its different modes is essentially normal. There is a small area on the front part of the dorsum of the left foot, which shows slight dulling of sensibility to touch, not to pain.

Reflexes.—No elbow or knee jerk, abdominal reflexes well marked, plantar exaggerated, no disturbance of visceral reflexes.

Eyes.—Reaction of pupils normal; when looking to either side some nystagmus-like jerkings occur—about seven in five seconds, irregular in time. They are also seen when she looks up or down; sight is normal.

Hearing.—The hearing distance varies, but is usually about three inches on each side (*watch test*).

The Tongue, when protruded, is constantly in a state of movement—not the irregular jerks of chorea, but a fine oscillatory movement.

Speech.—Her voice is low-pitched, not harsh, monotonous. She keeps her teeth rather close in speaking. Some words are slurred—"doer" for "doctor," "I don't emember" for "remember." The speech is consequently somewhat slipshod; it is very like her brother Hugh's.

Vasomotor System.—Feet tend to be cold; psychical condition normal; muscular sense normal.

Electrical reactions. *Faradic current* much better borne on legs than normally; somewhat better on arms. Muscles of arms contract well under its influence, but the interossei act sluggishly.

Galvanic current.—In flexors of right forearm KCC 6, ACC 7; in extensors of right forearm KCC 6, ACC 7, KOC 8, AOC 8; in antero-lateral muscles of right leg KCC 6, ACC 10. The interossei, especially those of the third and fourth spaces, respond feebly to galvanism.

B. AN "ISOLATED" CASE OF FRIEDREICH'S ATAXIA.

Case IV.—William M., aged 13, admitted to Dundee Royal Infirmary, August 30, 1894.

Family History, &c.—Patient is third of a family of four, the

FIG. 5.

Case III.—Scoliosis. Patient requires to be supported.

youngest three by the same mother. His father died some years ago of inflammation of the lungs, after eight days' illness. His mother, sisters, younger brother, and nephew are all healthy. A slight nystagmus can be detected on extreme lateral deviation of eyeballs in his brother and younger sister. There is no history of any nervous affection in any of the other members of the family, nor is there any evidence of syphilitic or tubercular taint. His home surroundings are good; he has not been overworked at school, and has shown average intelligence.

Present Illness.—About two years ago his mother noticed that he became easily fatigued, and did not care to join in the games of his companions. His gait was a little unsteady; if he ran or walked quickly he was apt to stumble or fall. This was no worse in the dark, but if his feet were bare there was decided aggravation. There was no definite onset, no pain, shivering or feverish attack. His condition did not change much for some months, and then his upper extremities were seen to be slightly affected. He could not dress himself so quickly as usual, his hands were a little unsteady, and he was apt to spill the contents of a saucer in drinking. His handwriting was rather shaky, there was a want of freedom in using his pen or pencil and a difficulty in forming some letters. Both sides of his body seemed equally affected. The only time he complained of pain was a year ago, when his back ached at times, but this feeling passed off after the back was painted with iodine. He had no headache, or shooting pains in his limbs. During the past few months his arms and his legs have been getting thinner. No change has been noticed in his speech. There has been a steady deterioration from the beginning of the illness, the difficulty in locomotion increasing especially in trying to go downstairs.

State on Admission.—A fairly well-developed boy, with florid complexion, sad or pensive expression; brow transversely wrinkled, eyebrows often involuntarily drawn up while under observation. Weight 5 stone.

Nervous System (the only system affected).—The boy is quite intelligent, and has a keen sense of his unfortunate condition.

Motor Functions.—Patient stands with his feet about twelve inches apart, but if his heels are brought together he sways from side to side, and would fall if unsupported. Head and upper part of body bent forward. Closure of eyes increases the unsteadiness; when sitting in bed he has difficulty in balancing himself. Various muscles are seen to be constantly working as he tries to steady himself when sitting or standing. When laid on his back

he can raise himself up fairly easily by means of the trunk muscles alone. When standing he can stoop forwards, and pick any article off the floor.

Gait.—He walks with head bent forwards, and eyes fixed on the ground, his legs widely straddling. He brings his feet down heavily with a slap, first the toes, then the heels, reeling from side to side as he goes, and swaying his arms irregularly. His eyes are a great help to him in walking. He walks backwards with great difficulty; it is impossible to get him to keep a straight line. His gait has something of a festinant character; his steps are irregular in length, force, and direction.

Co-ordination, as tested by making him bring the tips of his fingers together, or touch other parts of his body, or place the heel of one foot on the opposite knee, is fairly good.

Writing is shaky; he presses his hands heavily on the paper. Nutrition of muscles, good; no hypertrophy or atrophy; no true fibrillar contractions, though muscular twitchings are noticed here and there even when he is lying at rest. No deformities about feet, but slight lateral curvature of spine, dorsal convexity and lumbar concavity to left.

Speech is low-toned and somewhat slow, not jerky nor scanning nor monotonous. The muscles of his lips seem slightly weak in their action, but there is no paralysis of the facial muscles.

There is distinct nystagmus (better marked on some days than others), brought out when the eyes are fixed on an object away from central line of vision.

Reflexes.—Patellar reflexes gone; no ankle clonus; plantar reflexes present on both sides, rather better marked on left; epigastric and cremasteric reflexes present; visceral reflexes normal.

Sensory Functions.—Has no abnormal sensations such as cold, heat, numbness or prickling. Sensibility to touch seems a little dull in lower limbs, the light touch of the head of a pin being frequently unfelt. Sensibility to pain acute, can readily localise a pin prick, can easily distinguish heat and cold.

Sight good; pupils equal, regular, reacting freely to light, but somewhat slightly to accommodation.

December 13, 1894.—Tongue thick and flabby, constant tremors in it, especially on protrusion. Frequent involuntary jerkings occur in head and upper limbs. Complains of feeling weaker, and often lies in bed for a day. Has occasionally been troubled with cramp in muscles of calf. His walking and general power of balancing are much worse, and he often falls. In trying to touch

an object he is more uncertain in his aim, from the irregular jerkings in his arms during voluntary movement. There is a corresponding difficulty in carrying out the heel-on-knee manoeuvre.

Fundi of eyes show nothing abnormal. (Dr. Macgillivray.)

February 19, 1897. — Since last note patient has had fair general health. He has grown much taller, but there is no improvement as regards any of the symptoms described above. His speech is rather worse, the lips still seem somewhat at fault. The voice is more monotonous, and is low in pitch, distinct enough when speaking to me, but his mother says that when he is reading aloud to her, as he is fond of doing, she often has difficulty in understanding him. There is a thickness, a want of crispness about his utterance.

The ataxia of the hands is still as marked, if not more so. He can put in buttons, and write his name—always with the same awkward procedure. The superadded movements in using his hands are much as before.

He can walk, but only with strong support. His soles feel the ground quite well: the feet are raised higher than normal and brought down with a stamp, parietic not spastic. Romberg's sign is present.

When in bed he lies on his back; he has often a painful cramp, which causes the left leg to draw up of itself. The pain is immediately above and below the knee. The limb has first to be rubbed over the knee and then pushed down. If it were simply pulled down, the pain would be severe.

In the morning he has some pain in the lumbar region, when bending to get on his clothes with his mother's help. This is probably due to some stiffness, developed through his position in sleep. His back is very weak, and shows still a slight scoliosis; this is not of a permanent character.

His favourite position is sitting on the floor, with his back against the wall.

A recent symptom is a difficulty in commencing the act of micturition; there is no dribbling afterwards.

A brief *résumé* of these cases will be useful, and will show the main diagnostic points in Friedreich's ataxia.*

* (I take this opportunity of acknowledging the valuable aid given me in taking Cases I. and IV., by Dr. C. Kerr, then house physician in Dundee Infirmary, and by Dr. A. Croll, resident physician in the Parochial Hospital, in recording Case III.

REMARKS ON SOME OF THE IMPORTANT SYMPTOMS.

Disturbances of the motor system are by far the most important in the symptomatology of Friedreich's ataxia. So much is implied in the name, and Friedreich himself pointed out that the ataxia appears in two forms—ataxia or voluntary movement, and static ataxia. Choreiform jerks, nystagmus, tremors of tongue, and possibly disturbances of speech, are further evidence of wide-reaching motor invasion, while in the later stages of the disease certain muscles are apt to become paretic or paralysed.

Ataxia of voluntary movement is well shown in the gait. It has been often described, and has a character which marks it off from that of tabes dorsalis. In two of my cases (Cases I. and IV.) alone could it be examined. The patient in walking keeps his legs apart, and has difficulty in keeping to a line. He reels a little, sometimes a good deal, like a drunken person. The feet are brought down with a slight slap, first the toes and then the heels. The steps are short and vary in their length. The arms sway to some extent to aid in balancing, and for the same reason the eyes are fixed on the ground, a little in front; the head is bent forwards. In tabes the movements of the limbs are more disorderly, there is more excess of range, more exaggeration; the foot is raised higher, and brought down with more noise. The patient can keep to a line, if not asked to put one foot directly in front of the other. The character of the gait in each case varies, however, according to the stage of the disease, and, in tabes, according to the muscles worst affected. The point about the gait of Friedreich's ataxics which is most important from a diagnostic point of view, is its cerebellar character. This occurs also in Marie's disease, in multiple sclerosis ("rarely," Leyden), and of course in cerebellar disease.

But, added to the cerebellar or tabeto-cerebellar character, are various movements of the limbs and trunk best studied when the patient stands. These movements come under Friedreich's designation of *static ataxia*.* They are perhaps

* In one of Senator's cases (34) examined when the disease had made only very slight progress, the locomotor ataxia was very slightly marked, while the static ataxia was pronounced.

CASE I.	CASE II.	CASE III.	CASE IV.
<p>(1) Family character.</p> <p>(2) Onset soon after a fever, possibly typhoid, when 7 years old.</p> <p>(3) First symptom, difficulty in walking.</p> <p>(4) Very slow advance of ataxia; cerebellar character, slightly marked; worse in the dark (Romberg's symptom).</p> <p>(5) Slight change in speech, beginning somewhat early.</p> <p>(6) Slowly increasing ataxia of upper extremities.</p> <p>(7) Swaying of body; choreiform jerks.</p> <p>(8) Nystagmus.</p> <p>(9) Absence of knee-jerks.</p> <p>(10) Slight weakness of interosseal in hands, slight scoliosis. No talipes, or hyperextension of great toe, but unusually high instep.</p> <p>(11) No marked paresis unless in hands.</p> <p>(12) Intelligence and sensory functions (general and special) normal. Perhaps some impairment of muscular sense.</p> <p>(13) Visceral functions normal. Pupil reflexes normal.</p>	<p>(1) Family character.</p> <p>(2) Severe cerebral affection in infancy. Onset of present illness at 12.</p> <p>(3) First symptom, unsteadiness in walking.</p> <p>(4) Advance of ataxia, with remission, between 17 and 19; again rapid advance, with paralysis of legs at 20.</p> <p>(5) Marked articulatory defect.</p> <p>(6) Ataxia of hands at 13.</p> <p>(7) Choreiform jerks and swaying.</p> <p>(8) Nystagmus.</p> <p>(9) Absence of knee-jerks.</p> <p>(10) <i>Main en griffe</i> on extension of fingers. Kypho-scoliosis, very marked. Talipes equino-varus.</p> <p>(11) Great loss of power in arms, trunk, and especially in legs.</p> <p>(12) Intelligence normal. General and special sensory functions normal. Muscular sense affected.</p> <p>(13) Delay in commencing act of micturition since age of 20; visceral reflexes otherwise normal. Pupil reflexes normal.</p>	<p>(1) Family character.</p> <p>(2) Acute illness (? meningitis) at 15, immediately followed by symptoms.</p> <p>(3) First symptom doubtful.</p> <p>(4) Ataxia of gait, gradually increasing. Paraplegia nearly complete at 26.</p> <p>(5) Articulatory defect.</p> <p>(6) Ataxia of hands since the age of 15.</p> <p>(7) Choreiform jerks and swaying.</p> <p>(8) Nystagmus.</p> <p>(9) Absence of knee-jerks.</p> <p>(10) <i>Main en griffe</i>. Kypho-scoliosis. Talipes equino-varus.</p> <p>(11) Parietic condition in back and legs.</p> <p>(12) Intelligence normal. Sensory functions normal. Muscular sense normal.</p> <p>(13) Visceral functions and micturition normal. Pupil reflexes unaffected.</p>	<p>(1) No family character.</p> <p>(2) Onset, insidious at 11.</p> <p>(3) First symptom, affection of gait.</p> <p>(4) Ataxia of legs (tabeto-cerebellar), going on to complete inability to walk alone. Static ataxia. Romberg's symptom.</p> <p>(5) Speech only slightly affected and that recently.</p> <p>(6) Ataxia of hands and fingers, noticed after some months' affection of legs.</p> <p>(7) Constant swaying of body. Choreiform jerks.</p> <p>(8) Nystagmus.</p> <p>(9) Absence of knee-jerks.</p> <p>(10) No deformity of hands or feet. No permanent scoliosis.</p> <p>(11) Weakness of muscles, especially in back and legs; no atrophies.</p> <p>(12) Intelligence normal. Sensory functions normal.</p> <p>(13) Visceral reflexes normal, till quite recently, when a difficulty in beginning the act of micturition came on (as in case II.). Pupil reflexes unaffected.</p>

most strikingly shown when one looks through the camera at the patient. The inverted image is seen to be in a constant state of unrest, from swaying movements which are in inverse proportion to the amount of support afforded to the body. When patient lies in bed they are absent; when sitting with the back well supported by pillows, the head and neck move; if sitting unsupported, the movements are more marked, as the trunk is now swaying; and when standing unsupported, they may be seen at their maximum. A rough examination, then, shows the patient to be standing with legs apart, head and upper part of body bent forwards, tendons about ankle and on dorsum of feet in a state of perpetual motion, toes often with an athetoid movement, grasping the floor.

A more careful examination may show the various muscles of thighs, hips, and back in a state of irregular contraction, in the endeavour to preserve equilibrium.

Ataxia of movement is shown in the lower limbs when patient is lying by such common proceedings as putting one heel on the other knee. This test is carried out by Cases I. and II. very well, much better than one would expect.

Ataxia of upper limbs is easily brought out by writing, threading a needle, putting in buttons, picking up a small object.

The super-added oscillatory movements are present in carrying out any such attempt, and the movements are clumsy and awkward. The patient when lying seems to have more control over the test movements of the feet than of the hands, possibly because those of the latter are more delicate. Senator²⁵ notes that in his first case, incoordination is less marked in the lower extremities when lying or walking, than in the upper.

Case I. illustrates the bad effect of rest, for she is always worse as regards the ataxia in the morning, quite apart from the degree of light.

Tresidder²⁶ notes the same to a much greater degree in his Case I. He says "When 23 years of age, the unsteadiness in walking was so pronounced that a London physician was consulted, who thought that the lateral curvature of the

spine was producing paralysis, and the patient was ordered to lie down continuously. She did so for thirteen months. After this she began to walk again, and was at first very weak, but in the course of some months regained her strength until she could walk nearly as well as before the thirteen months rest."

Besold's second case,⁵⁴ a hawker, believed that his long walks (20-25 kilometers) were of great service to him.

These facts are of interest and importance, in view of the excellent results in the mechanical and gymnastic treatment of tabes attained by Leyden, Frenkel and others.

Organic changes are not affected, but the function of the muscles is improved by this method of "compensation-therapeutics." (Leyden and Goldscheider, *Erkr. d. Rückenm.*, page 589.)

The test of making the patient close the eyes when standing or walking (Romberg's sign), while of very great value in locomotor ataxia, is not so useful in Friedreich's disease. Increase of unsteadiness through this is often absent, as in three out of four of Friedreich's own cases, in Rosenbaum's⁸³ elder case, in Bramwell's¹² first case. In my own cases that can walk (I., IV.) there is no doubt about the existence of this sign. In Senator's first case it was better marked on the second examination than on the first, a year and a half intervening between the two.

Where the patient can no longer walk, as in Case III., the same test can be applied to the upper extremities; here there is increase in the superadded movements when the patient is using her arms, with eyes shut.

The so-called *nystagmus* of Friedreich's disease seems best classed with static ataxia. It is not a true rhythmical nystagmus, in the sense of the ophthalmologist, though this occurs in rare cases (noted in three of those analysed by Griffith). The symptom must be carefully looked for, and brought out by directing the patient's gaze to one or other side. When the muscles of the eye-balls have their tension thus increased, distinct jerkings are seen—slow they may be and irregular, as in Case III., when the rate was seven in five seconds. Sometimes the movements can only be elicited

by turning the patients three or four times round, as in Geigel's²⁸ two cases (brothers, ages 20 and 18). On turning the patients to the right the eyeballs both turned to the right, and the short jerking movements of nystagmus to the left followed, lasting a quarter of a minute.

In two of Mendel's¹⁰ cases, the movements thus brought out were very marked, and lasted some minutes.

They are found in most cases of Friedreich's disease—"not in all" Gowers), but they are likewise often found in persons who have weak health, and even in the perfectly healthy.

This nystagmus I found in two quite healthy members in each of the families to which my cases belong, but no pathognomonic stress is to be attached to this.

On Schultze's instigation, Ausschlag⁶³ examined 200 unselected cases, and of these only 25 showed no tremor, though frequently there was only a single movement too far. "Only a strikingly high degree and special frequency of movements can be regarded as pathological."

Even modified nystagmus ("nystagmoid jerkings) is very uncommon in locomotor ataxia, so this sign may in certain cases be a help in diagnosis.

The involuntary movements, like an irregular tremor shown in the protruded tongue, may, without doubt, be referred to the same category as those we have been considering, and some would similarly treat the speech disturbances.

Besides the swaying movements of static ataxia, there are frequent *involuntary jerkings, choreiform in character*, but not so constant or violent as in chorea.

Thus in Case I., occasional twitchings are seen in the face, especially drawing up the right angle of the mouth. Very much the same facial movement is seen in her sister.

In Case IV., the brow is often raised involuntarily. The transverse wrinkles so marked in his brow may be associated with the bent position of his head, and may have been produced by his raising his eyebrows along with his upper eyelids. A wagging or wobbling of the head is constantly to be seen in the two worst cases, and to a slighter degree in the others. Even when lying at rest

occasional muscular twitchings, not fibrillar, are seen in Case IV. He is subject to a cramp in the left leg at night.

The other three cases have all been subject to jerkings in the legs at night. In the case of the eldest (Case II.), they seem to be more of the nature of cramps now, and another member of the family has often to get up to pull his legs straight.

The choreiform jerkings are an almost invariable part of the clinical picture of Friedreich's ataxia, and are of great diagnostic importance. They have sometimes led to a false diagnosis of chorea, as will be noted later on (*v. Diagnosis*).

In one case (Zabludowski's⁸⁴) are noted "choreiform movements, especially when going to sleep," and also at that time, twitching in face and ears, and flexion of thighs on abdomen.

Painful cramp like contractions when in bed have been noted by others (Bramwell,¹⁸ *Atlas*, vol. i., p. 46; Besold⁵⁴ Case II.). Nocturnal restlessness is noted in Bramwell's¹⁹ Case I. Some of these exaggerated movements at night are probably analogous to the jerks which most people experience on dropping off to sleep.

Athetoid movements occur, too, sometimes in a quite minor form, as in one of Griffith's cases where was no tremor but a slow ataxic movement, resembling athetosis, which appeared when the hands were lying passively in the lap.

Athetoid movements of the toes in walking are occasionally present (Ladame's case,¹ p. 477). In both of Wallace Anderson's³⁶ cases, athetosis was distinctly marked in fingers and toes. These cases were peculiar in other ways; the eldest was twelve years old, but looked six or seven, and his mental condition corresponded to this age. The knee-jerks were present, though faint in both, and so was rigidity of legs with spastic gait. The onset in both was after measles two years previously.

The most remarkable movements of this class were observed in three cases reported from France. The first

(Chauffard³⁸) was a boy of eight; symptoms since age of three; no heredity, sisters unaffected. No nystagmus, gait neither cerebellar nor tabetic. One leg made to pass before the other with a jerky and exaggerated convergent movement. Among usual symptoms were the progress, absence of jerks, absence of ocular symptoms, comparative preservation of muscular force, slight equino-varus. Further there was marked lordosis (dorso-lumbar), with slight cervico-dorsal kypho-scoliosis. No choreiform instability in repose, but sudden involuntary flexions of lower extremities at hips, legs remaining extended. Intelligence good, no grimaces. "Hovering" inco-ordination in picking things up. Arms took up athetoid position in walking with support. Forearms in position of pronation, hands flexed, their backs being directed down, fingers extended or flexed in the palm. The movements were not the usual slow convulsive movements of double athetosis. They only appeared when other voluntary movements were made—a muscular synergy, an athetoid synkinesia.

Chauffard easily separates this case not only from athetosis, but from tabes, insular sclerosis, cerebral diplegias, and chorea. But though there are several points of contact with Friedreich's ataxia, it is difficult to get over the uncommon features. The other two cases, reported by Londe and Lagrange,⁷⁹ are more like the ordinary type: Two sisters, aged 15 and 16, both took ill at six or seven. At first they were regarded as chorea. Absence of jerks, hollow-foot, marked inco-ordination, no nystagmus, internal strabismus in both; gait staggering; involuntary facial movement; head continually moving in various directions; choreiform jerks in head and limbs; *danse des tendons* in standing; slight scoliosis, no sensory or sphincter troubles, no affection of pupils or fundi; mental dulness. Athetoid attitude exactly as in Chauffard's case.

It is curious that in no one of these three cases was nystagmus present; in the latter two the internal strabismus, the mental dulness, and the athetoid attitude confer a special family mark.

Deformities of extremities and spine.—Certain deformi-

ties affecting the feet and spine have been always regarded as characteristic of Friedreich's ataxia. To these must be added the affection of the hand shown in my first three cases. None of these conditions is essential; the talipes (equinus or equino-varus) was noted by Griffith in only 27 cases out of 143. In a number of instances besides these there was dorsal flexion of all or some of the toes. The latter condition in itself might, as Besold suggests, be attributable to "modernes Schuhwerk," unless uncommonly well-marked. In Rüttimeyer's families, Blattner and Kern, early prominence of the extensor propr. hallucis was regarded as a sign of very bad augury—it was one of the first symptoms (Ladame). Baskett's case,⁷⁸ a girl of 13½, was treated at a hospital for double talipes varus at the age of 8½. Her symptoms began when she was six, but usually deformity of the foot is of late occurrence. It is well-marked in Cases II. and III., but not in the other two. In Case I., however, there are indications in the high instep and occasional dorsi-flexion of the great toe that the shape of her foot will become like her brother's and sister's. The general appearance in a marked case corresponds closely to *pes cavus*, "hollow claw foot." The claw position may or may not be assumed by the terminal phalanx of the great toe. In Case II. (photo) the right great toe is over-extended as regards both phalanges. This is so, too, in Case III. In Plate X. in *Med. Annual* for 1895, representing a case under the care of Mackay, of Devizes, the great toes, like the others, have the second phalanges flexed, and this seems to have been the way in Ladame's patient.

Dreschfeld⁵⁰ describes a case which showed no talipes, but hyper-extension of the great toes "resembling that seen in the early stages of peripheral neuritis."

Deformities of the hands have been rarely observed. Bramwell²³ gives a family of three showing *main en griffe*. Griffith (*Amer. Journ. of Med. Sciences*, 1888, p. 377) describes one of his own cases as having "grasps strong but claw-like," another as having claw-like hands; but he may be simply referring to the mode of prehension so common in Friedreich's ataxia. In two of Rook's⁷ cases, the hand

was claw-like and one had wrist-drop. One of Taylor's⁵⁵ cases had a little difficulty in extending the fingers.

In Cases I., II., and III., the attempt to extend fingers in line with the forearm at once brings out a more or less pronounced *main en griffe* (see fig. 3). Let me recapitulate the description of the condition in this case. When the hand is lying passively, nothing wrong is noticed. When he is told to straighten his fingers, flexion is produced at the wrist, with some bending towards ulnar side. The near phalanges are over-extended in the four inner digits, more so in the fifth than the fourth, in the fourth than the third, and so on. The middle phalanges are flexed, and the terminals are nearly in line with these. In regard to these joints, too, the deformity is more marked in the inner than in the outer digits. The fingers are separated a little from one another. The thumb is simply bent back. The more strongly patient tries to straighten the fingers, the greater does the alteration in form become. The extensor com. dig. tendons can be seen standing out on the back of the hand, and can be traced up the lower third of the forearm. The palmaris longus tendon, too, stands out prominently. The fingers can be adducted and abducted, but to a much less degree than normal.

What is the mechanism of this condition? It has nothing to do with the proper flexors and extensors of the wrist, for when the hand is closed, movement may occur at the wrist in the normal four directions—flexion, extension, ab- and adduction. It is not a paresis of the extensor com. digitorum; its tendons are inserted mainly into the near ends of the near phalanges, and Duchenne hence says they might be called *extensors of the near phalanges*. In the above case, so far from there being any sign of their failure to act, the chief trouble would appear to be their over-action (over-extension of proximal phalanges).

The cause is, I think, a paresis of the interossei—not paralysis—though the condition seems to come nearer this the older the patient becomes (compare condition in the three first cases). The interossei flex the first phalanges, and extend the second and third; they also abduct and

adduct the fingers. When they are weakened, and when the common extensor is thrown into action, the fingers assume the position above described. To keep the first phalanx in line with the metacarpals, antagonistic action of its extensor and flexor is needed. "When the interossei fail, the unopposed extensor causes an exaggerated extension of the near phalanges, and the extension of the middle and far phalanges does not only not occur, but they are, on the contrary, flexed in direct proportion to the degree of extension of the near phalanges." (Duchenne, *Syd. Soc. Trans.*, p. 265.)

The flexion at the wrist may be thus explained: when the hand is extended, not only do the proper muscles of the wrist act together, but also the long extensors and flexors of the fingers. The combined effect of their synergic action is to place the metacarp in line with the forearm. If the interossei are weakened, and the near phalanges are over-extended, the balance is upset, the common extensor acts at a disadvantage as regards its extending action at the wrist, and there is the appearance of a slight drop-wrist. This can be imitated to some degree in a healthy limb.

The greater amount of distortion in the ring and ulnar fingers corresponds to the condition in injury of the ulnar nerve, and is accounted for in this case by the escape of the first two lumbricals, which are supplied by a branch of the median. That the whole terminal part of the ulnar is not affected is proved by the intact sensibility, among other points.

The simpler case here considered leads to the consideration of the "Friedreich foot," about which a good deal has been written. I believe that it is analogous to the condition just described, and that its formation can be referred to an affection of the interossei of the foot (including the short flexor and adductor of the great toe). There may also be a real weakening of the antero-lateral muscles of the leg; there is certainly some retraction of the calf muscles, and tightening of the tendon-Achillis.

Paretic condition of various muscles besides the interossei is present in advanced cases. This is shown in the help-

lessness of Cases II. and III., especially as regards the whole of the lower extremities. The muscles that act on the spine may also be involved. In early cases, decided loss of power is usually absent.

Curvature of the Spine is very common in Friedreich's ataxia. Soca, quoted by Hallion,²⁴ says it occurs in half the cases. The photos of Cases II. and III. show the usual marked conditions. In the latter, the main convex curve is to the left, and is greatest about the ninth dorsal vertebra; in the former it is to the other side; it is both longer and more marked, and it is worst at the sixth or seventh dorsal vertebra. Perhaps the convexity to the *left* in the young woman is associated causally with the greater weakness in her right arm from her acute illness. In the common lateral curvature of adolescents, the main dorsal convexity is usually an exaggeration of the natural curve to the right, as in the brother (Case II.). In both there is kyphosis in the cervico-dorsal region, in the man this being so marked that the vertebral spines are nearly horizontal when he is sitting up. Observe how close the numbers on the spines appear to be, the effect of photographic distortion (fig. 2).

Hallion²⁴ has made a careful study of the spinal curvatures in nervous diseases, especially syringo-myelia, infantile paralysis, Friedreich's disease and locomotor ataxia. In all, the gray substance of the cord is affected in a different degree: in Friedreich's disease a pathological change is found in Clarke's columns, where some localise vaso-motor centres. Have we here, too, he asks, nervous lesions capable of causing an alteration of the bones, and specially of the vertebræ? He quotes Soca, to the effect that muscular weakness alone is sufficient to account for the condition. The want of tone may involve ligaments and bones too, as in the common scoliosis, but of this there is no proof.

We do not require to postulate any marked paresis of the spinal muscles; adolescents, who are subjects of the ordinary lateral curvature, are fairly muscular. A want of balance about the groups of muscles is more likely to start the scoliosis.

REMARKS ON THE CONDITION OF THE SENSORY SYSTEM.

The *sensibility of the skin* is frequently quite normal as regards pain, touch and temperature, but on the other hand it is not uncommon to find a slight dulling. In Cases I. and IV. the condition is practically normal; in Case III. there is some delay in recognising touch in the lower extremities, but localisation is correct. At one time there was absence of sensibility to touch on the dorsum of the left foot and the middle toes.

Case II. shows slight diminution of acuteness in recognising touch in the legs, but pain is felt normally; in the upper extremities sensibility to touch is normal. The curious thing about his condition is, that at one time (1884 and earlier) there was decided anæsthesia to touch and pain below the level of the iliac crest, this being proved by the examination in the Infirmary (1884), and also by the fact that about the age of twenty (1883) he burnt his foot by standing on a hot coal without feeling any pain.

It is interesting to compare this analgesia with the condition in the three cases recorded by Klippel and Durante, quoted in Londe's Thesis.⁶¹ These cases are amongst the most important selected by Marie, from which to build up his type *héréd-ataxie cérébelleuse*, but they are distinguished from the rest of this group, as well as from ordinary Friedreich's ataxics, by the disturbances of sensibility.

In Louis H., in 1892, tactile sensibility was abolished in legs and feet, much diminished in forearms and hands, and diminished in face; no plantar reflex; sensibility to heat recognised everywhere; retraction of field of vision. In 1894, Londe found analgesia in hands and legs; sensibility to touch present; sensibility to heat and cold weakened in the analgesic parts, retraction of field of vision hardly perceptible.

In François, sensibility to touch and cold, abolished below the knee in 1892, two years later no affections of sensibility.

In their sister, complete anæsthesia to all forms of sensibility on the internal aspect of the leg, in 1888; whereas a few years later, sensibility in her too, was normal.

Londe gives various reasons for his opinion that these sensory disturbances were due to hysteria. Certainly one is justified in regarding the condition as functional, and it has to be remembered, that many of the symptoms in nervous diseases, which have a well-marked organic foundation, are in all probability purely functional.

One of Stintzing's cases (*Münch. Med. Woch.*, No. 21, 1887) had complete anæsthesia in the lower extremities, and Senator's⁵⁵ second case felt on the feet needle pricks and differences of temperature, only slightly or not at all.

Déjerine⁴ describes a brother and sister who were affected with marked sensory troubles of the extremities, diminishing towards the trunk, and also corresponding muscular atrophies; these he regards as true cases of Friedreich's ataxia, with very marked exaggeration of certain symptoms, sometimes met with in this disease, though only to a very slight degree. He refers them to lesions of the peripheral nerves and sensory roots.

It is interesting to note that Risien Russell, by his recent experiments (*Trans. Phil. Soc.*), showed that anæsthesia and analgesia occur in the partially paralysed lower extremities, after removal of the cerebellum.

Subjective sensations are also occasionally found. In Case II. these used to be, formication in the soles of the feet, and tingling sensations in the legs. Case I. had for a short time pains in the back of the thighs.

Lightning pains occurred in one of Déjerine's cases, very much the same as in locomotor ataxia. Bramwell¹³ reports a family group of three cases, all of whom suffered from lightning pains at one time or another, one from gastric crises and all from exaggerated sensibility. As he notes, one of Charcot's showed lightning pains.

Other cases have involuntary spasms, associated with itching or other sensations (Tresidder's).

MENTAL CONDITION.

A normal mental condition may still be looked upon as characteristic. Tresidder's³³ three cases showed remarkably high intelligence, one of them having taken an

honours degree at Cambridge. Ladame's¹ case had a very vigorous memory and intellect. Hector Mackenzie's⁴⁸ was bright and intelligent for his station in life. My own cases are not affected mentally in the least degree. The *expression* of face is often unintelligent (as in Case IV.), and likely to mislead the observer. Cases that have normal mental power may be easily excited to uncontrollable laughter (Brock,⁴¹ Ladame, Bramwell).

Mental defect is noted in some cases. In one case, epileptic attacks and dementia (Hoffmann, of Heidelberg⁷¹); in another imbecility (Power, quoted by Gowers, *St. Bart.'s Hosp. Reports*, 1882, p. 305); in two sisters, marked mental defect (Londe and Lagrange⁷²); in one of Taylor's⁶⁵ increased mental irritability, with pronounced feebleness of intellect; in the other, not related to the first, epileptic attacks; in Nolan's⁶³ three cases, congenital idiocy; in Vinaj's⁶⁰ mental degeneracy.

In some of these, however, the diagnosis is more than doubtful. In itself it would not be surprising to find mental defect added to the usual symptoms; it would represent a further stage of the neuropathic tendency.

An American professor (Halbert,⁶⁸ Hahnemann Hospital, Chicago) gives a report of two cases of inherited syphilis with spastic-paralytic symptoms, as cases of Friedreich's ataxia, and in regard to the elder—eleven years of age—makes the quaint observation that "his mental condition is not greatly impaired, except that it is infantile from the lack of development!"

More will be found on this subject of mental condition under the heading of diagnosis (page 105).

REFLEXES.

The condition of the *superficial reflexes* in Friedreich's ataxia may be regarded as normal.

In Case II. plantar reflexes are well marked on the right, slightly on the left. The cremasteric and abdominal reflexes are well marked. In Case III. all are well marked, and the plantar exaggerated. In Cases I. and IV. the superficial reflexes are normal.

In Berdez's⁷⁹ case, superficial reflexes were very lively. In three of Besold's cases⁵⁴ they were normal, in the fourth, abdominal reflexes were absent.

Variations in degree of cutaneous reflexes are of course of constant occurrence in healthy persons, and the point under discussion is chiefly of importance as helping to distinguish Friedreich's ataxia from tabes. In the latter affection the plantar reflexes, especially, are often absent, this being due to the loss of cutaneous sensibility.

Much more important is the *condition of the deep reflexes*, of which the knee-jerk alone may be considered here, as it is easily tested.

In the four cases I have described the knee-jerks are completely absent, and this is the case in the great majority of examples of Friedreich's ataxia.

Crozer Griffith (*loc. cit.*), in his analysis of a hundred and forty-three cases, describes the abolition of the knee reflexes as very early and very constant. In ninety cases only was their condition mentioned, and curiously enough in as large a proportion as twenty-two of these they were present, in six being actually exaggerated. It is true that Griffith's statistics are compiled from an unsifted mass of cases, but one can certainly not go as far as Ladame, who holds that the abolition of patellar reflexes is an absolute rule in this disease, and that one would do right to mistrust the diagnosis of Friedreich's ataxia whenever the knee reflexes are not lost, although all the other symptoms would seem to confirm it (BRAIN, vol. xiii., 1890, p. 492).

When one comes across a typical family of Friedreich's ataxics, such as that described by Tresidder,⁸⁸ and finds the knee-jerks completely absent in one and present in the others, one is forced to the conclusion that this test is not infallible. Moreover, it seems probable that the disappearance of the knee-jerks is, sometimes at least, not one of the earliest symptoms.

In Senator's^{34 35} first case, these were found to be simply weakened one year after the onset of the illness. After the lapse of another period of eighteen months the right was quite gone, the left could be obtained only by

Jendrassik's method. A sister of this patient, who had been ill for twenty years, had no patellar reflexes.

Brock's⁴¹ second case, a man aged 27, had knee-jerks, though they were absent in his ataxic brother and nephew.

Erb⁹ and Mendel¹⁰ both showed cases in 1890 with tendon reflexes, in Erb's exaggerated. In one of Mendel's the jerks were present in May, but absent in the following November.

Londe⁹ would class Erb's cases with Marie's hereditary cerebellar ataxia simply on account of the exaggeration of the reflexes, though vision was normal and talipes present.

In the cases I have quoted, as in two of Bramwell's,³² the presence of the jerks may not be sufficient to blur the picture of Friedreich's ataxia, but it is a different matter when one finds that, along with retention of knee-jerks, there are other important deviations from the type. Bouchaud,⁶⁰ for instance, records a case which differs from Friedreich's ataxia not only in retention of the jerks, but in the impaired mental condition, hallucinations, affection of vision, nature of gait, and a too-pronounced nystagmus. I might refer further to the cases reported by Wallace Anderson,³⁶ Vinaj,⁸⁹ one of Nolan's,⁶³ T. S. Wilson's³⁹ first and fourth, as examples that are open to the above criticism.

However, the pathological anatomy of the disease varies, as we shall see later on, to such a degree that it is easy enough to suppose the knee reflexes might be free from interference in an occasional case. Such a case might be comparatively free from sclerosis in the lumbar portion of the cord.

Of the *visceral disturbances*, that noted in Cases II. and IV., viz., slow inception of the act of micturition, is not so very infrequent (thirteen times in Griffith's analysis).

ORDER IN WHICH THE SYMPTOMS APPEAR.

This is not easy to fix, but usually the ataxia of the legs shows itself first. This seems to have been the way in all my cases; in Case I. the speech was very soon involved, but so slightly that it would have remained unobserved,

but for the brother's thorough knowledge of the symptoms that might be expected.

In James's,⁴⁰ and one of Brock's⁴¹ cases, the spinal curvature seems to have been the first symptom noticed. Both cases were lads of 18. One of Besold's⁵⁴ showed the ataxia first in the arms. In one of Mendel's,¹⁰ the ataxia of the arms was noticed in the thirteenth year (writing becoming worse and more illegible), while it was two years later before the gait was observed to be changed. Griffith, in his analysis, states that the lower extremities were first attacked in 114 out of 143 cases, the arms in 10, and arms and speech simultaneously with the legs in 8.

MODE OF ADVANCE OF SYMPTOMS.

Some writers speak of the steady progress of the symptoms of the disease, and contrast this with the remissions of insular sclerosis. The mode of advance of the disease varies, however, to a great extent in different cases, and one may often distinguish remissions, or at least periods in which the symptoms are quite stationary.

Case II. showed slowly advancing symptoms till he was seventeen; then for two years there seems to have been a remission. At nineteen there was rapid deterioration; scoliosis and paresis, with unfitness for work, developed. Then the disease passed into its present chronic and almost unchanging state.

In Case III. the periods are less distinct; for years there has been hardly any alteration, but at about the same age as in her brother's case the extreme difficulty in walking developed rather rapidly.

The other cases (I. and IV.), show a steady advance, much more rapid in the latter than in the former. The age of nineteen or twenty will probably be critical in the history of Case I.

PERIOD OF ONSET.

The date of onset of Friedreich's ataxia is by no means easy to state. Friedreich himself, judging from his own cases, considered that it was to be regarded as a disease of

puberty. Most cases undoubtedly begin before this, the seventh and eighth years being most common (Gowers⁸⁸), and it is rarely that one commences after sixteen. The onset is so gradual, unless precipitated by an acute illness, that it may be almost impossible to definitely mark it. One of the latest cases seems to be Auscher's,—the woman being twenty-five when symptoms first appeared, but as we shall see, the diagnosis of this case must be accepted with reserve.

Londe (*Thèse*, p. 154) refers to two cases lectured on by Charcot, in 1893—brothers, in whom the onset was at twenty-five and twenty-one respectively.

In my cases, the fourth took ill about the age of eleven, but the beginning was insidious.

Case I. was seven when she had the illness, which seems to have been very soon followed by slight leg symptoms.

Case II. had a prolonged illness, possibly hydrocephalus, between his third and sixth year; there was, however, a long interval of good health before the ataxia became noticeable at the age of twelve.

Case III., like her sister, took ill immediately after a severe illness, though in this patient it seems to have affected the brain. She was then fifteen years of age.

The general statement holds good that it is a disease of the developing period.

DIAGNOSIS.

The diagnosis of Friedreich's ataxia seems to become more involved as time goes on, and cases accumulate. In typical family cases, such as the three first above recorded, there can be little difficulty if one is at all cognisant of the group of symptoms that are required to define the affection. But the symptoms are by no means few, and not infrequently some are wanting. Hence, on the one hand there are close relationships with some other nervous disorders, and on the other more or less atypical cases occur, as to whose exact position there is disagreement amongst the highest authorities. And even in regard to some cases

where an autopsy has been made, and where very decided lesions have been found, there are differences of opinion; the clinical symptoms may put a case into one category, and the pathological anatomy into another.

Senator³⁵ considers that Friedreich's disease is a clinical conception, which would exist without any anatomical substratum, just as the notion of diabetes mellitus is independent of anatomical ideas. The pathological anatomy of some forms of diffuse myelitis ("combined sclerosis") is very like that of some cases of Friedreich's disease, but the clinical appearances are so distinct that no one could confound them.

The following case, one would almost think, had been diagnosed as Friedreich's disease after the *post-mortem* examination. It forms the basis of an elaborate paper by Tedeschi,³⁶ entitled "*Die Friedreich'sche Krankheit — kritische und pathologisch-anatomische Untersuchung.*"

The patient, an orphan boy, aged 17, was admitted into the asylum at Florence in May, 1893. Two cousins in an asylum with mania. First symptom five years before, through a fright; tremor of right hand gradually involving arm, then left arm, next the legs. The tremor was like chorea. No pain; no remarkable affection of sight or hearing; no deformity. In asylum, increase of choreic movements, head involved. Walking difficult through inco-ordination, like an advanced tabetic. Speech slow and distinct; irritable, quarrelsome; of limited intelligence; could hardly understand what was said. Died in February, 1894, of tuberculosis.

One would like information about the knee-jerks, nystagmus, scoliosis, ataxia of upper extremities, and more about the sight and sensory system. But even with the details given—the mental affection, the uncommon onset, the asymmetry of onset, the absence of cerebellar type in the gait—I would have no hesitation in saying one does not, from the description, get the clinical idea that is requisite for the diagnosis.

The autopsy showed, besides tuberculosis in various thoracic and abdominal organs, a degenerative process in

the posterior ganglia and posterior roots; marked sclerosis of the posterior columns and Lissauer's zones (reminding one in extent and arrangement of the lesions in tabes dorsalis); marked atrophy of Clarke's columns and a slight change in the direct cerebellar tract in the cervical region. The cord was strikingly small, and there was chronic lepto-meningitis spinalis, specially posteriorly. The histological examination is most carefully described, but does not help the student of Friedreich's ataxia, if the above criticism is justified.

This, like Case IV., is a so-called "isolated" case—that is, he is the only one of the family affected with the disease. We are thus deprived of one important aid in diagnosis, namely, the "family" character.

Isolated cases are found to be not uncommon, and a survey of a large number in the more recent literature makes it clear that there are no outstanding characters which distinguish them from the true family cases occurring in collaterals. Divergences from the complete clinical type occur here and there, just as they do in the family form; and just as each family affected with Friedreich's disease shows its own special peculiarity (Marie²⁶), so one must make a certain allowance for individual difference in the isolated form. This much being granted, there is still room for the criticism that in some of the published cases the narrator has shown a somewhat excessive desire to fit his irregularly shaped peg in the round hole of the "clinical picture." Take for instance the personally observed case given by Dana¹⁷ in his article on Friedreich's disease in Keating's "Cyclopædia of Children's Diseases."

A lad, 18 years old, got an injury to the back of his head five years before. Soon he began to feel dull, heavy, and drowsy, had pains about knees and ankles, severe pains in thighs, pains in lumbar region, constant and severe cephalalgia, formication in legs, back, and shoulders. Stomach symptoms, vomiting occasionally, staggering gait, ankles unsteady. Felt as if tongue were tied back, consequently speech thick. Face became and remained flushed; much

dizziness; persistent polyuria (180-190 oz. daily, s.g. 1005-1010, no sugar or albumen). Diplopia.

So much for the history.

Physical examination showed a vaso-motor paresis of face and hands; "drunken" gait; no choreic or tremulous movements, except that head oscillated to and fro when he stood; slight tendency to flat-foot; no scoliosis; olecranon and *patellar reflexes normal*. Sometimes headaches, often vertigo; trunk ataxia; motor ataxia in using arms and legs; some nystagmus, which disappeared. Intelligence good.

Whatever this case is, it is not a typical case of Friedreich's ataxia. The history gives unusually copious and varied abnormalities of sensation, and as the author himself says, the origin from a blow on the head, and the cerebellar ataxia, are suggestive of tumour. The other symptoms all seem to point in the same direction (vomiting, polyuria, retention of knee-jerks, giddiness, &c.).

A case is reported by T. S. Wilson³⁹ (Case III.), as interesting from showing the symptom of glycosuria. The patient died at the age of 10 of diabetic coma, had been ill for five years, showed ataxia of gait, and slowness and hesitation in speech; a certain amount of clumsiness of hands; absence of knee-jerks; equino-varus; lateral curvature. There was no choreiform unsteadiness; no tremor on movement: great emaciation (from the diabetes, no doubt).

The case is certainly very interesting, and though the details are meagre, I should be more inclined to class it with the peripheral neurites than with Friedreich's ataxia. The sensibility was normal, but this may occur in a case of diabetic neuritis.

Case IV. of the same physician seems really to be what it was sent into hospital as, viz., St. Vitus' Dance. The unsteadiness became, in a fortnight from onset, so marked that the least thing would knock him down. There were muscular weakness; lordosis; increased knee-jerks; spastic tendency. Much improvement in hospital. Whole duration of illness to time of examination, only eight months.

Putting aside such cases as extremely doubtful, one finds considerable differences in those that remain.

Ladame's,¹ M'Caw's,⁸¹ Baskett's,⁷⁶ Mackenzie's,⁴⁸ and Berdez's,⁷⁹ to name only a few, may be looked on as quite typical.

My Case IV. shows the essential points, namely, onset in childhood; first symptom (ataxia) in legs; progressive course involving legs and arms; affection of speech; swaying and choreiform jerks; absence of knee-jerks; Romberg's sign; nystagmoid movements; and amongst characteristic negative signs, absence of sensory defect; clearness of intellect; normal pupils; normal ocular fundi.

On the other hand, there are some points of divergence. Though the disease is now far advanced, there is no club-foot or extension of great toe and no marked scoliosis. Cases reported by Dreschfeld,⁸⁰ D. Inglis,²⁰ Besold,⁸⁴ (Cases III. and IV.), Zabłudowski,⁸⁴ and many others quite undoubted, show no abnormality in this respect.

The hesitancy in beginning the act of micturition is not very uncommon, and is present in Case II. The cramp, curiously enough, affects only the left leg. According to Griffith, spasmodic contraction of muscles occurred in 21 of his 143 grouped cases, and was usually in the lower extremities.

Chorea is one of the diseases for which Friedreich's ataxia has several times been mistaken—Ladame, Londe and Lagrange,⁷³ Bramwell,¹² first case, one of Ormerod's (*Med. Chir. Trans.*, vol. lxxviii., p. 149), &c.

It is to be distinguished by the slowly progressing character, the absence of knee-jerks, and the character of the involuntary movements. In marked cases of either disease there can be no difficulty. The tongue, for instance, in a choreic is jerked up and down, in and out, while in an ataxic it shows an irregular tremor with no rapid jerks. The face is usually not affected in Friedreich's disease, though in Cases I. and II. slightly marked exceptions may be noted. *Chorea* does occur in families. "Though the proportion is not great, the family tendency in some cases is very marked." (Gowers.)

Huntingdon's *chorea* is hereditary, begins in adult life, is accompanied by mental symptoms, and is always fatal.

Hereditary Cerebellar Ataxia.—It is by no means certain, as yet, whether this is to be looked on as a definitely marked disease with independent pathology. So far it would appear that the symptoms may vary to an extraordinary extent, and the autopsies on which Marie⁴⁶ based his reasoning amount to only two amongst the sixteen cases that he reckoned up as belonging to the new type.

Londe wrote a lengthy thesis on the subject (*De l'héréditaire ataxie cérébelleuse*), and added three new cases.

Marie recognises in all the character of staggering and cerebellar inco-ordination, and connects this with the cerebellar atrophy found in the two autopsies. He is supported in this by the most recent observations on the function of the cerebellum.

Both Friedreich's disease and the new type are family diseases, the latter beginning more commonly in adult life, the former in childhood. But as we have seen, Friedreich's disease may begin at twenty-five, and on the other hand, Fraser's patient (*Glasgow Medical Journal*, 1880) showed signs in infancy. This is one of the two cases of cerebellar hereditary ataxia in which an autopsy was made.

The various other points, when gone into, show a close parallelism in the two types. The symptoms of hereditary cerebellar ataxia may be summarised thus from Londe :—

(1) Slow and gradual affection of legs, in standing or walking.

(2) Occasional pains in legs and loins.

(3) Usually in three years or so, uncertainty of hands and arms.

(4) About the same time speech and vision involved. (In Londe's own cases the eyes were not affected, if one excepts nystagmiform movements).

(5) Reflexes remain or are increased. This is of course very exceptional in Friedreich's disease.

(6) Sometimes mental enfeeblement.

(7) Death at advanced age, usually from intercurrent disease, frequently of the lung.

It is much easier to say that the cases collated by Marie and Londe are not cases of Friedreich's ataxia, than it is to

admit that they should be put together to form a homogeneous group. But the theory put forward by Marie is at least plausible—that the same kind of degenerative process may be at work, involving the cerebellum only or mainly, in his cases—the spinal cord in Friedreich's disease.*

RELATION OF MENTAL DEFECT TO FRIEDREICH'S ATAXIA.

In cerebellar hereditary ataxia *mental enfeeblement* is often present. This leads one to consider the question "What weight should be attached to this point in making a diagnosis of Friedreich's ataxia?"

The pure type of this affection involves no mental change unless occasional irritability and a tendency to uncontrollable laughter; even this is exceptional. In fact, if we exclude the nystagmus and the speech affection, in regard to the causation of which we are quite in the dark, we have no evidence of involvement of the cerebrum, one might almost say, of the encephalon.

When the mind is affected we may take it for granted that the higher cerebral centres are diseased. We have seen that this condition is now and again present in cases that in other respects correspond to Friedreich's ataxia, but an examination of some cases recently reported makes one disposed to think that they should be regarded, not as Friedreich's ataxics, but as transitional to other types, *e.g.*, cerebral diplegia.

Amongst these I would reckon Nolan's⁶³ three cases of Friedreich's ataxia with congenital idiocy. These were a sister, aged 22, and two brothers, aged 15 and 10. Their mental condition was one of low intelligence; volition rarely manifested; memory defective; no delusions; speaking only in answer to queries; anxious to please; good humoured; asexual. All had nystagmus, internal strabismus; imperfect laboured articulation, many sounds not being uttered; ataxia in the arms and legs, with Romberg's sign.

* Compare Dr. A. Meyer's paper in the last (Autumn) number of *BRAIN*, with a reading of which I have been favoured by the Editor.

The thyroid was enlarged in the two elder. "Universal partial analgesia," and "drop-ankle" were present in the same two. Knee-jerks absent in oldest and youngest, exaggerated and combined with ankle clonus in the other. No scoliosis or Friedreich's foot; a "suggestion" of *main en griffe* in two; no visceral disturbances, or ocular troubles other than those referable to the external muscles. The true jerky or choreic movements do not seem to have been very marked. The disease was observed soon after birth.

There are indications here of an extensive defect of development in the whole nervous system. The cases seem to be more closely allied to cerebral diplegia than to Friedreich's ataxia.

Nearly related to these are the cases of a brother and sister (Bouchaud⁸⁰), who at the ages of six and seven began to show mental enfeeblement, inco-ordination of the four extremities, and tabeto-cerebellar gait. No paralysis, nystagmus, or kypho-scoliosis. Speech slow but well articulated. The dementia steadily increased. Sight became lost; speech unintelligible; chewing and swallowing slow; rigidity of muscles increased, though knee-jerks were in one case absent, in the other diminished. The autopsy showed double sclerosis of lateral pyramidal tracts, diminution of sheathed fibres in cortex, anterior and posterior columns healthy; ganglion cells of anterior and posterior horn diminished in number.

The same category may be said to include Vinaj's⁸⁹ family, about which he asks "Is this a hitherto unrecorded kind of case or a modification of Friedreich's ataxia?"

Father, mother, and four children alive, second dead. Signs of nervous predisposition.

(1) Eusebio, well till age of 5, then difficulty in speaking and walking; intelligence affected, ending in idiocy; steady degeneration. When 11, paralysis of legs; paresis of arms; physical degeneration; no control of bladder or bowel.

(2) Marietta showed same symptoms at 5; died of accident at 9; by that time had lost the power of speech.

(3) Iside, age 6; slight nystagmus; knee-jerks normal;

gait uncertain; staggering; speech hesitating; scanning; memory weak.

(4) Egidio, aged 5, large head; unequal pupils; nystagmus; hesitating speech, pronunciation like a child of three; normal jerks.

(5) Healthy child of fifteen months.

In these cases the evidence against a diagnosis of Friedreich's ataxia consists in retention of knee-jerks, absence of talipes and scoliosis, and, most important of all, early and marked loss of intelligence. The family tendency, age of onset, ataxia, affection of speech, nystagmus and preservation of sensibility bring it into relation with that disease. The author rejects as diagnoses:

(1) Little's disease (origin; absence of special gait; absence of spastic rigidity; psychic phenomena).

(2) Spastic tabes.

(3) Primary progressive myopathy.

(4) Chronic encephalitis: primary cerebral sclerosis of infancy (time of onset; normal mode of birth; absence of convulsions, of contractures, of paresis or paralysis; the mental phenomena).

The cases are quite analogous to a number of others described by various authors, many of which are brought together by Higier⁷⁸ in his study of the rarer forms of paralysis of a hereditary and family character. After detailing four cases of his own in one family, he reviews the cases of cerebral diplegia reported by Sachs, Freud, Strümpell, Erb, Tooth, &c. The age of onset and the character of the affection are pretty constant in all, but each family has its distinctive characters. If one takes as the chief symptoms, onset in early childhood, spastic or flaccid paraplegia, idiocy, nystagmus, affection of speech and sight, one sees that Vinaj's and Nolan's cases are perhaps more nearly related to this group than to Friedreich's ataxia.

Higier's own cases are specially interesting, in relation to the point now under discussion, from the fact that they showed disturbances of co-ordination; weakness and awkwardness in upper extremities; intention-tremor; great

difficulty in sewing, knitting, and writing; in lower extremities weakness and stiffness followed by spastic paresis in gait, in one spastic ataxia for a time. He draws attention to the "little-known fact" that inco-ordination does now and then occur with increased tendon reflexes, quoting reports by Sachs, Haushalter, König, Dreschfeld. I may abridge the descriptions of Haushalter's cases:—Parents near blood relations, father alcoholic. Four brothers and sisters, normal births. Two eldest affected, now 11 and 10 years old. Symptoms first while at breast; rapid advance; inco-ordination of four extremities; fixed position of arms, of hands and fingers; slight rigidity in extremities, more in tongue and face; choreiform jerks when at rest; increased knee-jerks; *pes equinus*; anomalies of speech; disturbances of swallowing; imbecility. Nystagmus only in eldest.

These cases differ from Friedreich's ataxia in the retention of knee-jerks, the imbecility, the rapid advance at an early age and some other minor points, but they are transition forms between this disease and cerebral diplegia.

Collins⁷⁸ reports a case quite allied to those transition forms:—A child, 11 years old, of tubercular heredity (whose brother died in his second year, defective mentally and nystagmic), began to walk at end of first year. From third year gait stiffer, often falling; high myopia in fourth year; did not get on at school; when seen at age of 11, choreic movements of lower extremities; speech indistinct; exaggerated action of mimetic muscles under emotion; old face; gait cerebellar-ataxic, legs kept apart; ataxia of arms, especially left; increased knee-jerks; occasional ankle clonus; explosive ataxic speech; imbecile; sensibility and sphincters intact; no muscular atrophy; progressive paresis; slight nystagmiform movements; slight club-foot.

Here there is clearly a defective developmental condition of the nervous system in both cord and brain.

It is interesting to note how often in these cases which present a combination of the symptoms of Friedreich's ataxia with idiocy, the knee-jerks are retained. This tends to confirm the very high position of Westphal's sign as a diagnostic mark.

To the above list I might add the two cases of Londe and Lagrange,⁷⁰ who showed imbecility with the exceptional athetoid attitude in walking, and Hoffmann's⁷¹ case, where there were imbecility and epilepsy, a case he himself records as doubtful in regard to diagnosis.

One sees from this review that when the degenerative process has involved not only the spinal cord, but the higher cerebral functions, other symptoms besides simple imbecility or dementia are almost invariably added to the symptoms of Friedreich's ataxia. And in such cases there must always be a doubt as to how far the changes in the cord, with their accompanying symptoms, are secondary to the cerebral affection, a doubt which a *post-mortem* examination might not clear away. Mental defect seems a much more important sign than the retention of the knee-jerk in making one doubt a diagnosis of Friedreich's ataxia.

DISSEMINATED SCLEROSIS.

We pass now to consider the differentiation of Friedreich's ataxia from disseminated sclerosis. This disease is said never to occur in families (Marie), but a difficulty may arise when one meets an isolated case of Friedreich's ataxia. A good number of cases have been recorded of its occurrence in early childhood, but the proof of this anatomically was first given by Eichhorst in a paper "*über infantile und hereditäre multiple Sklerose*" (*Arch. für Path. Anat. und Physiol. &c.*, Bd. 146, Hft. 2, p. 173). A well-marked case of insular sclerosis can present no difficulties; the intention tremor is coarse and jerky, much more so than in Friedreich's disease; the nystagmus is more pronounced; the course of the disease is liable to fluctuations in much greater degree; the tendon reflexes are exaggerated; ocular paralysees and affections of the optic nerve are frequent.

But the group of symptoms due to insular sclerosis varies much. The gait is at times cerebellar in type; knee-jerks are occasionally absent; scoliosis may be present; the speech may be similar to that in Friedreich's disease.

Brissaud (quoted in Londe's thesis) says it may be abso-

lutely impossible to distinguish an isolated case of *hérédo-ataxie cérébelleuse* from a case of *sclérose en plaques*, and the following cases may show the difficulty in diagnosing it from Friedreich's disease. Bouchaud¹⁰ describes this one as "Maladie de Friedreich":

Boy, aged 18, always rather delicate, fairly sharp at school. First symptoms at 12, viz., awkwardness of hands; tremors of upper extremities; difficulty in walking. Apprenticed to saddler at 14, but made no progress. On examination asymmetry of face (left smaller); left arm and leg more awkward and weak than right; intelligence much below normal; speech difficult to understand; does not speak much; no choreiform instability when at rest; static ataxia when trying to stand; inco-ordination; intention tremors; gait not tabetic; staggering, requires support; "tendency" to talipes equinus; exaggerated extension of great toe; knee-jerks both retained, left exaggerated; nystagmus marked; slight lateral curvature; acuteness of vision much diminished; hallucinations; no evidence of loss of sensibility.

This case differs from Friedreich's disease in quite a number of particulars, especially the mental condition, retention of knee-jerks, affection of vision, gait, asymmetrical appearance of face, greater awkwardness on one side. On all these points a diagnosis of disseminated sclerosis would be much more adequate.

Ewart¹¹ gives a case of a female, aged 19, of "isolated" type. Symptoms began at 16. Menstruation first irregular, then ceased; slight affection of speech, only temporary, a year ago; knee-jerks exaggerated; slight tremor of hands on purposive movements; slight lateral nystagmus; marked ataxia of lower limbs; slight loss of sensation; functions of brain intact.

He thought disseminated sclerosis could be excluded by the insufficient degree of tremors and nystagmus and the normal cerebral condition. Friedreich's disease is, however, a doubtful diagnosis, on account of the disappearance of speech affection and the exaggeration of the jerks. These points would bring it nearer to disseminated sclerosis.

Probably, in a case like this, continued observation of the course would settle the question of diagnosis.

LOCOMOTOR ATAXIA.

The diagnosis from ordinary locomotor ataxia can present little difficulty. A comparative table of the symptoms is given in Ladame's paper. A difficulty comes up when cases of Friedreich's disease have to be distinguished from tabes commencing in childhood. In the latter there is usually a syphilitic heredity, but not in the former. In a summary of ten cases Hildebrandt of Berlin (27) found that six began before tenth year, four before fourteenth. In some cases ataxia of extremities takes the leading place, lancinating pains in two, girdle sensation in four. Loss of patellar reflexes in all cases, eyes three times normal, nystagmus twice present, reflex paralysis of pupil twice, disturbances of sensibility in only half, disturbances of bladder six times. Moreover, the speech often shows disturbances on account of slight cerebral affection. Fortunately the symptoms are so grouped usually that there is not much difficulty in reaching a decision.

Raymond⁹¹ discusses this subject at some length in a recent paper. He first records two cases, father and son. The father was a well-marked tabetic, showing, in addition to pre-ataxic symptoms, disturbance of superficial and deep sensibility, loss of knee-jerk, urinary incontinence, ataxia of the four limbs, characteristic gait, Romberg's sign, and marked eye affections. Onset 1883, a few years after his marriage. Of his family, two girls and one boy were healthy, but the youngest boy had a nervous disease. He had always been rather weak in mind. In October, 1895, at the age of 15 he noticed one evening that he could not run, though he could walk. There was loss of muscular power; he soon found difficulty in climbing a ladder; he let things fall, and lost sensation of touch in his hands. He had no pains, but a sensation of wearing tight boots, even when his feet were bare. Gradually he lost confidence in movements, his gait became uneven, and his speech was embarrassed. In

January, 1896, this was his state: Development good; slight upper dorsal scoliosis; convex to right; tabeto-cerebellar gait; Romberg's sign; much impaired sense of touch in hands; impaired sense of posture; loss of deep reflexes; fibrillar and jerky tremors of tongue; nystagmiform jerks; frequent foolish laughter; intellectual functions limited. At present (September, 1897) condition rather worse; sight gone; (discs "tabetic"); jerky movements of limbs and trunk more pronounced, especially on effort.

In discussing this case, Raymond points out certain resemblances to Friedreich's ataxia, viz., gait, scoliosis, speech affection, nystagmus, jerkings, loss of knee-jerks. On the other hand, it differs in the marked affections of sensibility of skin and joints, and the optic atrophy. The case resembles tabes in these respects, in the loss of jerks and in the motor inco-ordination, but differs in the absence of a pre-ataxic stage with lightning pains, Argyll-Robertson phenomenon, urinary troubles; also in the swaying character of the gait, the nystagmus, the scoliosis and the speech affection. It is, in short, a hybrid case, a transition form between tabes and Friedreich's ataxia. Going into the question of juvenile tabes, he points out its great rarity; he himself has not met with a case in five hundred of locomotor ataxia. Only ten cases can he find in literature, with a doubtful eleventh, but these are all very characteristic, not differing from the adult form unless in a greater tendency to early development of optic atrophy, this being unaccompanied by any delay in the advance of the malady. Most of the cases published as juvenile tabes are cases of Friedreich's ataxia, or closely allied thereto, and amongst those Raymond would place two recorded by Freyer, and quoted by Hildebrandt in the paper above referred to.

But tabes in the adult may take abnormal forms. Auscher's³⁰ "case of Friedreich's ataxia," with autopsy, presents several points of close correspondence to locomotor ataxia. The onset was not till the twenty-fifth year; there was mental degeneracy, and abolition of light reflex. The patient was a prostitute, and there were no other members of her family affected. She showed marked ataxia of limbs;

choreiform movements of head and trunk ; double equinovarus ; loss of knee-jerks ; nystagmus and affection of speech. Sensibility was retained, though in a case of marked mental defect it must be difficult to test it completely. Some lightning pains were present. The autopsy may be summarised thus:—Meninges normal ; cord much diminished in size, like that of a child seven years old ; cerebrum and cerebellum much smaller than normal, but free from microscopic changes ; *posterior horns* much atrophied in whole length, worst in dorsal region ; atrophy of fibres of Clarke's columns, number of cells in them diminished ; *posterior columns* sclerosed in whole length, worst in lower part, Goll's columns being more changed than Burdach's ; only a few medullated fibres in Goll's columns in a network of fine neuroglia fibres ; *Lissauer's zone* normal in lumbar and cervical portions, perhaps slightly degenerated in dorsal ; posterior roots and antero-lateral columns normal ; motor nerves normal ; sensory nerves showed a great number of non-medullated fibres, or fibres with an extremely thin sheath, perhaps embryonal.

The sclerosis showed what Déjerine had a few months before described as a pure gliosis : great abundance of fine neuroglial fibres arranged in whorls, absence of granule cells, no thickening of vessel walls or of septa connected with pia mater.

From the clinical account, as well as from the autopsy, one would prefer a diagnosis of locomotor ataxia, with an ill-developed brain, the congenital mental defect being possibly aggravated by a progressive paralysis, as Schultze suggests.⁵⁸ Criticism of a case like this is most desirable, for from it the following conclusions have been drawn :—

- (1) Friedreich's disease may start as late as 25.
- (2) Cerebellar defect may account for the clinical signs of Friedreich's ataxia (Senator).
- (3) Mental degeneracy, abolition of light reflex and other exceptional conditions may occur in Friedreich's disease.
- (4) The motor nerves may be normal, and the sensory show certain changes in Friedreich's disease (Leyden and Goldscheider).

(5) A full confirmation of Déjerine's views on the peculiar character of the histology in Friedreich's disease.

In making such inferences from a single case, the doubtful nature of the diagnosis must be borne in mind.

Disease of Cerebellum may cause symptoms like those in Friedreich's disease: reeling gait with a zig-zag character, difficulty in standing, nystagmus, loss of knee-jerk (usually not constant); but there is no tabetic character in the gait; the choreiform movements are probably absent; "the arms are usually steady, rarely they present some jerky inco-ordination" (Gowers), and in the case of tumour, pressure symptoms are generally present. Some cases of simple atrophy or hypoplasia of cerebellum have been described, but the symptoms are variable. In König's⁶⁶ case there was congenital cerebral diparesis, but no affection of inco-ordination at all, although the hypoplasia was very well marked.

"Oppenheim has observed cases of *cerebellar sclerosis*, and has had one anatomically investigated by Arndt. He regards cerebellar sclerosis as associated with a clinical picture composed of vertigo, cerebellar gait, ataxiform disturbances of movement, weakness of bladder, difficulty of articulation and motor weakness of the extremities" (Leyden and Goldscheider, *loc. cit.*, p. 597).

Although there is undoubtedly a general resemblance here, to cases of Friedreich's ataxia, there would probably not be much difficulty in distinguishing actual cases.

Cardossi¹⁹ publishes a case of "*a rudimentary form of Friedreich's disease*," which he admits is by no means typical.

Giuseppe P., aged 40, has a son aged 18, who has had tremor in the right hand from the age of 13. He himself began to have tremor in writing at 10. Right arm and body affected; never so bad as to prevent him from attending to his usual occupation; gait, tabeto-cerebellar; has had phthisis for last ten years, not much advanced; oscillation of body when standing, brief, rhythmical, not well marked; intention-tremor in upper limbs, as in disseminated sclerosis; slight tremors of tongue; speech slow

and hesitating; no nystagmus; pupils normal; sensory system normal; knee-jerks much diminished; no paresis or paralysis; no deformities of feet.

The symptoms are of too mild a type and too few for Friedreich's disease of thirty years' duration, and they can be resolved into tremor, but apparently not that of chronic chorea, nor of paralysis agitans. Possibly a functional disorder allied to one or other of those.

In this discussion on *diagnosis*, I have not attempted to set forth exhaustively the differences between Friedreich's disease and the various affections likely to give rise to difficulty. My desire has been rather, by giving groups of cases from recent medical literature, to bring out the close relationships that bind one type to another. From a textbook stand-point this would be most confusing, but one cannot be too often reminded of the somewhat arbitrary character of the boundaries. The more numerous the doubtful cases that are published the better, though unfortunately there is sometimes a tendency to magnify this symptom or minimise that, so as to get the case included in a favourite type.

PATHOLOGICAL ANATOMY.

The following are the more important records of *post-mortem* examinations in Friedreich's disease :—

(1-5) Friedreich and Schultze. (6) Everett Smith, *Boston Med. and Surg. J.*, 1885, vol. cxiii. (7) Newton Pitt, *Guy's Hosp. Rep.*, vol. xlv. (8-9) Rüttimeyer, *Virch. Arch.*, vol. cx. (10) Letulle and Vaquez.⁵ (11) Blocq and Marinesco.⁶ (12) Déjerine and Letulle.² and ⁶ (13) Guizetti.⁴⁹ (14) Mirto.⁵⁷ (15) Michell Clarke.⁵⁷ (16) Burr.⁷⁴ (17) Dana.⁷⁷

Autopsies by Menzel, Brousse, Kahler and Pick, Erlicki and Rybalkin, Rossi and Auscher, are rejected from the list as dealing with cases that can hardly be reckoned examples of Friedreich's ataxia.

For adequate reasons see Ladame,¹ Tedeschi,⁵⁵ and Schultze.⁵⁸

Most of the cases that have been fully described have shown extensive system lesions. They have been summarised by Ladame in 1889, and more recently Tedeschi has given an analysis of 15 up to 1896. The most common conditions are diminution in bulk of the cord, sclerosis of posterior columns (especially Goll's), and degeneration of Clarke's columns. Next in frequency is degeneration of the direct cerebellar tract, and the antero-lateral ascending tract of Gowers. The crossed pyramidal tract may be partially involved, but it is thought that the true motor fibres are unaffected unless in very advanced cases. In one of Friedreich's cases the direct pyramidal tracts were involved. Apparent involvement of motor tracts is explained by the presence in them to a greater or less degree of neurons with other than motor functions. The "system" arrangement of fibres is, in short, not an absolute one. Occasionally changes in the spinal ganglia have been observed. The posterior roots have sometimes been found atrophied. Changes in form of the central canal, and sometimes peri-ependymal sclerosis, have been seen (Déjerine and Letulle).

Ladame's opinion that the zone of Lissauer (fine fibres at outer side of posterior column stretching from apex of posterior horn to periphery) always remains intact can no longer be supported. He looked on this as a point of distinction from tabes. But in several cases (Letulle and Vaquez, Blocq and Marinesco, Guizzetti, Mirto) it has been found more or less extensively altered.

The degeneration of Clarke's column, especially of its ganglionic cells, seems associated with the sclerosis of the direct cerebellar tract. The view that these cells are trophic centres for the tract fibres is thus confirmed. In early tabes Clarke's columns are affected. There is an absence of the fine fibrous network, while the coarse fibres, the ganglion cells, and the fibres of the direct cerebellar tract arising from them are unchanged. In advanced tabes Clarke's columns are almost entirely degenerated, and then the cerebellar tracts are said to be sclerosed (*see* Leyden and Goldscheider,⁸⁷ p. 509).

There appears to be no constant anatomical character

which would allow one to say from examination of a section of the cord that one had to do with Friedreich's disease, and not with tabes, or a combined sclerosis. Déjerine and Letulle have endeavoured to find a distinctive character in the nature of the sclerosis. Those authors believe that the sclerosis in the posterior columns is purely neuroglial, a gliosis, consisting of fibrillæ intercrossed in different directions, and forming whorls like those described by Chaslin in the brains of epileptics. Further, the pial prolongations into these columns are unaltered, and not more evident than in a healthy cord. In tabes, on the other hand, there is thickening of the septa of the pia mater, and of the prolongations which start from them, thickening of the neuroglial tissue between the nerve tubes, very pronounced changes in the vessels, &c. A vascular sclerosis such as this they found in the *lateral columns* in their cases.

To meet the possible objection that they are making an extensive generalisation from two cases, they point out that similar changes have been seen by several authors, but that they have not interpreted them in their true sense. They quote to this effect from the histological descriptions of Friedreich himself, Newton Pitt, Rüttimeyer, and Blocq and Marinesco. Auscher, in his doubtful case, showed a similar condition at the Société de Biologie, July 26, 1890; and Déjerine, after examining his specimens, pronounced them examples of pure neuroglial sclerosis.

Schultze, of Bonn, re-examined by recent methods the cord of a case observed and described by Friedreich and himself in 1881, and he failed to find any justification for the distinction in the nature of sclerosis thus defined by Déjerine and Letulle. The whorls of fibrils were not present. The vascular thickenings were found in the posterior columns, especially between the columns of Burdach and Goll. He sums up his re-examination thus: No proper chronic meningitis. The degenerations are essentially a degeneration of the posterior roots and their connections in the posterior columns and posterior horns, disease of Clarke's columns, of the cerebellar fibres, and the pyramidal tracts—this so far as concerns the spinal cord.

Michell Clarke⁶⁷ describes the autopsy on a patient aged 14, a typical case except that the knee-jerks were normal, and that during the last few weeks of life he showed symptoms of cerebral tumour (cerebral vomiting, intense headache, optic neuritis, coma, &c.). A round-celled sarcoma was found growing from the under surface of the right half of the cerebellum, pressing on the pons and medulla. Apart from this no change in cerebellum. In cord no naked eye change except diminution in size. "No complete degeneration in any section, but a partial change chiefly in posterior columns, next most marked in lateral columns with involvement especially of the margin of the cord, with thickening in the pia mater and the walls of the vessels, and probably with some excess of the neuroglia."

A brother and sister were affected with the same disease. Had it not been for this, in view of the presence of a tumour in the cerebellum, one would have been apt to ask could not this have accounted for the symptoms. Certainly the symptoms were very severe at the end, and the tumour was apparently growing rapidly then, but it might have taken on an active character after a long comparatively latent course. The family had a strongly neuropathic tendency, and it is of course quite possible that two members of such a family should have Friedreich's disease and a third a cerebellar tumour without Friedreich's disease. Clarke himself states that the changes in the cord were by no means so marked as he was led to expect from the symptoms.

Dana's⁷⁷ autopsy is briefly recorded: it is accompanied by some photographs of cord-sections. In addition to smallness of cord, thickening of pia, sclerosis of posterior columns, sclerosis of lateral columns (crossed pyramidal and direct cerebellar tracts) and marginal sclerosis, he observed vacuolisation in the gray and white matter all through.

There is no note of the condition of Clarke's columns or Lissauer's tracts, nor could the author say whether the sclerosis was really a gliosis.

Burr's⁷⁴ autopsy was on a case that was eighteen

years ill. He found (1) marked degeneration of Goll's columns, slighter of Burdach's, from top of cervical to lower lumbar cord; (2) degeneration of lateral and anterior pyramidal tracts; (3) degeneration of direct cerebellar tracts. The sclerosis of lateral columns was not systematic. (4) posterior roots much degenerated, anterior intact; (5) slight affection of posterior horns with degeneration of ganglion cells, while in anterior horns only a very few cells were atrophied; (6) degeneration of cells in Clarke's columns; (7) walls of vessels slightly thickened.

The following are the chief points in Guizzetti's⁴⁹ case, who died of heart disease at the age of 28, after the nervous affection had lasted eighteen years: small size of posterior roots, spinal ganglia and cord.

(1) Lumbar region—sclerosis of posterior columns except a portion immediately behind posterior commissure and inner border of posterior horn, and also Westphal's posterior limiting zone; numerous degenerated fibres in pyramidal tracts.

(2) Dorsal region—posterior columns (with exception of part as in lumbar region), pyramidal tracts (numerous fibres), Clarke's columns and Lissauer's band all involved.

(3) Cervical region—Goll's columns as seriously involved as lower down, Burdach's less so, more healthy fibres in pyramidal tracts. Longitudinal fibres degenerated even into the nuclei of Goll's and Burdach's columns in the bulb, the horizontal fibres remaining unaffected. Some altered fibres even in anterior pyramids of bulb, contrary to usual condition.

Posterior horns atrophic. Microscopic condition like that described by Déjerine and Letulle.

In peripheral nerves, diminution of coarse nerve fibres and presence of numerous fine fibres.

Mirto's⁵⁷ case (death after five years' duration of illness, at age of 19) had systemic degeneration of posterior columns (Goll's, Burdach's and Lissauer's) of lateral cerebellar tracts, of Gowers' tracts; slight degeneration in crossed pyramidal tracts disappearing in cervical region, advanced sclerosis of Clarke's columns; rarefaction of

reticulum nervosum in anterior and posterior gray columns with partial cell-atrophy, partial degenerative neuritis in motor nerves (sensory not examined)—spinal ganglia affected as regards both cells and reticulum.

The latter two abstracts are taken from Tedeschi's critical digest, and I have not seen the clinical reports of these cases. But, supposing them to be typical, nothing of much importance has been added to what was known in 1889 to Ladame, or indeed Friedreich himself. The posterior roots and ganglia have shown marked changes, and in this respect and the coincident or consecutive affection of the posterior columns the similarity to tabes is obvious. In two cases (Guizzetti's and Mirto's) the principal nerves showed embryonic characters. In Mirto's the sensory were not examined, but the motor nerves were affected. Auscher's doubtful case presented similar appearances in the sensory nerves.

PATHOLOGY.

Various theories have been suggested to establish some causal relationship between the anatomical basis and the clinical symptoms. The general assertion that in a case of typical Friedreich's ataxia the symptoms are mainly spinal is one which has been widely though not universally accepted. As for the nystagmus, the affection of articulation, the giddiness and certain choreiform movements, though their origin is in all likelihood intra-cranial they have as yet no morbid anatomy. The strongest critic of the spinal origin of most of the symptoms is Senator.^{34 35} He has maintained the view that the disease is essentially *cerebellar*, that in its most outstanding features it depends on an arrest of development of the cerebellum or of single physiologically important parts of it, and that it is only in a somewhat advanced stage that other symptoms appear, referable to super-added degenerations, in cord and cerebrum. The cord changes may be concomitant or consecutive. In confirmation he quotes Marchi's conclusion from experiments: "Destruction of cerebellum entails degeneration of

lateral portions of antero-lateral columns, of anterior part of direct cerebellar tracts and of a certain number of fibres in the pyramidal bundles." He says that we know nothing of the symptoms associated with disease of Goll's columns alone, that a separate or simultaneous affection of Burdach's would give a clinical picture like that of tabes dorsalis, and would be more likely to do so as certain other peculiar changes in tabes (affection of Clarke's columns, of Lissauer's zone, atrophy of posterior roots and degeneration of peripheral nerves) are not wanting in hereditary ataxia. Moreover, by reference to the autopsies he finds that the condition of the cerebellum was not mentioned in eight out of the fifteen he analysed. In three cases of the other seven there was a recognised atrophy (Menzel's, Rossi's, Auscher's). In three cases nothing wrong was seen microscopically, and in one nothing either with naked eye or microscope. This is the weakest point about the theory—the want of anatomical evidence—and this objection has not been removed by the later *post-mortem* examinations, though Clarke's case given above is interesting in this relation. As regards the three cases he quotes, we have seen how unusual in some important respects Auscher's case was; as for Menzel's, it is generally regarded as not a case of Friedreich's disease, and Rossi's also seems to deviate considerably from the classical type.

The small size of the spinal cord in Friedreich's disease seems to have struck all observers, and Friedreich and Schultze first advanced the view that the columnar degenerations were secondary to an *arrest of development in cord and medulla*. Shrinking of the cord no doubt occurs subsequent to the degeneration as the connective tissue takes up less bulk than the nerve tubes, but it is questionable if this would account for all the difference; it would hardly explain why the cord is so much less in this disease than in other scleroses, even if the points emphasised by Schultze are allowed their full weight. This author speaks of the general atrophy of the cord in tabes in the dorsal and lumbar regions as due to long disuse of the lower extremities. So in Friedreich's ataxia one would, he says, still more look

for such atrophy, on account of the inactivity enforced in childhood and the longer duration of the disease than in the case of locomotor ataxia.

The theory of developmental error is supported by other facts—the presence of numerous fibres without sheaths in peripheral nerves (Guizzetti, Mirto), anomalous development of central canal of cord (Déjerine, who quotes similar observations by Ormerod, Pitt, Everett Smith); condition of syringo-myelia (Friedreich's fourth case, quoted by Schultze).

No satisfactory explanation has yet been given as to why the cord should be able to perform its functions for a series of years varying from two to twenty or more, and then show signs of inadequacy. The demands made on the spinal cord of a young lad are great enough to test its capabilities most thoroughly. In certain families, and certain isolated cases, parts of the central nervous system seem to be badly constructed. They may be looked upon as unfit to bear a long strain, and their decay and death gradually come about; not that the change is analogous to a senile change; at least the differences are so great as almost to hide any analogy that may be supposed to exist. If the process of disease or decay has once been started, it is likely to be aggravated by the struggle for existence among the cells that is taking place constantly at all ages, and so actively in youth. The neuroglia rapidly increases, and not only fills the space formed by shrinking of the nerve-tubes, but compresses and destroys healthy nerve tissues (Roux and Weigert, quoted by Edinger).

Edinger⁹⁰ a few years ago brought forward his *repair* theory, to explain the condition found in several nerve affections. In any exertion there is waste of nerve-cell and nerve-fibre, compensated in health by repair. If nutrition is defective through any cause, as a toxin (syphilis, diphtheria), or injury, or hereditary nervous weakness, repair will be insufficient and degeneration may be the result. Some of the cord tracts seem more easily acted on than others, viz., the posterior columns, pyramidal tracts, and, less frequently, the cerebellar tracts. A greater strain may be put on certain nerve-lines than on others. Thus, taking up

tabes, he would explain the involvement of the nerve apparatus connected with the erect posture, the anæsthesia of soles (pressure on soles), affection of muscle sense, of eyes, &c.

In Friedreich's ataxia he thinks the progress of the disease may be explained as the legs and arms are used more and more with the progress of years. There is some abnormal weakness, either congenital or precipitated by an acute disease. The posterior and lateral tracts which are most used thus become affected. Here, however, Edinger admits there are serious objections to the adequacy of his theory. Why are other parts of the nervous system exempted through life? Why do the pupils, the bladder and the brain not show more disturbance than they do? No answer is attempted, but the very elementary nature of our knowledge of the physiology of certain parts of the nervous system is emphasised.

Dr. C. Macalister (*British Medical Journal*, April 8, 1893) suggests that the tendency to degenerate in certain diseases, such as pseudo-hypertrophic paralysis, may be due to the diminution in the functional activity of some gland whose secretion exercises a controlling influence over the nerve elements. Stawell⁶⁷ would extend the theory to cover cases of Friedreich's ataxia.

It will be evident that the above theories, however suggestive, do not by any means suffice to explain the whole disease in any large and general way. This symptom or that may be elucidated through the aid of other affections, perhaps more thoroughly understood. Atrophy of muscles, for instance, as of the interossei, may be referred to degenerations of their trophic cells in the anterior horns. Atrophies in this disease are not so extensive or frequent as to lead one to expect any very striking evidence of wasting of ganglion cells, but such wasting has been discovered.

The *ataxia* seems to be different from that of tabes, both clinically and pathologically. The cerebellar character may be referred, if not to the cerebellum, at least to the tracts conveying impulses to the cerebellum. The direct cerebellar

and the antero-lateral ascending tracts are longitudinal commissures between successive spinal segments and some part of the brain (Foster, "Physiology," par. 580). Another suggestion is that the posterior columns are the main tracts for conveying motor impulses downwards (Inglis, *Journal of Nervous and Mental Diseases*, January, 1892). Hughlings Jackson's view that cerebellar ataxia is due to weakness of the back muscles, and to the consequent attempt of the legs to correct the weakness of the back, is strengthened by such cases as those we have been considering. (For valuable discussion of this subject see paper by Ormerod, *British Medical Journal*, 1894, vol. ii., p. 1291.)

But the ataxia is not only cerebellar, it is tabetic also. The muscle-sense to whose impairment the most widely accepted theory refers the ataxia of tabes, is usually, though as we have seen not invariably, well preserved in Friedreich's disease. With a pathology so alike, as regards the posterior columns, one is tempted to think that whatever explanation is adequate for the ataxia and absence of jerks in tabes is likely to be sufficient *pro tanto* in the case of Friedreich's disease.

THERAPEUSIS.

Almost all writers on this part of the subject speak in the most hopeless tone. However, Zabłudowski in his patient, a little girl of 9, obtained decided improvement by a four weeks' course of massage. The case was an isolated one, and was exceptional in some minor points; for instance, regurgitation of fluids through the nose occurred in swallowing, formerly solids also had been thus returned. The "cure" so improved her that she seldom fell, walked straight, and could feed herself without letting the spoon drop. Massage increases the functional capacity not only of healthy muscles but of those whose activity is impaired by direct or indirect causes. The nerves are better nourished through the improved circulation. Nerves and muscles only partially diseased may take up the work of those that are further gone. To this one might add Weigert's concep-

tion of the mode of action of a stimulus. "Neither cell nor organ is directly strengthened by the stimulus; the apparent strengthening is due to the fact that in action portions are used up and replaced by new fresh portions, a substitution which in many cases occurs in excess." (Edinger.⁸⁰) Improvement thus obtained can hardly be anything but temporary. Case IV. had a month's massage in hospital but without benefit. It is clear, however, that rest is the worst thing possible for such cases; they should be encouraged to take exercise, and to go through such actions as will best bring their co-ordinating faculty into use. Touching definite points, picking up small objects and placing them on marked spots, writing, walking without or with aid—these and other exercises employed in the mechano-therapeusis of tabes should be useful in Friedreich's ataxia.

LITERATURE.

- (1) LADAME, "Friedreich's Disease." BRAIN, vol. xiii., p. 467, 1890, a critical digest, with full bibliography to 1889.
- (2) DÉJÉRINE ET LETULLE. "Etudes sur la maladie de Friedreich; sclérose nevrologique pure des cordons postérieurs." *La Méd. Moderne*, vol. i. No. 17. 1890.
- (3) LETULLE ET VAQUEZ. "Un cas de maladie de Friedreich avec autopsie." *Comptes rend. de la Soc. de Biol.* p. 21, 1890.
- (4) DÉJÉRINE. "Sur une forme particulière de mal. de Friedreich avec atrophie musculaire et troubles de la sensibilité." *La Méd. Moderne*, 1890. No. 25. Two excellent cases with paralysis of hands and feet.
- (5) AUSCHER. "Anatomie path. de la mal. de Friedreich." *Sem. Médicale*, No. 32, July 30, 1890 (with remarks at the Société de Biologie, by Déjerine).
- (5a) BLOCC ET MARINESCO. "Sur l'anatomie pathologique de la maladie de Friedreich." *Archives de Neurologie*, pp. 118 and 331, 1890. Case of a girl, ill from age of 9, died at 19.
- (6) DÉJÉRINE ET LETULLE. "Sur la nature de la Sclérose dans la mal. de Friedreich." *Semaine Méd.*, March 12, 1890.
- (7) ROOK. "Four cases of Friedreich's disease." *Journal of Nerv. and Ment. Dis.*, p. 173, 1890. Claw-like hands in two. Autopsy on one, by Dana, 1896; see below.
- (8) BROWN, C. H., of New York. Two cases, not related. *Journal of Nerv. and Ment. Dis.*, 1890 (abstract in *Neurol. Centralblatt*, 1892, p. 209). Both cases exceptional, doubtful.

- (9) ERB. "Two cases of Friedreich's ataxia." *Deutsche Med. Woch.*, No. 46. 1898. Short account, knee-jerks.
- (10) MENDEL. "Friedreich's ataxia." Two brothers and a male cousin. *Deut. Med. Woch.*, No. 47, 1890.
- (11) EWART. *Lancet*, vol. i., p. 991, 1891. "Case of Friedreich's disease, with increased knee-jerk."
- (12) BRAMWELL. *Atlas*, vol. i., p. 28, 1891-2. Case of Friedreich's disease, with essay on whole subject.
- (13) BRAMWELL. *Atlas*, vol. i., p. 45, 1891-2. "Three cases of Friedreich's ataxia." Subsequent history up to death of two of those given by Herbert Bramwell, in *Brit. Med. Journ.*, October 2, 1897. Also a note on the histological characters in the cord of one of them, by Byrom Bramwell. The main interest of this is a confirmation of Déjerine's description of the wheel-like arrangement of the neuroglial fibres in the posterior columns. Otherwise the appearances were such as are commonly found.
- (14) BRAMWELL. *Atlas*, vol. i., p. 178, 1891-2. "Notes of three new cases of Friedreich's ataxia."
- (15) LYMAN. "Juvenile or hereditary ataxia." *Intern. Clinics*, 1891 (knee-jerks increased).
- (16) BAUMEL. "Considérations cliniques sur la maladie de Friedreich." *Nouv. Mercredi Méd.*, vol. ii., p. 49, 1891.
- (17) DANA. *Keating's Cyclop. of Dis. of Children*, vol. iv., pp. 716-725, 1891. Article, with one doubtful case.
- (18) FRIEDENREICH. "Et tilfalde af hereditær ataxi. Friedreich's sygdom." *Hosp. Tid.*, 1891 (*Neurol. Centralblatt*, 1892, p. 211). One case, second in family, from birth, never steady on legs. Optic atrophy. Intelligence defective.
- (19) DESTREE. "Un cas de maladie de Friedreich." *Journal de Méd. de Bruxelles*, Nov. 5, 1891. One case with fulgurant pains, strabismus, and slight left ptosis.
- (20) INGLIS. "Friedreich's ataxia, its relation to conducting paths in cord." *Jour. of Nerv. and Ment. Dis.*, vol. xix., p. 30, 1892. An isolated case, no nystagmus.
- (21) SANGER BROWN. *BRAIN*, pp. 250-282, 1892. Twenty-one cases in five generations of a hereditary ataxia, not Friedreich's; cerebellar gait, increased jerks, visual troubles, affection of deglutition, late onset.
- (22) BRAMWELL. *Atlas*, vol. ii., p. 22, 1892. "Two cases of Friedreich's disease in which the knee-jerks were not lost."
- (23) BRAMWELL. *Atlas*, vol. ii., p. 70, 1892. "Three cases of Friedreich's disease with *main en griffe*."
- (24) HALLION. "Des déviations vertébrales névropathiques." *Nouv. Iconog. de la Salpêtrière*, 1892.
- (25) LEEGAARD. "Friedreich's sygdom (hereditær ataxi)." *Norsk. mag. f. Lægevidensk.*, 1892, s. 444 (ref. *Neurol. Centr.*, p. 618, 1893). Three cases, acute onset, with fever and pains in limbs; spastic symptoms, with pains in limbs supervening on the ataxia.

- (26) MARIE. "Leçons sur les maladies de la moelle," 1892.
- (27) HILDEBRANDT. "Über tabes dors. in den Kinderjahren." Berlin, 1892 (abstr. in *Neurol. Centr.*).
- (28) GEIGEL. "Über Friedreich's hered. ataxie." *Sitzungsber. d. Würtsb. Phys. Med. Gesellsch.*, 1892 (*Neurol. Centr.*, p. 50, 1892). Two cases.
- (29) BURGESS. *Brit. Med. Jour.*, April 9, 1892. "Two cases of Friedreich's disease."
- (30) AUSCHER. "Sur un cas de maladie de Friedreich avec autopsie." *Arch. de Phys. Norm. et Path.*, April, 1893, p. 340 (see also No. 4, same case).
- (31) M'Caw. "Notes of a case of Friedreich's disease in a girl 8 years old." *Lancet*, vol. ii., p. 484, 1893. An isolated case.
- (32) KRAUSE. "Et tilfælde af Friedreich's sygdom." *Hosp. Tid.*, 1893, 4r., I., 30. Isolated case, classic sympt. (*Neurol. Centr.*)
- (33) CHAUFFARD. "Maladie de Friedreich avec attitude athetoïdes." *Sem. Med.*, vol. xiii., p. 409, 1893. Isolated case, marked jerky movements on exertion.
- (34) SENATOR. "Über hered. ataxie." *Berl. Klin. Woch.*, vol. xxx., p. 489, 1893.
- (35) SENATOR. Second article. *Berl. Klin. Woch.*, vol. xxxi., No. 28, 1894. Two cases in one family in those two papers.
- (36) ANDERSON, WALLACE. "Friedreich's ataxia." *Glas. Med. Jour.*, vol. xl., No. 8, 1893. Two cases, retained knee-jerks, mental impairment, slight athetosis.
- (37) MIRTO. "Atassia di Friedreich e atassia volgari." *Giorn. Del. Ass. Dei Medici e Naturalisti*, Anno iv., puntati 3 e 4, 1893.
- (38) TRESIDDER. "Three cases of hereditary ataxia." *Lancet*, vol. ii., p. 304, 1893.
- (39) WILSON. "Notes on Friedreich's disease." *Birm. Med. Rev.*, vol. xxxi., October, 1893. Four cases, two isolated, all somewhat atypical.
- (40) JAMES. "Lecture on a case of Friedreich's ataxia." *Edin. Med. Jour.*, vol. xxxiv., p. 521, 1893. An isolated case; spinal curvature first symptom.
- (41) BROCK. "Three cases of Friedreich's ataxia." *Lancet*, vol. i., p. 139, 1893.
- (42) FRY. "Two cases of Friedreich's ataxia." *Med. Fortnightly*, St. Louis, December 15, 1893. Typical; sisters.
- (43) BENTHALL. "A case of Friedreich's ataxia." *Brit. Med. Jour.*, vol. i., p. 1,163, 1893.
- (44) LUNZ. "Ein Fall von Friedreich'scher Krankheit." *Deut. Med. Woch.*, August 17, 1893. A typical case with no nystagmus.
- (45) RAUZIER. "De la maladie de Friedreich." *Nouv. Montpellier Méd.*, vol. ii., pp. 725-748, 1893.
- (46) MARIE. "Sur l'héréd-ataxie cérébelleuse." *Sem. Med.*, No. 56, 1893.
- (47) BRISSAUD ET LONDE. *Rev. Neurol.*, 1894. One case of hereditary cerebellar ataxia.

- (48) MACKENZIE, H. W. G. "A case of non-hereditary Friedreich's disease." *Amer. Jour. of Med. Sci.*, vol. cvii., p. 371, 1894.
- (49) GUIZZETTI. "Contributo illa anatomia patologica della malattia di Friedreich." *Il policlinico*, p. 438, 1894.
- (50) DRESCHFELD. *Lancet*, vol. i., p. 1,014, 1894. A short account of two isolated cases.
- (51) NAMMACK. *Archiv. of Pediatr.*, New York, vol. xi., p. 386, 1894. An isolated case.
- (52) MACKAY. "The isolated type of Friedreich's disease." *Amer. Jour. of Med. Sci.*, vol. viii., p. 151, 1894.
- (53) SCHULTZE, FR. "Über die Fried. Krank. und ähnliche Krankheitsformen." *Deut. Zeitschr. für Nervenh.*, vol. v., 1894. Heft 1-3, pp. 27 and 194. A criticism on Senator's case (No. 31 above), a re-examination of the cord of one of Friedreich's cases, with criticism of Déjerine and Letulle's observations, and a description of three new cases, which he calls a variety of Friedreich's disease (*ähnlich*).
- (54) BESOLD. "Klinische Beiträge zur Kenntniss der Friedreich'schen Krankheit." *Deut. Zeit. für Nerv.*, vol. v., p. 157, 1894. Four isolated cases.
- (55) TAYLOR, JAMES. *Pract.*, November, 1894. Two cases, one with marked family tendency; the other isolated, with slight mental degeneration, and retained, though weak, knee-jerks.
- (56) KRAFFT-EBING. "Über Fried. Krankheit." *Allg. Wiener Med. Z.*, No. 14, 1894. One case.
- (57) MICHELL CLARKE. "Case of Friedreich's disease, with necropsy." *Brit. Med. Jour.*, December 8, 1894.
- (58) SCHULTZE. "Erwiderung auf den zweiten Artikel von Senator." *Berl. Klin. Woch.*, No. 33, 1894.
- (59) TIRASEK. "Česopis lékařů českých." Praha. *Sajous' Annual*, 1894. One case of Friedreich's disease in a family with frequent inter-marriages.
- (60) BOUCHAUD. *Rev. Neurologique*, No. 1, 1894. (Ref. in Higier's article). Two cases allied to diplegia and Friedreich's disease.
- (61) LONDE. "De l'héréd-ataxie cérébelleuse." *Thèse de Paris*, 1895.
- (62) GUTHRIE. "Case of early Friedreich's ataxia." *Brit. Med. Jour.*, vol. i., p. 813, 1895. (Also in *Trans. Med. Soc., Lond.*, xviii., p. 382, 1894-5.) A boy of 9; no family predisposition, no nystagmus or speech defect.
- (63) NOLAN. "Three cases of Friedreich's ataxia, associated with genetous idioey." *Dublin Jour. of Med. Sci.*, xcix., p. 369, 1895.
- (64) SMALL, S. J. "Friedreich's ataxia, with cases." *Med. Rec.*, New York, xlviii., p. 85, 1895. Four advanced typical cases in a family, with strong neuropathic tendency. Onset at 13 or 14.
- (65) KÖNIG. "Cerebrale Diplegie der Kinder, Friedreich'sche Krankheit und multiple Sklerose." *Berl. Klin. Woch.*, xxxii., p. 716, 1895. A case like Friedreich's ataxia and disseminated sclerosis in some respects; also another case of hypoplasia of cerebellum and cord without inco-ordination.

- (66) BOUCHAUD. "Maladie de Friedreich. Amblyopie, persistance du reflexe rotulien." *Jour. d. Sci. Med. de Lille*, No. 14, p. 1,313, 1895. The parents were first cousins. An isolated case.
- (67) STAWELL, R. K. "Two cases of Friedreich's disease." *Austral. Med. Jour.*, N. S., xvii., p. 45, 1895. Brothers, typical cases.
- (68) HALBERT. "Two cases of Friedreich's ataxia." *Clinique, Chicago*, No. 16, p. 636, 1895. Congenital syphilis, imbecility, spastic symptoms. Not examples of Friedreich's disease.
- (69) CARDOSI, S. "Contributo alla forme rudimentale del morbo di Friedreich." *Gazz. d. Osp. Milano*, No. 16, p. 158, 1895.
- (70) VERHOOGEN. *Jour. de Med. de Chir. et de Pharm.*, Bruxelles, January 19, and March 9, 1895. Two typical cases of Friedreich's disease.
- (71) HOFFMANN. *Münch. Med. Woch.*, February 19, 1895. Two cases, one typically Friedreich's disease, the other doubtful. Epilepsy and mental weakness in this. Both isolated.
- (72) LONDE ET LAGRANGE. *Annales de Med.*, Paris, March 7, 1895. "Two cases of Friedreich's ataxia." Sisters; athetoid attitude in walking, mental defect.
- (73) COLLINS. *New York Med. Rec.*, December 21, 1895. "A contribution to the study of hereditary cerebellar and hereditary spinal ataxia, with the history of an atypical case of the first-named disease."
- (74) BURB. "A contribution to the pathology of Friedreich's ataxia." *Univ. Med. Mag.*, Philadelphia (*Neurol. Centr.*, p. 376, 1895).
- (75) MACKENZIE, H. W. G. "Friedreich's disease." Article in *Med. Annual*, 1895. A short but valuable descriptive account, with a list of references.
- (76) BASKETT. "A case of Friedreich's disease." *Brit. Med. Jour.*, vol. ii., p. 749, 1896. No family affection.
- (77) DANA, C. L. "A case of Friedreich's ataxia, with autopsy." *Post-graduate*, New York, vol. xi., p. 328, 1896.
- (78) HIGIER. "Über die seltene Formen d. hered. u. fam. Hirn und Rückenmarkskrankheiten." *Zeitschr. f. Nervenheilkunde*, vol. ix., pp. 1-61, 1896.
- (79) BERDEZ. "Un cas isolé de mal. de Friedreich." *Rev. Med. de la Suisse rom.* (Genève), vol. xvi., p. 304, 1896.
- (80) FRY, F. R. "Additional cases of Friedreich's ataxia." *Med. Rev.*, St. Louis, vol. xxxiv., p. 19, 1869. One an isolated case, the others sisters; of these one is very doubtful, the main feature being post-diphtheritic paralysis.
- (81) ORMEROD. "On Friedreich's ataxia." *Clin. Journ.*, London, vol. viii., p. 118, 1896-7.
- (82) RIGGS, C. E. "Two cases of Friedreich's ataxia." *North Western Lancet*, St. Paul, vol. xvi., p. 264, 1896. Cases not in same family, description very brief; one probably not correctly diagnosed (spastic gait, ankle and rectus clonus, but no knee-jerk).
- (83) ROSENBAUM. "Friedreich'sche Krankheit." *Deut. Med. Woch.*, vol. xxii., p. 483, 1896. Two cases of typical form; sisters.

- (84) ZABLUDOWSKI. "Ein Fall von Friedreich'scher Krankheit. Behandlung durch Massage." *Berl. Klin. Woch.*, p. 757, 1896.
- (85) TEDESCHI. "Die Friedreich'sche Krankheit, kritische und pathanat. Untersuchungen." *Beiträge z. Path. Anat., etc.*, vol. xx., pp. 51-84, Jena, 1896. A useful summary of autopsies. A careful pathological account of the case criticised above, and a bibliography.
- (86) MAGNUS-LEVY. *Deut. Med. Woch.*, No. 28, 1896. *Ver. Beil.*, No. 19. Isolated case, doubtful. Details scanty.
- (87) LEYDEN UND GOLDSCHIEDER. "Erkrankungen des Rückenmarkes." Wien, 1895-6.
- (88) GOWERS. "Diseases of the nerves and spinal cord." London, 1892.
- (89) VINAJ. "Malattia di Friedreich." *Osservazione Clinica. Estratto dalla Gassetta Medica di Torino*, No. 61, 1894.
- (90) EDINGER. "Eine neue Theorie über die Ursachen einiger Nervenkrankheiten, insbesondere der Neuritis, und der Tabes." *Samml. Klin. Vortr.*, No. 106.
- (91) RAYMOND. "Tabes juvenile et tabes héréditaire." *Le Progrès méd.*, 1897. No. 82 and 83.

A Case of Friedreich's Ataxia, with Autopsy and Histological Examination. By JULES SIMON. (*Le Progrès Méd.*, 1897, September 4, No. 36.)

JULES SIMON relates the following case:—Boy, aged 10, only child, admitted to Enfants-Malades in 1889 for pains in legs. Nothing hereditary of importance. Had pains in legs at 2½. dysentery at 5, after which he walked badly. In 1891, following was observed: strumous aspect, chronic blepharo-conjunctivitis, old keratitis, not intelligent-looking, speech slow, jerky. No nystagmus, club-foot, scoliosis, sensory troubles or sphincter affection. Knee-jerks absent. Standing difficult, especially if eyes shut. "Drunken" gait, often falls. Diagnosis of Friedreich's ataxia, confirmed by Joffroy. In 1896, slight tremor in head, jerky and fibrillary movements in hands; motion and sensation normal in upper extremities. Lower limbs cannot support him. Atrophy of calves and muscles of soles, club-foot, no knee-jerks, normal sensibility, cold feet. Pupils normal, transverse nystagmus on lateral fixation. Tends to slaver, especially on laughing, slight deviation of mouth to right. Cannot whistle now but could formerly. No scoliosis. The pains he has from time to time consist in intense heat and pricking in calf regions. Intelligence mediocre. Often has headaches followed by vomiting preceded by giddiness. Incontinence of urine frequently.

Death sudden and unexpected.

Post-mortem.—Normal naked eye appearance of cerebrum and cerebellum; weight of latter normal. No histological examination (unfortunately *Rep.*).

Examination of cord by Philippe (three months in Müller's fluid, Weigert-Pal stain).—I. Posterior columns show loss of myeline sheaths. (1) Sacral region, level of third sacral root: rootzones more affected than the zones of endogenous fibres, the short commissural fibres; myelinic reticulum of posterior horns very poor. (2) Lumbar region: rootzones specially affected, also the centrum ovale of Flechsig; cornu-commissural zone nearly intact; myelinic reticulum of posterior horns affected as in sacral region. (3) Dorsal region: decoloration of posterior columns at maximum in upper dorsal region; spec. the lumbo-sacral fibres, near the posterior fissure; the myelinic reticulum in Clarke's column scanty. (4) Cervical region (sixth root): Goll's column nearly completely decolorized; the postero-external column also well decolorized but less so than in dorsal region. (5) Bulbar region: fasciculi gracilis and cuneatus much decolorized, the former more, representing as it does Goll's column.

II. Posterior roots, loss of myeline everywhere, more so where the adjacent posterior columns are most affected.

III. Lateral columns, affected all along cord, but less so than posterior column; pyramidal tract and cerebellar tract affected; direct pyramidal tract in upper dorsal and cervical regions.

Where the nerve tubes have gone, their place is taken by a dense sclerosed neuroglia, arranged in places in whorls (both in posterior and lateral columns); connective tissue thickening of pia mater over posterior aspect; considerable diminution in number and volume of cells in posterior horn and at base of anterior horn; also of column of Clarke in lumbar and dorsal regions.

Conclusion.—The cord is the seat of a combined sclerosis affecting the posterior columns, the pyramidal bundle and the direct cerebellar tract. This corresponds fairly well with the lesions hitherto found in cases of Friedreich's disease. In addition to the combined sclerosis of the white matter, there is a lesion of the gray substance.