he is unable to answer. It is, he said, very difficult to speak concretely on the subject of seborrheids, as we have not a very clear comprehension of what seborrheids really are, nor are we in accord as to the etiologic factors that have been described in connection with seborrheic processes. Some years ago he examined the skin of fifty patients who came to the dermatologic clinic, for various reasons, and in 45 per cent. of these cases, the sebaceous glands contained the microbacillus of Sabouraud. Whether this organism bears any special pathogenic relationship to the class of eruptions under discussion future investigation must determine. It has not been positively proved that the microbacillus is not a normal inhabitant of the sebaceous glands, although Sabouraud certainly makes out a strong case for its pathogenicity in relation to oily seborrhea.

Dr. W. H. Mook, St. Louis, said, in regard to the pityriasis affections of the face and neck which are embraced under the second heading, he would insist that this is certainly a distinct condition from eczema, and should be so regarded. The idea of his classification of this form of acne is to distinguish it from the other forms, as it is, in his opinion, a distinct entity.

THE TECHNIC OF OPERATIONS FOR CARCINOMA OF THE BREAST.*

MILES F. PORTER, A.M., M.D.

Surgeon to Hope Hospital, Professor of Surgery in the Indiana Medical College, Department of Medicine, Purdue University.
FORT WAYNE, IND.

The chief object of operation for cancer of the breast is, of course, the cure of the cancer. This is as it should be, but it is also important that the convalescence of the patient be as comfortable and as rapid as possible, and that the resulting disability be as slight as consistent with radical removal of the growth. It is especially with a view of aiding somewhat in the accomplishment of the last-named objects that this paper is written. The chief basis of the paper—aside from a study of the literature—is my personal experience in this line of work, something over fifty cases of radical removal.

SUPERFICIAL INCISION.

This must be varied often to suit the case in hand, but the skin forming the floor of the axillary space should not be incised except in cases wherein the removal of a portion of it is necessary. There are two reasons why it is desirable to avoid incising the skin in this area: First, because infection is more likely to follow incisions made in this area, and, second, because the resulting scar is likely to interfere with the movement of the arm.

Figure 1 illustrates these points. There was a slight infection of the wound in the apex which increased the amount of scar tissue and exaggerated the bridge which interferes with the upward movement of the arm. Incisions so made as to form flaps with acute angles are objectionable, because they are difficult to coapt and are prone to slough on comparatively slight provocation. Rectangular incisions for removal of the breast, as compared with circular or elliptical incisions, are wasteful of tissue. The incision which I think the best, and one which is adapted to most cases, is one commencing at the humeral insertion of the pectoralis major muscle and following its lower border until the breast is reached, then surrounding the gland in a somewhat elliptical fashion so that when coapted the line of coaptation will occupy about the position occupied by the lower border of the normal pectoral muscle, save that

it extends farther inward and downward. This incision gives ample room for the clearing of the axilla and the removal of the pectorals, and leaves a minimum scar, which may be fixed well up in front of the joint, where it will be least likely to interfere with abduction of the arm.

Figure 2 shows the result after an incision of the kind described. As stated above, the lines of the incision must of necessity be changed sometimes to meet the requirements of the case at hand, but the principle of so planning the incisions as to allow of coaptation with a minimum amount of tension made in a vertical direction, especially in the outer aspect of the wound, must be adhered to as nearly as is consistent with complete removal, if the best results are to be attained, so far as concerns the movement of the shoulder.

PART TO BE REMOVED AND METHOD OF REMOVAL.

All of the pectoralis major muscle, except the clavicular portion, and the abdominal portion when present, should be removed, together with the whole of the pectoralis minor, the axillary fat and glands, and the breast, with its coverings.

The pectoral insertions are most quickly and safely divided by scissors, using the finger as a guide. The cleaning of the axillary space should be from above

Fig. 1.—Showing scar tissue in axillary space interfering with upward movement of arm.

downward and from without inward. The larger part of this dissection can be most satisfactorily done by wiping the fat and gland away from the vessels. I owe my knowledge of this gauze dissection to Dr. A. J. Ochsner, and regard it as an important improvement in the technic. By ligating the vessels near the main trunks, as advised by Jackson, the field may be kept free from blood and the work much expedited. Clamps are placed, when necessary, on the distal ends of the vessels. As the mammary gland is lifted from the chest wall the perforating arteries may be caught with forceps before they are divided.

CLOSING AND DRESSING THE WOUND.

The rapidity and completeness of the patient's recovery, as well as her comfort, depends, in a large measure, on the way the wound is closed and dressed. The most perfect wound treatment is that which reduces to the minimum the scar tissue, and disposes that which is unavoidable in such manner as not to interfere with function. With this object in view the wound should be completely closed and no drainage used. In a very few cases, indeed, will it be impossible to secure com-

* Read in the Section on Surgery and Anatomy of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

plete coaptation of the margins of the wound, and in still fewer will drainage be necessary.

Hemostasis must be practically perfect. In accomplishing this, the use of hot compresses is invaluable. These are applied to oozing surfaces while the larger vessels are being tied or twisted and the wound is being closed. Not only does the heat assist in preventing the bleeding, but it is of material value in preventing the untoward results, pneumonia, shock, etc., which may follow the exposure of so large a raw surface to cold. I have already said that in all but a very few cases complete coaptation of the margins of the wound can be secured. In six patients operated on in Hope Hospital since October, 1905, complete closure of the wound has been secured in all. In order to do this, the skin must be drawn taut over a larger or smaller area, but the skin will stand much more tension without breaking down than most operators seem to think. In most of these cases in which the wound is completely closed, the skin will

Fig. 2.—Showing scar following incision recommended by author.

have to be stretched until it is white, but by placing tension sutures properly, and in sufficient number so as to distribute the tension as evenly as possible, no harm will come from the tension, and the wound will heal by first intention within a week. For tension sutures I prefer silk-worm gut and buttons. The buttons have a tendency to produce pressure necrosis if the tension is great, but this can be so nearly overcome, by placing under the buttons a little gauze, that it amounts to but very slight annoyance at most.

Figure 3 is from a photograph, taken before operation, of a case of recurrent adenocarcinoma of the left breast. One can see from the picture how extensive a removal of the skin must have been necessary in this case, and yet, following the method described, complete closure of the wound was secured and the wound was

found to be completely healed when the dressing was changed on the seventh day after the operation.

Figure 4 is from a photograph of this patient taken seven days after operation; the wound is healed, and the hand can be placed on the head. Equally good results, as to wound closure and movement of the arm, were obtained in the case of a Mrs. H., in which, in addition to the removal of the tissues ordinarily removed, the floor of the axillary space, the fascia of the serratus magnus, and subscapularis, and the edge of the latissimus dorsi were removed, together with the supraclavicular glands. These latter were removed through an incision above the clavicle.

With the tension sutures properly placed, the edges of the wound can be brought together without marginal tension. In coaptation, adhesive straps are of much value. Indeed, the coaptation of the margins of the wound is, for the most part, accomplished by adhesive straps, aided not infrequently by subcuticular sutures of catgut, and, in rare cases, a few through-and-through sutures are used.

Of no less importance than the coaptation of the margins of the wound is the coaptation of the skin to the underlying surface. If this part of the coaptation is perfect, there will be little likelihood of subsequent

Fig. 3.—Recurrent adenocarcinoma before operation, showing how much tissue it was necessary to remove.

trouble from lack of motion in the shoulder or from venous stasis, except, of course, the stasis due to compression of the vein by recurrence of the carcinoma. Lymphatic edema will occur in a small percentage of cases as a direct result of the operation, owing to the cutting off of the communication between the lymph vessels of the arm and those of the neck, and, under these circumstances, is rather an indication of a complete operation than otherwise. Edema of this kind usually manifests itself soon after the operation, and is usually at most a source of some inconvenience. The point is that we have removed the anterior wall, together with the cellular and glandular contents of the axilla, and to secure subsequent freedom of movement of the shoulder nothing more nor less is necessary than to apply the skin flaps accurately to the remaining walls and secure their union without infection. Of especial importance is it to secure coaptation of the skin and underlying cellular tissue to the apex of the axilla. Failing in this, there will be formed a smaller or larger wedge of connective tissue in the apex of the axilla, closely adherent on all sides, or a bridle will be formed

at some distance below the joint and firmly attached to both inner and outer axillary walls. The natural result of either condition will be inability to raise the arm and more or less interference with the return of blood from the arm. To secure the opposition necessary to prevent those untoward results, I relied on a snug application of the external dressings until something less than a year ago, but it was found, not infrequently, that they were too tight for comfort and produced considerable swelling of the arm, forearm and hand. To overcome this difficulty, I conceived the idea of using subcuticular stitches of catgut to secure the apposition necessary, and have used them in six cases with perfectly satisfactory results in all save one, which is shown in Figure 1. In this case, the incision is faulty, but, notwithstanding this, a proper placement of the stitches would have prevented the inhibition of movement which is present. (The arm can not be raised higher than it appears in the picture.) The line of incision in the skin should

Fig. 4.—Recurrent adenocarcinoma seven days after operation, showing degree of freedom of movement in arm.

have been fixed well up in front of the joint. Two or three stitches are usually sufficient.

I described this method to several of my friends and requested them to try it, and I would be glad to hear of their results. The advantages of this technical device over others, whose object is the same, are:

1. The certainty of the good results.
2. The comfort it secures to the patient.
3. Its simplicity and ease of execution, and
4. More rapid recovery of usefulness of the arm.

Figure 5 is from a photograph taken fourteen days after operation. The wound area in these cases is so large that with the most perfect hemostasis practicable there will be enough oozing to cause considerable accumulation of blood under the flaps while the margins are being coapted, unless measures are taken to prevent it. The use of hot compresses has already been referred

to, but I wish here to speak of their use more particularly. As soon as the axillary space is cleared, a hot compress is placed in it and held snugly by an assistant while the removal of the parts is being completed, and the same thing is done with the remaining wound area as soon as it is cleared. The aim is to keep these compresses applied as continuously and as long as is consistent with the reasonably rapid completion of the work. The one in the axilla is not disturbed until one is ready to place the sutures that anchor the skin in the apex, and only just enough to permit this to be done. The margins are sutured over the compress until its lower margin is reached when it is withdrawn. During the suturing and after the withdrawal of the compress, an assistant is directed to keep the flaps closely applied to the underlying compress and chest wall, respectively. The compresses in the lower and inner aspect of the wound are removed for the placing of the tension sutures. Before the last stitches are tightened, the flaps are methodically compressed toward the opening to remove any blood that may have accumulated; the stitches are tightened, the adhesive straps placed, and the dressing put on. Care should be taken to apply this so that it will hold the raw surfaces snugly in apposition. The arm is carried in a sling for a week, when the patient is allowed to use it with caution for

Fig. 5.—Showing result of using subcuticular stitches of catgut. Fourteen days after operation. Hand can be raised to head.

another week, after which she is allowed to use it as she pleases.

The first dressing is allowed to remain undisturbed for a week, when it is taken off and another applied for a week longer, when no further dressing is needed unless it be a few layers of gauze to protect the tender surfaces. The patient is put to bed, with the head and trunk well raised, and is allowed to assume the full sitting posture as soon as she will. Patients are usually walking about by or before the end of the week.

CONCLUSIONS.

In conclusion, permit me to emphasize the following points:

1. Complete closure of the wound after removal of cancer of the breast may be secured and drainage dispensed with in nearly all cases if proper attention be paid to the placing of the incisions, the securing of hemostasis and the use of tension sutures.

2. Venous stasis in the arm and lack of motion in the shoulder, as postoperative results, may be avoided in nearly all cases by avoiding infection and keeping the wound surfaces in close apposition until healing is complete.

3. The best means of accomplishing the last-named object is a few buried sutures in the apex of the axilla and a properly adjusted compressive dressing held in place by adhesive plaster.

DISCUSSION.

DR. WILLIAM L. RODMAN, Philadelphia, agreed with Dr. Porter as to the necessity of the free removal of the overlying skin, even in such cases as show that we can not get primary union. He inclined to the belief that possibly the greatest mistake that is made in practice generally is to subordinate the probabilities of a radical cure to securing primary union in every case. He said that we should keep constantly before us that Gross and Pancoast and the older surgeons reached 10 and 15 per cent. of cures, but only by very freely removing the skin. A complete axillary dissection was never done, and muscles were not destroyed. Whatever good was in these operations must have been in the free removal of skin. Therefore, he thinks that we should not accentuate the necessity for getting primary union, even in a majority of the cases. It is all very well if it can be done and without making too much tension, and this, he said, can be done, as Dr. Porter says, by undermining the skin, or by skin grafting, which can always be resorted to to cover in any defect that may exist.

Dr. Rodman has done the operation according to the method described by Dr. Porter a number of times, and he has also advocated strongly the operation of Warren, which is practically the same. He liked it very much and thought it the best operation he knew, inasmuch as it has generally made primary union possible, but he has found in a number of cases a bridle of skin which fetters and prevents the uplifting of the arm beyond that extent, thus interfering with the doing up of the back hair, etc. While Dr. Rodman still believes that that operation is one of distinct value, he has obtained recently a better result from the operation of Jackson, and prefers to place the incision in that way. He believes that it is a mistake to cut on to the arm, or, at least, to go low down on the arm.

Dr. Rodman emphasized the necessity of removing the muscles in every case of cancer of the breast, although he was not prepared to say that the muscles will be involved. Researches have demonstrated that the fascia of the muscles usually will stop limitations of the growth, but recent anatomic discoveries show that Halsted built better than he knew when he insisted on the removal of the muscles in every case. For, in the first place, a complete axillary dissection can not be made with the muscles *in situ*, and Gross and Pancoast demonstrated that occasionally there will be lymph glands which can not be felt with the muscles in position. Therefore, it is necessary to remove the muscles in every case, otherwise the operation is incomplete.

As to drainage, Dr. Rodman admitted that in the majority of instances the wound can be closed without drainage, and yet it certainly is the custom of American surgeons to drain after amputations of the breast. In preparing a paper on this subject some time ago, he wrote to fifty American surgeons, and of the twenty-one who replied, there were three who did not use drainage. Certainly drainage is not necessary in case of a small breast particularly, but Dr. Rodman believes that, as a rule, it is just as well to drain.

Speaking of the necessity of invading the neck, Dr. Rodman stated that he is satisfied that we do not invade the neck as often as we should. Recent anatomic investigations show that if the tumor is located superiorly, in the upper hemisphere of the breast, it is drained by the lymphatics which pass from the clavicle into the glands above the clavicle and not directly into the axilla. Therefore, in the case of tumors that are situated superiorly, it is unnecessary to enter the neck in every case.

DR. MILES F. PORTER, Ft. Wayne, Ind., was very glad indeed that Dr. Rodman emphasized the necessity of not sacrificing the chance of a perfect recovery for the sake of securing primary union. In the case he reported the breasts were removed as widely as any one removes them, and yet, in the majority of cases, union was achieved. Dr. Porter invades the neck, when it is the proper thing to do. As to drainage, he said that it is never too late to mend. The sooner those who make it a rule to drain, make it a rule not to drain, the better.

PLASTIC RADIOGRAPHY.

A PRELIMINARY COMMUNICATION.

WALTER G. STERN, M.D., AND E. ROSENBERG, M.D.

CLEVELAND, OHIO.

Radiographs, usually spoken of as skiagraphs, are not, as the latter name would imply, mere shadow-graphs or silhouettes; they are, rather, the result of the superimposition of semishadows or half-tones as they are called in photography, whose size, shape and intensity depend on the size, shape and molecular density of the object to be radiographed. But such pictures have never been interpreted as perspectives or "plastic" reproductions, and successful attempts have never been made to obtain "relief" effects except by means of stereoscopic reproductions.

Fig. 1.—A "plastic" radiogram, showing distinct shadow of soft parts, with high-lights and shadows of the bones, giving relief effects. Print on bromid paper direct from the combination of the negative and positive.

Béla Alexander, a country practitioner hitherto unknown, living in an obscure part of Hungary, first published¹ a method of obtaining "plastic x-ray" photographs. It is through the efforts of one of us (Rosenberg) that the substance of this and other contributions of Alexander and other Hungarian writers in the various Hungarian medical journals has been collected and abstracted. The method as they give it is long and tedious, full of archaisms and omissions of most important details such as the time and methods of exposure, copying, registering, printing, etc. We have tried

1. Orvosegyesület, March 10, 1906.