SECTION II.

ENGINEERING AND ARCHITECTURE.

ADDRESS

BY BALDWIN LATHAM,

M.INST.C.E., F.G.S., F.S.S., F.R.MET.SOC., &c.

PRESIDENT OF THE SECTION.

IN presiding over this section, which is devoted to engineering and architectural subjects, it is necessary that I should say that I have been requested to address you upon a subject which properly belongs to the Climatological Section, namely, upon the probable influence of ground water on health. This request has no doubt been made in consequence of some observations which have been made in this city by Mr. North, the Medical Officer of Health, who has traced some connection between an outbreak of typhoid fever which occurred here in 1884 and the movements of the subsoil water at that time.

Having devoted much time, during the past eleven years, specially to the study of the question of underground water, and having established and maintained a number of stations for observing the relative height of subsoil water in various parts of the country, and having also collected the past records which are available in this country and elsewhere, I am in possession of facts not easily obtainable, and am able, therefore, to draw some definite conclusions as to the probable influence of ground water upon health.

Turning to what may be called historical records, periods of great drought clearly indicate a low state of the ground water, and in ancient records there are some remarkable references to the influence of drought in producing disease.

In the Volume of Sacred Writ, Zechariah, Chap. 14, v. 17 and 18, the effect of the absence of rain in producing plague is clearly set forth, and in a passage in Revelations, Chap. 11, v. 6, men-

tion is made of the with-holding of rain and the production of plague.

Aristotle placed Zoroaster at 6,000 years before Plato, and the latter lived about 360 years before Christ. In the Avesta we have clearly laid down the fact that the absence of rain produces disease.

In these sacred writings it will be observed that the pollution of the earth, water, or air, from any cause, is looked upon as a deadly sin.

The influence of light in destroying noxious properties arising from decomposing matter is also clearly indicated, and it is pointed out that the effect of keeping back the waters of the sky and not suffering them to be poured down on the earth, would be that the noxious animals which live in the water would pollute it.

Again, it is pointed out that the star Tistrya raises the vapours out of the sea and draws the clouds forward, causing the rain to fall. This star was looked up to as the distributor of water, and it is recorded that when the star went to the sea a burning demon came out and scared Tistrya from it. Then the deity of heaven gave strength to Tistrya, and he returned to the sea, and the burning demon came out against him, but was defeated and driven away. Tistrya then raises the vapours of the sea and draws the clouds forward, causing the rain to fall, and so drives away sickness.

No inquiry into the question of the influence of climate on disease would be complete without reference to the labours of Hippocrates. It is curious to note in his works, written upwards of 2,000 years ago, that there are conditions recorded, attending healthy and unhealthy seasons, which are identical with the conditions which may be observed in this country at the present day. Hippocrates taught that all disease may be traced to natural causes, and he counted it impiety to maintain that any one more than another is an infliction of the Deity; and he pointed out to his followers that if they wished to study medicine properly, they must in the first place study the seasons of the year, and the effects which they produce. He also stated that acute diseases occur in periods of drought, and that you could tell what epidemic diseases would attack a city either in summer or winter, and what sickness each individual would be in danger of experiencing. He went much further than our knowledge at the present time enables us to go, for he stated that the changes of the seasons may be predicted from the rising and setting of the stars, so that we could know beforehand what sort of a year would ensue. Hippocrates also pointed out the conditions affecting the health of any period: with rain

in autumn, a mild winter, neither very tepid nor unseasonably cold, and rain in spring and summer, the year is likely to prove healthy; but if the winter is dry, and the spring showery, the summer will necessarily be of a febrile character. If, at the rising of the Dog-star, rains and wintry storms supervene, there is reason to hope that disease will cease, and the autumn will be healthy.

It is curious that a dry winter is often the precursor of disease, not at that time, but in the following autumn. As a rule a short supply of rain in December has a most marked influence upon the stores of underground water, and a deficiency of rain in this month has probably a greater effect in influencing the future health of any particular district than it has in any other month of the year. As a type of a healthy season the present year is an example, and fully complies with the conditions laid down by Hippocrates.

In classic times pestilence was ascribed to the anger of the Sun-God Apollo, or to his twin-sister Diana, the moon, and certainly with very good reason.

“Along Olympus’ heights he passed, his heart
 Burning with wrath; behind his shoulders hung
 His bow and ample quiver; at his back
 Rattled the fateful arrows as he moved.
 Like the night cloud he passed; and from afar
 He bent against the ships, and sped the bolt;
 And fierce and deadly twanged the silver bow.
 First on the mules and dogs, on man the last
 Was poured the arrowy storm; and through the camp
 Constant and numerous blazed the funeral fires.”—*Homer*.

It may be interesting to note that some years ago Dr. Laycock published an interesting account showing the incidence of disease in York, from which it appeared that this city was always susceptible to violent outbreaks of disease, traceable to local sanitary circumstances combined with peculiar climatological conditions, and that there appeared the same incidence in the prevalence of the sweating sickness of 1550-51, the plague of 1604, and the cholera of 1832, to which may be added the typhoid fever of 1884.

The results of my prolonged investigations on the subject of ground water in this country and elsewhere, clearly show that there is generally a direct parallelism between the conditions of health and the volume of ground water. The years in which there has been a large quantity of ground water present have invariably been the healthiest years, while those in which there

has been a small quantity have as invariably been the most unhealthy periods.

As a rule the lowness of the ground water indicates the future health, and not the state of health at the particular time of lowness, that is, the unhealthy period, as a rule, follows the period of low water—the degree of lowness indicating the intensity of future disease, especially fever. Sometimes an unhealthy period runs concurrently with the period of low water, but in all these cases there is clear evidence that percolation has recommenced before the unhealthy period takes place. These results are entirely confirmed by observations which were carried on in Paris between the years 1868 and 1883, and which have been collated and published by M. Durand Claye, Chief Engineer of the Municipality of Paris, with the object of putting all the facts and circumstances in connection with the outbreaks of fever in Paris at the disposal of those who might choose to investigate the subject—a course strongly contrasting with the conduct of some authorities in this country, who desire rather to hide the true facts from public view.

The observations which have been published by Professor Pettenkofer, and which were commenced in 1854, differ from the experience gained in this country, as he has shown that typhoid fever in Munich commenced with the fall of the subsoil water, and reached its greatest intensity with the greatest degree of lowness, and with the rise of the water there was a diminution of fever, a result exactly contrary to experience in this country. Professor Pettenkofer's observations, however, agree with the observations made here in the fact that the greatest intensity of typhoid fever coincides with the periods of the greatest degree of low ground water, that is those years in which the subsoil water has fallen to its lowest level are those in which there has been the most fever.

With regard also to the experience in this country as to subsoil water, it may be pointed out that there is clear evidence that the lowering of the subsoil water by artificial means will produce, and does produce, a tendency to the development and dissemination of typhoid fever. The effect of drainage works, during their construction, in lowering the subsoil water where precautions have not been taken to speedily and permanently get the water back to its proper level, has been, in many instances, to cause outbreaks of typhoid fever, which at the time have been attributed to the sewerage works and to sewer gas, even in cases where no connections had, at the time of the outbreak, been made with the sewers.

It may also be pointed out that at the time of the outbreak of cholera in East London, in 1866, as to the cause of which

there has been so much dispute, the very district which was most afflicted with cholera, had, at the time, its subsoil water unduly lowered by the construction of the main drainage works in that part of the Metropolis, but on the completion of this work, when the sewers were brought into operation, the epidemic ceased.

In regard to outbreaks of cholera in this country, I have not been able to find any record showing the actual state of the subsoil water in 1832, the time of the first outbreak of cholera; but in 1831, there was clearly a deficiency of rainfall, which points to the fact that in the following year the subsoil water would probably be low. In 1847, the year preceding the cholera outbreak of 1848-9, the subsoil water over a large area was undoubtedly low. In the cholera outbreak of 1854, we had a very low period of subsoil water over a large part of the country, and in 1865, immediately preceding the outbreak of cholera in 1866, a similar state of things occurred. With reference to cholera, it should be noted that the conditions of underground water which affect the outbreak, should be observed in those countries where the disease first makes its appearance.

It is curious that in recent times as a rule there has been, every ten years, a marked period of low water; for example, in 1834-5, 1844-5, 1854-5, 1864-5, 1874-5, 1884-5. The lowest water in this series probably occurred in 1864-5. In 1844-5 the low water was not intensely low, but it was low compared with the period. In addition to these periods, there are other times of low water, and in investigating the subject it should be studied locally and comparison made with local vital statistics, for the largely varying distribution of rain tends to equalise results when spread over large areas, as the same conditions rarely occur over extended areas at the same time.

I have carried on, as many of you know, very extensive observations at Croydon on ground water, and in consequence of the results there obtained I have extended them into various geological formations in different parts of the country. I propose now, however, to deal with some of the Croydon results, not that Croydon possesses any exceptional circumstances either with regard to health or disease, but that my intimate knowledge of the district enables me to study all the attendant conditions, and, moreover, in Croydon and its neighbourhood we possess a long record of the condition of the ground water.

The register of Croydon goes back to the year 1539, and with the exception of years in which there has been revolution, or disturbing causes of a kindred character, the record is complete. A tabulation of the whole of the burials and baptisms, extending from this early period to a date overlapping that

when registration of births and deaths commenced, clearly indicates that years of drought are, without exception, the most unhealthy periods.

We possess too, in this country, rainfall records extending back over a period of 200 years, and we have a large amount of concurrent history telling of the state of the weather from very early periods. For example, in 1539, the first year of registration at Croydon, there is a record that in this particular year the springs were remarkably low, so low that the River Lea was nearly dried up, and writers of that age remark on the great drought and heat of that period. In that year the number of burials recorded in Croydon was 50, and the number of baptisms 55, indicating a probable death-rate of 25·6 per thousand. In the following year on the rise of the water, the burials rose to 87, and the baptisms 72, indicating that the death-rate was nearly 32 per thousand. Coming to the period when we have rainfall records, the year 1741 was a very dry time, the rainfall at Lyndon for the year being 15·7 inches, and in that year the burials in Croydon were 271, and the baptisms 113, giving a probable death-rate of 63·7 per thousand, while in the two years preceding this year, the death-rate was 27·7 and 40·7 per thousand, and in the following year 24·2 per thousand. As we arrive at more recent periods when we have the certain records collected by the Registrar-General, registration having commenced in the dry year of 1837, the death-rate at Croydon was 30 in the thousand; in that year and in the following year a similar rate occurred. In the years of very low water 1854 and 1855, we had death-rates of 26·84 and 21·14 respectively, while in 1851, the death-rate was only 18·72 per thousand. In the dry periods 1864 and 1865, we had death-rates of 21·5 and 22·7 per thousand, while in 1860, a wet year and one of high springs, the death-rate was only 17·27 per thousand. The year 1871 was another dry period, but the low water at Croydon was not so intense in this year as in other years. The death-rate was not so high, viz., 18·89 per thousand, but in 1873 when the springs were very high, the death-rate fell to 16·59 per thousand. After the dry period of 1874-5, the death-rate rose again to 21·10 per thousand. From 1876 up to the beginning of 1884, there was unprecedented high water, and during the whole of this period the country generally enjoyed a high state of public health. The conditions, however, which have secured for us the estimable boon of good health, by removing the cause of much sickness, suffering, and death, have been disastrous to the agriculturist, as the large quantity of water passing into the ground has washed away the fertilising elements.

By reference to the returns of the Registrar-General, which

show the deaths in every quarter of the year, it will be seen that, with the single exception of the September quarter of 1849, the period of the cholera epidemic, the most unhealthy periods are invariably those quarters immediately following the periods of low water; as for example, the March quarters of 1838, 1845, 1847, 1853, 1855, 1864, 1865, 1866, 1875, and the increase of deaths in these quarters is so large as to clearly indicate a direct connection between the sanitary influence which the state of underground water denotes, and that of public health. With regard to deaths from fever, in recent times the years in which there has been the lowest subsoil water generally in the country are shown to be the most unhealthy, as for instance the two years 1864, 1865.

The year 1884 was an exceptional year in Croydon, and it is by means of such exceptional periods that we are able to draw some conclusions as to the probable influence of ground water upon health. It is clear to my mind, after the most careful consideration of this subject, that ground water itself has no influence, either for good or evil, upon health, but that the lowness or the highness of the water in the ground is the index of conditions which greatly influence the health of all communities. We have periods of abundance of water, and periods of low water with both healthy and unhealthy conditions. Ground water has been shown by Professor Pettenkofer to be chemically more impure in periods of high water when the conditions were favourable to health than when there is a low state of the ground water, and the conditions are unfavourable to health. The records also show that we have had periods when rain has brought malignant diseases into existence, while, on the other hand, we have similar heavy rainfalls accompanied by a high state of public health, as in the present year.

The records clearly point out that it is not one circumstance alone which produces disease, but that there are at least three factors necessary for the production and distribution of disease, especially typhoid fever, viz. :—

1. The elements which produce disease, such as a polluted state of the ground.
2. The conditions which are necessary for the development of disease, such as a period of dryness of the ground in those regions which are usually occupied by water, combined with a comparatively high degree of temperature.
3. Conditions which will lead to the spread of the disease such as the probable influence of a storm or rain in driving impurities out of the ground into our water

supplies, or through the instrumentality of ground air passing into our habitations, and its reception by a population which is in a condition to receive such germs of disease.

If any one of these conditions is absent, diseases like typhoid do not occur.

The long period of washing and purification which the ground has passed through since 1876, has generally so purified it from the producing elements of typhoid fever that, with the exception of occasional cases of impurity where the ground has become fouled from the leakage of sewage from the imperfect sewers and other direct sources of pollution, as in the case of this city, Beverley, Kidderminster, and some other places which suffered from epidemics in the low water period of 1884, the country has enjoyed comparative immunity from diseases of this class. On the other hand, if we take a period when there has been marked low water for a number of years, followed by unusually low water at particular periods, these are the times when typhoid fever is most rife, as for example between 1854 and 1865, excepting only the years 1860 and 1861, when we had a high state of ground water. The intensity of the fever rates of 1865 and 1866 point out a lesson which ought to be learnt by every sanitarian—that we must keep the ground free from impurity, if we wish to secure conditions which are essential for the promotion of health.

When we come to deal with local conditions preceding disease, we find that not only cholera and typhoid fever, as pointed out by Prof. Pettenkofer, are amenable to the conditions indicated by the highness or lowness of the ground water, but probably all other zymotic diseases, with the exception of diarrhoea, are influenced by the conditions which produce low ground water. At the Leamington Congress of this Institute in 1877, in a paper which I read on this subject, I quoted and reproduced a table from the work of Dr. Macpherson, showing that the incidence of small-pox and cholera was identical at Calcutta, and that both these diseases became virulent at the period of lowest ground water. Small-pox in this country appears to occur when the ground is drying, and is most prevalent in the years of lowest ground water. The conditions affecting measles are almost identical with those of small-pox. Scarlet fever and diphtheria follow almost on the lines of typhoid. They are the least prevalent in the years of highest ground water, and rise and fall with the increase and diminution of percolation. Whooping cough is ordinarily most prevalent at those periods of the year when the ground water is at its highest position, but is also prevalent in years of low ground water. The period of

percolation is that when most of the zymotic diseases run their usual course, that is, they commence with percolation and gradually decline as it ceases. Fifty years of percolation observations at Nash Mills carried on by Messrs. Dickenson, and more recently by Dr. John Evans, F.R.S., when put into tabular form, show percolation to be almost co-incident with zymotic diseases.

We must not, however, lose sight of other conditions which are at work, such as—

(1.) *The influence of light.* It will be observed with reference to the period of percolation that it is almost parallel with the time the sun is below the horizon. The influence of solar light is well-known in malarious countries, which may be traversed with impunity while the sun is above the horizon, but not after night-fall.

(2.) *The influence of temperature.* There is no doubt that, in winter time, many diseases are aggravated by the intensity of cold, but cold is not a general cause of disease, especially in children under five years of age. This was shown by the late Dr. Farr, and in the volume recently published by the Sanitary Institute, it is pointed out that the death-rate of children in Norway is lower than that of England, while the death-rate of children in England is lower than in Italy, indicating that at this period of existence cold is not detrimental to life. When we come to isolate the deaths in particular months, and compare them with the periods of low water, it often happens that extreme low water in winter corresponds with periods of great cold, and low water in summer also corresponds with periods of great heat. It is only at such times when we are able to discount these influences by comparing them with periods when we have a normal state of things with reference to temperature, and abnormal in regard to ground water, that the influences measured by the ground water are brought into prominent relief.

In the records which have been published by M. A. Durand-Claye, for Paris, data are supplied which give information upon a variety of conditions likely, or unlikely, to affect or influence the development of typhoid fever, such, for example, as barometric pressure, wind direction and force, temperature, rainfall, observations with the actinometer, hygrometer, evaporation, geological character of soil, physical character of soil, height of ground water, density of population, distribution of water of various kinds, distribution of baths, lavatories, wash-houses, distribution of furnished apartments, length of sewers both in district and constructed during the period of epidemic in 1882.

Prof. Pettenkofer, in his researches, has not ignored the conditions which affect epidemic disease, for example the conditions

of wealth as contrasted with poverty. He clearly shows that poverty has a marked influence in the dissemination of disease, and that epidemics fall more lightly upon the wealthy classes than upon those who are not so happily circumstanced. However, occasionally typhoid fever attacks the wealthy classes more violently than those less favourably circumstanced, as was the case in Croydon in 1875. In this instance the best districts were in the position of receiving the poison in great intensity.

The records of Paris show that typhoid fever is lowest in June and highest in December and January. The increase generally takes place in August, or the period when percolation first commences. An exactly similar state of things is shown to be the case in Croydon. When the number of deaths is small the curve is not clearly indicated, but the lowest number of deaths occurs in June and the highest in January. The following figures show the proportion in the respective months in the period of registration at Croydon from 1837 to 1886, and in Paris 1865-69 and 1872-81.

Month.				Croydon.		Paris.
January	...	...	...	152	...	1,478
February	...	...	...	97	...	1,585
March	...	...	...	100	...	1,496
April	...	...	...	118	...	1,278
May	...	...	...	99	...	1,080
June	...	...	...	73	...	933
July	...	...	...	87	...	1,200
August	...	...	...	89	...	1,904
September	...	...	...	106	...	1,884
October	...	...	...	116	...	2,014
November	...	...	...	130	...	1,975
December	...	...	...	110	...	1,641
Totals				1,277		18,468

The different length of each month must be taken into account, and in the case of Croydon, where the numbers are small, the influence of epidemic periods interferes with regular order; but when these separate interferences which tend to swell the numbers in particular months—as, for instance, in April—are studied, they become clear indices, and point more directly to the conditions attending this particular disorder.

A very marked circumstance in connection with ground water, and the period of percolation, is shown in the case of deaths of children under five years of age. While there may have been mistakes with reference to the causes from which a child dies, very little error occurs with regard to its age. I am of opinion,

that the proper way of estimating the sanitary state of any period in any district, is by taking the number of children under five years of age, and calculating the deaths by the number living at these ages. The figures show, especially after deducting the deaths from diarrhœa, which are influenced by high temperature, that there is an almost exact parallelism between the period of percolation, and that of deaths occurring at those ages, the smallest number of children dying in the month of June, and the largest number in December and January. Moreover, the death-rate from year to year, fluctuates in a very marked manner with the fluctuations of the ground water. The most healthy periods are those in which there is the most ground water, and the least healthy are those in which there is the least ground water in any year. These results corroborate the strong relation which apparently exists between the state of water in the ground and zymotic diseases. It also shows that there are influences at work, which can be measured by the quantity of water in the ground, and which are destructive to young life, but which may be guarded against, as these influences indicate themselves many months before they begin to affect the population, therefore "to be forewarned is to be forearmed."

The fluctuation of the water line therefore is an essential condition in the development of disease, especially typhoid fever and cholera. It has been pointed out by Professor Pettenkofer that in those districts in which the rivers are held up at uniform levels by weirs, the conditions are favourable to health, and in such districts cholera rarely becomes epidemic. In a great measure this is corroborated in this country by the state of health at our sea-side resorts, which being the natural outflow for ground water, and owing to the uniform height of mean tide level, are without exception placed in a condition favourable to health. We have also the record in connection with the City of York, in which it is clearly shown by Dr. Laycock, in his report on York, published in the first volume of the Health of Towns Commission, that previous to the construction of the lock at Naburn below the city, the tide used to flow up above York, and there were considerable variations in the level of the waters from time to time, but after the construction of the lock in question the health of York materially improved. The health of districts, such as the Wandle Valley, is proverbial. In the latter district there are a large number of mills in a comparatively short length holding up the water to a uniform level.

With such examples for our guidance, it is clear that sewers may be of great advantage in maintaining uniformity in the water level. On the other hand, leaky sewers are liable not only to pollute the ground, but to cause considerably greater varia-

tion in the levels of underground water than would otherwise occur in various parts of the district. Good land drainage has a tendency to produce uniformity of water level, but such uniformity should rarely be attempted to be secured through the instrumentality of sewers carrying polluted matters. The influences which are observed clearly point out how important it is to guard districts against pollution of the earth. How little regard, however, has been paid to this point, for it is only within the last ten years that the importance of making sewers as water-tight as possible has received serious consideration, and still, in many parts of the country, sewers are being constructed without any regard to water-tightness and their other influences on ground water. Moreover, a large number of burial grounds have been established, in quite recent periods, in positions with respect to underground water which must more or less exercise a baneful influence on the health of the localities in which they are situated. Cess-pools, ash-pits and middensteads are still permitted to poison the air, ground, and water. No wonder that the towns which possess the means of most readily polluting the ground have, without exception, the highest rates of mortality. There can be no compromise in sanitary matters. It should be the aim of all sanitarians to preserve the ground from all impurities, especially in districts where the soil is of a porous character, and, above all, no supplies of water for dietetic purposes should be taken from wells sunk in the immediate subsoil in populous places; and finally, to secure the full measure of health, our houses should be so constructed as to prevent the admission of ground air into them.

The President of the Congress, Sir T. SPENCER WELLS, Bart., said, from the way in which the address of the President of the Section had been received, it was almost unnecessary to move a vote of thanks to him, but as the address could not be debated, he would ask the Dean of York to speak.

The DEAN OF YORK, rising in obedience to the call which had been made upon him, said he really did not know how to express his sense of the vastness of the subject, nor of the marvellous care, intelligence, and perseverance with which it had been dealt with by Mr. Baldwin Latham, and he offered his own tribute of appreciation of the extraordinary labour displayed in this address. This question of water supply had been one which had received the greatest attention of all peoples from the earliest times. The speaker had referred to the writing in the Scripture of the time of Zechariah; but

the subject was one of interest to Abraham, and there were many interesting passages concerning the matter which the President of the Section had discussed in this paper. What he had heard had impressed him with a sense of responsibility in regard to doing all that was possible to keep our supplies of water pure and unpolluted, and it was to be hoped that all persons who had had facts of such importance laid before them would do all they could to strengthen the hands of all who, like the President of the Section, were endeavouring to keep pure the supplies of water.

Mr. ROGERS FIELD (London), in seconding the vote, said that though he was aware there was some difference of opinion on some of the points raised by Mr. Latham, he coincided generally with the views expressed in the address, and more especially in what had been stated as to the lesson to be drawn by every sanitarian from the facts related, viz., that the ground must be kept clear from pollution if we wished to secure conditions essential to health. The speaker gave particulars of two cases which had come under his notice bearing upon the subjects under review, one being in connection with a large mansion in the Midlands, and the other in relation to a lunatic asylum in the West of England, in which there had been pollution of the water supplies, one from the drainage percolating into the wells, and the other by the drainage from a graveyard.

*On "River Pollution," by Professor HENRY ROBINSON,
M.Inst.C.E., F.G.S.*

IN view of the near approach of legislation on local government, there is every prospect that the long delayed subject of the prevention of the pollution of the rivers of this country will receive the attention which it deserves. In an address on "River Pollution," which was delivered at the "Parkes Museum of Hygiene" in 1885, I called attention to the various Bills which had been brought before Parliament for the purpose of effecting some improvement on the existing unsatisfactory state of affairs. I referred to the Standards of purity which were proposed to be adopted, and which have been the cause of much difference of opinion even amongst those who are striving to accomplish some reform. As this address elicited an expression of opinion from many well known authorities who took part in the discussion upon the occasion,