attended with a very small death rate. I refer to Dr. John P. Brison, of St. Louis.

DR. PARKER SYMS, in reply—I would urge the importance of one thing; if operation is to be done for the relief of prostatic obstruction it should be done early. When physicians and the public understand this and act upon their knowledge a great change will be wrought, for the mortality will be as nothing compared with that of the present time.

Concerning the Bottini operation, I am glad to state my views emphatically. I believe it to be an absolutely unsurgical and unscientific procedure. Years ago it was demonstrated and acknowledged by the profession that internal urethrotomy should not be performed deeper than the pendulous portion of the urethra. Whenever urethrotomy is necessary beyond that point, it should be done by external incision so that drainage may render it safe. This is acknowledged and accepted by the profession. The Bottini operation leaves a wound at the neck of a bladder which is full of infective urine. It is certainly a dangerous procedure. It is an incomplete operation, and I think it will be seen so to be when cases so operated upon have been observed more fully.

I would like to say concerning the failure to remove the prostate in my second case, to which Dr. Fuller has referred, that it was at a time before I had invented my retractor, and it was partly on account of this failure that I did devise the instrument. I believe to-day that every hypertrophied prostate can be removed through a simple median incision in the perineum.

Years ago I learned that when a statement was made by men to the effect that they were able to do things which I thought impossible, not to boldly assert that these things could not be done, for it has been my happy experience in watching the advance of progressive surgery, to find the impossibilities of yesterday become the possibilities of to-day. I think it is proper when one realizes his inability to perform a certain act, to say, "I can not do it," not to say "it can not be done," for there may be found someone who is able to accomplish it.

Concerning the choice of method of prostatectomy, I must say that I think a simple, median perineal wound—which is all that is left after operation as done by the method I have described above—is a far simpler and safer procedure than is any form of suprapubic cystotomy. The drainage is along the line of gravitation and is complete, the wound is a simple one and not large.

The patients are fairly comfortable and recover promptly. I have had mine sitting up at periods varying from the fifth to the tenth day. In two patients, as I have stated, there has been temporary dribbling after healing, but this has been owing to the fact that I made an unduly large cleft through the prostatic urethra. I am not sure to-day whether this is absolutely necessary in some cases or not, but I believe it is not necessary. I think it may be avoided by making the opening in the membranous urethra not too far back. In my last case I was careful to avoid this and was careful to separate the bladder from the prostate before I began the final enucleation. In that patient I removed the tube from the bladder on the fourth day, when I removed the packing; he had complete control of his bladder at once and made a most speedy recovery. That case, when I first saw him, had 52 ounces of very foul urine in the bladder. I did not operate nor empty the bladder at once but I succeeded with great difficulty in introducing a catheter and gradually emptying the bladder and slightly reducing its capacity.

Regarding the final functions of these bladders, I think it may be truly said that we can never restore them to histological perfection, but from the reports of the operators who have done the best work in this line, I think it can be said that these bladders do recover function fully. A bladder that is contracted may be dilated; a bladder that is dilated may be reduced in size. Certainly in my experience the patients have regained their functions so far that they have been able to hold their urine from four to six hours, holding from six to ten ounces, and have been able to empty their bladders, showing practically no residual urine.

THE TREATMENT OF PNEUMONIA.*

DE LANCEY ROCHESTER, M.D.

ASSOCIATE PROFESSOR OF MEDICINE, UNIVERSITY OF BUFFALO.

BUFFALO, N. Y.

In order to formulate a rational plan of treatment for a given disease it is necessary to recognize the nature of the disease, its course and mode of termination when recovery takes place, and its course and mode of termination when death occurs; it is also necessary to be able to recognize by symptoms and physical signs the changes that are occurring in the organs and tissues of the body during the course of the disease and to know the prognostic value of such changes when recognized.

Pneumonia is an acute infectious disease manifesting itself as a general toxemia, with a local disturbance in the lung passing through the clinically recognizable states of congestion, hepatization and resolution or disintegration.

A brief review of the clinical course of a favorable case shows us the initial chill; the high temperature; the increased frequency of respiration; the pulse of moderate frequency, full volume, high tension and normal rhythm; the coated tongue; the anorexia; the sluggish liver and inactive bowel; a kidney not functioning properly; the diminished excretion of chlorids; the presence in the urine of albumin, and usually casts; the more or less pronounced nervous symptoms; the cough; the rusty sputum; the dry hot skin; the flushed face of slightly dusky hue, and the leucocytosis. For a week or ten days these symptoms persist, the most important change occurring in the increasing frequency of respiration, and loss of the tension and sometimes change in rhythm of the pulse.

At the end of this period there occurs a profuse sweat, the temperature falls, the respiration becomes less frequent, the pulse improves in rhythm, the patient falls into a quiet sleep and awakens refreshed and well. If the case does not go on to recovery or dies before the time of the crisis, death occurs either from the intensity of the toxemia or from interference with oxygenation from involvement of too great extent of lung or from some complication.

The chief cause of death in pneumonia, exclusive of complications, is failure of the circulation through degeneration of the myocardium.

Several factors enter into the causation of this myocardial degeneration, namely, the toxemia of the infection acting directly upon the muscle fibers; the same cause active in the general system and producing an increase in the arterial tension throughout the body, thus making harder work for the myocardium of the left heart already suffering from toxemia; the morbid changes in the lung causing obstruction to onflow of the blood from the right heart, thus throwing harder work upon the right myocardium also suffering from the toxemia. The amount of pulmonary involvement would be, of course, a decided factor from this point of view, as well as from that of interference with oxygenation, too great an interference with oxygenation producing death.

Sometimes death occurs from the effect of the toxemia upon the brain—wild maniacal delirium wearing the patient out, or a low, muttering delirium ending in coma and death. Sometimes the kidney bears the

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Materia Medica, Pharmacy and Therapeutics, and approved for publication by the Executive Committee: Drs. Thomas Hunt Stucky, Leon L. Solomon and N. S. Davis, Jr.

brunt of the attack and uremia is added to the toxemia of the infecting organism.

Occasionally the gastro-intestinal tract is most seriously involved, and there occurs a violent diarrhea or a marked fibrinous inflammation of the lining mucous membrane.

The most common and serious complications of pneumonia from which death may occur are meningitis, pericarditis, endocarditis, nephritis and empyema. These, however, will not be considered in this discussion, which is limited to the treatment of pneumonia.

This review of the course and modes of termination of pneumonia shows that the most serious element in the disease is the toxemia, and next to this in degree of seriousness is the amount of lung involvement existing at any one time in the course of the disease. The first aim of the treatment, then, should be to get rid of toxins. The natural avenues of excretion by which toxins may be carried off are the respiratory, the renal, the gastro-intestinal and the cutaneous. In all cases of pneumonia the lung and the kidney are involved in the morbid process to a greater or less degree, according to the case. It would be obvious folly then to attempt to stimulate either of these tracts to any great excretory activity.

The bowels and the skin remain and nature has pointed out by the profuse sweat that occurs at the time of the crisis the avenue that she chooses for the greater part of the work of excretion. Accordingly, the prime therapeutic measures for relieving the toxemia are opening up of the bowels and skin and keeping them active. For this purpose there should be administered an initial dose of calomel of 0.50 gram followed by a saline laxative. This measure should be repeated during the course of the disease for the same purpose as the first dose and for the further purpose of relieving the distension of the right heart when indications of that condition are present.

To keep the skin active there is nothing more efficacious and less harmful than the proper use of the hot mustard foot bath. This can be given with next to no disturbance of the patient, the tub being put into the bed, with most excellent results so far as sweating is concerned. These baths may be kept up at four-hour intervals throughout the course of the disease. This sweating, in connection with the stimulation to be referred to later, is, in the opinion of the writer, the most important of the general therapeutic measures.

Another measure of value in preventing further bacterial invasion, is the frequent and thorough cleansing of the mouth with a mildly antiseptic solution. I think this a measure of considerable importance, as we know the pneumococcus is present in the mouth in considerable numbers.

While attempting to get rid of toxins by these means, it is important to put and keep the patient in the best possible condition to resist and overcome the toxemia that is present. For this purpose care should be exercised in the selection of a proper diet. This should be fluid and so far as possible should be sterile, so that no new poisons may be introduced. A large amount of pure water should be administered so that there may be plenty of fluid to flush the excretory organs. It is best administered in small amounts at short intervals.

Besides food, stimulation is necessary, and among stimulants strychnin and alcohol are the most useful in pneumonia. Moreover, the administration of them should not be delayed until failing circulation peremp-

torily calls for them. As soon as the diagnosis is made the strychnin should be begun in doses of 0.002 gram at four to six hour intervals as indicated. The interval may be shortened and the dose increased as symptoms which call for such increase in dose appear. As much as 0.004 gram every two hours may be given hypodermically with the best results.

Next to strychnin, alcohol is our best stimulant. Nevertheless, in some cases it seems to produce decided cerebral disturbance, particularly in those addicted to its abuse. In such cases ammonium carbonate or aromatic spirits of ammonium is a most useful substitute. These drugs are usually well borne by the stomach if combined with liquor ammonii acetatis and administered in mucilage or milk.

The local treatment of the lung is of importance. A careful physical examination of the chest should be made by the physician at every visit. In this way he can tell when congestion is beginning in any part; and vigorous local treatment will in many cases prevent the further progress of the morbid process. As soon as a portion of the lung is shown by physical examination to be in a state of beginning congestion, the prompt and thorough application of leeches, wet cups or dry cups over the part, will often stay the process there; or if it does not stop it entirely, will delay its progress twenty-four or forty-eight hours and give an opportunity for the clearing up of any previously involved portion of lung. My preference is for leeching or wet cupping. If dry cupping is resorted to, it should be very thorough, the skin and subcutaneous tissue rising up in the cup to as great a height as possible and turning blue or purple while there. The cups should be left on for a half hour and the process repeated every four to six hours. I have seen, postmortem, the direct anatomical evidence of the local value of this procedure, when thoroughly done. During the intervals between the applications of the cups the chest should be covered with a light woolen jacket. The poultice jacket should be relegated to the dark ages.

With ice applications I have not produced the relief of congestion desired. If, in spite of these local measures to relieve congestion, the process still persists and we begin to have evidence of over-distension of right heart, such as increasing cyanosis, the pulse losing its tension and becoming small and sometimes irregular, the liver enlarged, the veins pulsating and the area of cardiac dullness increasing chiefly to the right, whether a murmur of tricuspid insufficiency is present or not, prompt relief can usually be obtained by the withdrawal of 8, 10, or 12 ounces of blood from the median vein of the forearm. This procedure may be repeated several times if need be in the course of the disease. The procedure is of value both for its local effect in relieving the right heart distension and also for its general effect in withdrawing a decided amount of toxic material. At this time a little digitalis, given with due regard to its physiologic effect, may be of decided value.

After the withdrawal of blood, and in cases in which heart failure is imminent without evidence of right distension, the subcutaneous injection of normal salt solution has on several occasions proven of inestimable value. The use of the salt solution by rectum has also been of value, but of course does not give the prompt result that the hypodermic administration does. Whether the value of the procedure is dependent upon the restoration of bulk to the blood after the relief to the right heart has been obtained or whether the salt itself is of value, is

worthy of investigation, the fact being established that in three cases that have come under my observation—in one of them three times—the administration of the salt solution certainly had a marvelous effect in averting an evidently impending death.

If more than one lobe is involved the regular administration of pure oxygen gas at stated times, shortening the interval and prolonging the duration of the administration if cyanosis persists or increases, or if the frequency of breathing progresses to dyspnea, has proven of undoubted value in a number of cases.

In case of sleeplessness and restlessness chloralamid, chloral, Dover's powder, morphin and hyoscin hydrobromate are to be recommended in the order named; though with the thorough attention to the relief of toxemia by keeping the bowels and skin active, it is seldom that recourse to hypnotics is necessary.

On the possible complication of pneumonia I do not wish to touch, as that would take me beyond the scope of the subject announced; but I do desire to say that under the eliminative supporting treatment outlined complications are rare.

Before closing I wish to utter a protest against the treatment of pneumonia by large doses of digitalis in the early stages of the disease as irrational and unscientific, being based on a wrong conception of the nature of the disease.

If our conception of the infectious nature of pneumonia is correct, by the time the chill has taken place the disease is established, and any efforts to jugulate the disease by the use of remedies employed to overcome or prevent pulmonary congestion are most certainly misdirected, for preventing the change in the lung—and it is doubtful if the digitalis treatment can accomplish this—certainly will not overcome the toxemia; two of the most serious cases of pneumonia that have come under my observation have had no pulmonary involvement demonstrable by physical signs.

The treatment of pneumonia by the pneumococcus antitoxin is the natural and rational plan from the recognition of the infectious nature of the disease, but so far the attempts to produce a reliable antitoxin serum that will preserve its antitoxic powers have not been entirely successful, and consequently there are very great variations in the reports as to the results obtained by its use. However, improvements are being made and eventually I have no doubt that pneumonia will be put in the same category with diphtheria, which was formerly dreaded and of great mortality, but now, when carefully studied, diagnosed early and properly treated with large doses of antitoxin, shows 100 per cent. of recoveries. Until that time comes, however, we will get our best results by recognizing the essentially toxic nature of the disease and treating it along the eliminative supporting lines here suggested, which may be briefly summarized as follows:

1. The sustaining of the metabolic processes of the individual by the administration of easily digested or predigested foods in small quantities at stated intervals, the administration of large amounts of pure water for eliminative purposes and the administration of oxygen gas by inhalation whenever the absorbing surface of the pulmonary mucosa is involved to such extent as to interfere with proper metabolic oxygenation.

2. Elimination, (a) by the liver and bowel through the vigorous use of calomel and salts: (b) by the skin through sweats induced by external heat: (c) through

withdrawal of blood when indicated by right heart distension.

3. Stimulation of heart by strychnin, alcohol or ammonium carbonate, and in suitable cases by the subcutaneous injection of normal salt solution.

4. The local treatment of the lung by leeching, wet cupping or dry cupping as indicated.

As the value of the method of treatment detailed above was questioned at the time of the presentation of the paper, I beg leave to add, in publication, the statistics on which its value rests.

In some of the cases the notes are, unfortunately, decidedly incomplete, even the date of crisis not being mentioned, but enough is present—date, age, diagnosis, complications, treatment and result, to give them the necessary value for this statistical report.

Onset.	Termination.	Result.	Age.	Crises, Complications, Etc.
1. 1, '91.	1. 24, 91.	Recovery.	76	
2. 1.	3. 10.	"	14	
2. 9.	3. 28.	"	39	
2. 15.	3. 14.	"	52	
4. 14.	5. 21.	"	56	
8. 20.	9. 25.	"	82	
10. 31.	"	"	34	
1. 16, '92.	"	"	15	
1. 18.	"	"	29	
1. 18.	1. 31, '92.	"	74	
1. 31.	2. 20.	"	54	
2. 5.	2. 23.	"	22	
2. 7.	2. 21.	"	60	
2. 9.	2. 11.	Died.	35	Alcoholism; D. T.
3. 4.	"	Recovery.	37	
3. 6.	5. 20.	"	55	Chronic nephritis.
9. 30.	10. 14.	"	20	
10. 13.	11. 1.	"	35	
10. 26.	"	"	47	
1. 6, '93.	1. 21, '93.	"	23	Erysipelas
1. 4.	1. 19.	"	21	
1. 9.	1. 25.	"	24	
2. 22.	3. 15.	"	24	
2. 26.	3. 11.	"	30	
3. 6.	3. 21.	"	22	
3. 27.	4. 11.	"	45	
9. 13.	10. 10.	"	48	Entire left lung: patient was bled 16 ounces twice.
11. 4.	11. 23.	"	23	
11. 13.	12. 19.	"	20	
11. 18.	12. 12.	"	35	
11. 23.	12. 8.	"	50	
11. 27.	12. 4.	Died.	62	Sick one week before treatment.
11. 29.	12. 28.	Recovery.	40	Tuberculosis.
12. 7.	12. 9.	Died.	60	Sick eight days before treatment.
12. 14.	12. 28.	Recovery.	57	
12. 16.	1. 4, '94.	"	45	
12. 26.	3. 29.	"	18	Ecto- and pericarditis.
12. 27.	1. 9.	"	24	
1. 7, '94.	2. 4.	"	81	
1. 23.	3. 15.	"	32	
3. 8.	3. 29.	"	38	
3. 24.	4. 4.	"	32	
5. 20.	6. 6.	"	48	
7. 29.	8. 22.	"	81	
7. 30.	8. 24.	"	22	
9. 25.	10. 29.	"	82	
10. 7.	11. 21.	"	20	
10. 15.	10. 19.	Died.	27	Typhoid fever.
11. 3.	12. 5.	Recovery.	54	
11. 7.	11. 15.	"	29	
12. 2.	12. 24.	"	38	
11. 30.	12. 11.	"	20	
12. 9.	1. 4, '95.	"	12	
12. 10.	1. 2.	"	40	Double.
1. 11, '95.	1. 30.	"	43	Crisis tenth day.
1. 15.	1. 28.	"	21	
1. 17.	3. 14.	"	25	Typhoid fever.
1. 24.	2. 2.	"	33	
2. 2.	2. 19.	"	50	Mitral stenosis.
2. 12.	3. 7.	"	32	Crisis seventh day.
2. 13.	3. 25.	"	80	
2. 18.	3. 12.	"	38	Crisis ninth day.
2. 28.	3. 19.	"	32	Crisis seventh day.
2. 25.	3. 25.	"	25	
2. 25.	3. 26.	"	38	Developed lung abscess.
3. 19.	4. 17.	"	27	Crisis seventh day. Empyema.
5. 26.	6. 2.	Died.	50	Entire left lung; lower lobe of right.
9. 22.	10. 5.	Recovery.	30	
10. 20.	11. 13.	"	5	Crisis ninth day.
10. 20.	11. 11.	"	14	Crisis twelfth day.
11. 5.	11. 27.	"	70	
11. 6.	12. 3.	"	23	Crisis seventh day.
11. 5.	11. 30.	"	33	Crisis nineteenth day.
11. 3.	11. 22.	"	18	Crisis twelfth day.
11. 12.	11. 28.	"	23	Crisis tenth day.
11. 29.	12. 7.	Died.	33	Sick fourteen days before treatment (Alcoholism).
11. 23.	1. 8, '96.	Recovery.	67	Crisis ninth day.
12. 2.	12. 9.	"	25	
12. 1.	12. 13.	"	24	Crisis third day.
12. 2.	2. 23.	"	24	Crisis tenth day, followed by serous effusion.
1. 1, '96.	1. 19.	"	48	
1. 1.	1. 22.	"	48	Crisis ninth day.

Onset.	Termination.	Result.	Age.	Crises, Complications, Etc.
3, 12,	3, 18,	Died,	48	Alcoholism; sick two weeks before treatment; crisis March 16.
6, 28,	7, 20,	Recovery,	50	
8, 12,	8, 31,	"	26	
8, 14,	9, 25,	"	83	
9, 17,	12, 8,	"	15	12 Pericarditis.
9, 18,	9, 25,	Died,	12	
10, 7,	10, 15,	Recovery,	30	
11, 4,	11, 16,	Died,	61	Chronic nephritis, myocardial degeneration.
11, 14,	11, 18,	"	25	
11, 14,	12, 16,	Recovery,	78	
12, 23,	2, 17, '97,	"	29	Double; crisis twentieth day.
1, 2, '97,	1, 25,	"	26	Crisis seventh day.
1, 9,	1, 26,	"	36	Crisis seventh day.
1, 17,	2, 16,	"	41	
2, 19,	3, 27,	"	25	Crisis ninth day.
3, 12,	4, 10,	"	82	Morphine habits.
3, 20,	3, 26,	Died,	24	Double, pericarditis. [ment.
3, 20,	4, 2,	"	65	Nephritis, sick 8 days before treatment.
3, 24,	4, 14,	Recovery,	26	Crisis seventh day.
3, 29,	4, 22,	"	44	
8, 30,	9, 22,	"	17	Crisis seventh day.
9, 25,	10, 15,	"	32	Crisis ninth day.
11, 15,	12, 10,	"	84	
11, 26,	12, 31,	"	65	Crisis ninth day.
1, 1, '98,	1, 18, '98,	"	8	Crisis ninth day.
1, 15,	1, 19,	"	20	Crisis seventh day.
1, 29,	2, 24,	"	21	Crisis tenth day.
2, 26,	3, 9,	"	24	Crisis seventh day.
3, 1,	3, 29,	"	44	Chronic nephritis.
3, 17,	4, 15,	"	37	Crisis tenth day.
3, 28,	4, 15,	"	14	Crisis seventh day.
3, 26,	4, 20,	"	28	Pseudo-crisis fourth day, real crisis seventh day.
7, 2,	7, 17,	"	16	Crisis seventh day.
7, 17,	7, 31,	"	82	Crisis seventh day.
10, 10,	11, 21,	"	35	Crisis seventh day.
11, 15,	12, 17,	"	82	Crisis tenth day.
11, 28,	12, 8,	Died,	46	Purulent pericarditis.
11, 11,	12, 9,	Recovery,	18	Crisis seventh day.
11, 27,	1, 22, '99,	"	30	Crisis fourteenth day.
1, 10, '99,	2, 6,	"	43	Crisis ninth day.
1, 11,	1, 13,	Died,	45	Sick seven days before treatment; entire left, upper right.
2, 3,	3, 2,	Recovery,	13	Crisis tenth day.
2, 5,	2, 25,	"	40	Crisis tenth day.
3, 11,	3, 20,	Died,	32	Alcoholism; D. T.
3, 26,	4, 29,	Recovery,	60	Crisis tenth day.
3, 27,	4, 21,	"	56	Crisis tenth day.
4, 19,	5, 15,	"	83	Crisis tenth day.
7, 5,	3, 14,	"	27	Crisis tenth day.
8, 8,	8, 19,	"	29	Crisis fifth day.
9, 21,	10, 27,	"	80	Crisis tenth day.
11, 25,	12, 16,	"	22	Crisis seventh day.
11, 25,	1, 8, '00,	"	51	Crisis tenth day.
12, 16,	1, 1,	"	35	Crisis tenth day.
1, 12, '00,	1, 16,	Died,	45	Double.
1, 24,	2, 27,	Recovery,	50	Crisis tenth day; cystitis.
2, 16,	3, 7,	"	54	Asthma.
2, 19,	3, 7,	"	54	Mitral obstruction.
2, 20,	3, 19,	"	62	Crisis tenth day.
2, 22,	3, 22,	"	35	Crisis tenth day.
2, 28,	4, 10,	"	17	Crisis tenth day.
3, 1,	3, 22,	"	34	Crisis seventh day.
3, 11,	4, 16,	"	34	Crisis tenth day; diphtheria.
3, 25,	4, 16,	Died,	38	Double.
3, 10,	3, 26,	Recovery,	23	
3, 20,	"	"	26	
4, 1,	4, 6,	Died,	89	Edema of lung; asthma; dilated heart.
9, 12,	12, 13,	Recovery,	35	Relapsing pneumonia; three distinct attacks, with crisis on seventh, tenth and ninth days respectively.
9, 29,	10, 12,	"	25	Crisis seventh day.
10, 9,	12, 1,	"	35	Entire right lung; crisis tenth day.
10, 16,	10, 26,	"	39	Crisis fifth day.
10, 5,	11, 3,	"	12	Crisis fifth day.
10, 21,	11, 24,	"	23	
10, 25,	11, 1,	Died,	67	
10, 31,	11, 27,	Recovery,	61	Crisis seventh day.
10, 25,	12, 18,	"	46	Crisis tenth day.
11, 14,	12, 5,	"	35	Crisis tenth day.
11, 26,	12, 17,	"	19	Crisis tenth day.
1, 3, '01,	3, 7, '01,	"	54	Crisis tenth day.
1, 3,	2, 28,	"	82	Crisis tenth day.
1, 10,	1, 27,	"	50	Crisis seventh day.
1, 11,	2, 22,	"	50	Influenza.
1, 19,	2, 14,	"	40	Crisis tenth day; double pneumonia.
1, 21,	2, 12,	"	50	
1, 30,	2, 1,	Died,	46	Alcoholism; nephritis; entire left lung.
2, 21,	2, 27,	"	45	Double pneumonia.
3, 13,	4, 1,	Recovery,	49	Crisis seventh day.

The cases reported are from the service of Dr. Charles G. Stockton in the Buffalo General Hospital, my own service in the Buffalo Hospital of the Sisters of Charity and from my private practice.

Altogether, 168 cases were treated with 17 deaths. This gives a death rate of 10 1/8 per cent. This is better than most other methods of treatment, and if the cases that died are analyzed it will be found that four had been sick a week or more before treatment was instituted, and so can not fairly be considered, that two suffered with chronic nephritis, one had asthma and a di-

lated heart as well as being 89 years of age; that one had typhoid fever as a complication and that two entered the hospital with delirium tremens, no history obtainable as to the length of illness and died in less than 48 hours. If these cases are excluded, as I think they fairly may be, it reduces the number of cases to 158 and the deaths to 7, giving a death rate of 4.43 per cent. better than any other plan of treatment that I have heard of.

Although the number of cases reported is small it covers a period of a little over ten years and includes cases from all walks of life and of all ages, and may therefore, I think, be considered fairly representative.

THE ABORTIVE TREATMENT OF PNEUMONIA.

A PLEA FOR THE USE OF CARDIAC DEPRESSANTS IN THE TREATMENT OF THE CONGESTIVE STAGE OF PNEUMONIA.*

W. L. DICKERSON, M.D.

ST. LOUIS, MO.

In presenting this subject I have nothing new to offer and for this I apologize, but in this day of rapid advancement in medicine we are apt to reach out after the newer remedies and forget the older ones; therefore, I offer this plea for the use of one of the old cardiac depressants which seems to be rapidly giving way to new, and, I believe, less valuable remedies in the treatment of the congestive stage of sthenic types of pneumonia, namely veratrum viride.

That it is applicable to the congestive stage only and that the physician is often called too late to use it, I am perfectly willing to concede. I must admit also that its use really is an abortive treatment and that in the stage of hepatization is worse than useless and may be said to be malpractice.

We must also bear in mind the fact that many cases of pneumonia run through their course without medication and terminate satisfactorily; also that when we apply treatment we are often uncertain whether we have benefited our patient or not. Perhaps he would have gotten along just as well without our aid.

We may apply our abortive treatment and our patient get along admirably, but almost every practitioner, especially of the country, where we so often find cases of pneumonia in the robust, will tell you that he has met cases in which he was certain the patient must have a full-fledged pneumonia, bound to run its course, and yet in three or four days find him looking about over his farm seemingly a perfectly well man. We can only apply our remedies, and if successful, we are happy; if not, who can say but that our patient might have been worse had such treatment not been instituted.

In the congestive stage of pneumonia we must suppose that we have to contend with a condition not unlike an inflammation in other parts. When such inflammation takes place the vessels begin to dilate in the inflamed area, and as their lumen becomes larger the blood is attracted in this direction and soon a much larger amount of blood is accumulated here. The heart becomes somewhat excited and with each stroke increases the amount. Under such conditions it seems that the vessels in other portions of the body are contracted, thus increasing the amount sent to this area and lessening the opportunity to get away from it. A condition of stasis exists—stagnation, if you please. Still, the heart

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Materia Medica, Pharmacy and Therapeutics, and approved for publication by the Executive Committee: Drs. Thomas Hunt Stucky, Leon L. Solomon and N. S. Davis, Jr.