these structures on his species *A. binucleata* in Germany. But the host form in this country, according to Leidy's observations and my own, is *A. proteus*. As Mr. Woodward suggests, this fact of itself creates a presumption of the fungoid nature of the filaments.

b. The occurrence of the form at a limited season of the year is in keeping with the plant nature of the filaments. While one of the specimens of Mr. Archer appears to be an exception, all the others were found, I infer, in the early part of the year, from January to May. My own were taken in February or March of three successive years. The same locality was searched for them at other seasons in vain, though uninfested *Amœbæ* were found.

c. The hyphæ take no part in the movements of *Amœba*. If they bend or diverge, such movements are entirely passive, being due to contractions of the protoplasm to which their bases are attached. This is the explanation of what Archer interpreted as the creature's power of bending and quickly again erecting the filament at the point of constriction.* All observers agree that they are non-retractile. Cf. Fig. 7, a and b.

d. The progressive development of a single tuft is sufficient of itself to establish the main point of the thesis. It is clearly indicated in Figs. 8, 9 and 10, which are three of a series of five drawings representing as many stages of development of the same cluster. The fourth and fifth are not shown. Mr. Woodward tells me that after the first week the cluster of filaments became too complicated in its branching to draw, "although they always retained their original character of springing from a basal U-shaped filament and not branching near the distal extremities." Furthermore, Mr. Woodward observed on the same slide with this large specimen a number of small ones

* *Quar. Journ. Micr. Soc.*: 6, 190.

which "only after a week were found to possess any filaments."

* * * * *

An interesting inquiry remains to be made into the life history and relationships of the fungus itself, but upon that inquiry I cannot enter now.

WM. L. POTEAT.

WAKE FOREST COLLEGE, N. C.

NEW NAMES FOR *SPERMOPHILUS BREVICAUDUS*,
CANIS PALLIDUS AND *SOREX CAUDATUS* MERRIAM.

THREE names given by me to new species of mammals prove to be preoccupied and are here replaced.

Spermophilus chrysodeirus brevicaudus Merriam (Proc. Biol. Soc. Wash. VIII., 134, December 28, 1893), from the San Bernardino Mountains in California, is preoccupied by *Spermophilus brevicauda* Brandt (Bull. Acad. St. Petersburg, 1844, II., 369) from the southern Altai. I propose in its stead *Spermophilus* (*Callospermophilus*) *bernardinus*.

Canis pallidus Merriam (Proc. Biol. Soc., Wash., XI., 24, March 15, 1897), from the plains of Nebraska, is preoccupied by *Canis pallidus* Rüppell (Atlas zu Reise in Nördl. Afrika I., 33, taf. 11, 1826) from Kordofan. I propose in its stead *Canis nebracensis*.

Sorex saussurei caudatus Merriam (N. Am. Fauna, No. 10, p. 84, Dec. 31, 1895), from Reyes, Oaxaca, Mexico, is preoccupied by *Sorex caudatus* Hodgson (Horsfield's Catal. Mammals Mus. East India Co., p. 135, 1851), from Sikkim and Darjeling, India. I propose to replace it by *Sorex saussurei mutabilis*.

C. HART MERRIAM.

A NEW NAME FOR *MICROTUS INSULARIS*
BAILEY.

IN a preliminary paper describing several new Voles (Proc. Biol. Soc. Wash., XII., 86, April 30, 1898) I gave the name *Microtus insularis* to a species from Great Gull Island, N. Y. This name proves to be pre-