in the right arm and right leg is still very pronounced, the knee-jerks are both exaggerated and there is marked rigidity of the right leg. Paralysis of the sphincters persists. The apasia has changed very slightly, if at all. There is no optic neuritis.

Recently she was examined by Dr. Charles A. Oliver, who confirmed the existence of a right-sided hemianopsia and also demonstrated the presence of Wernicke's hemipupillary inaction sign.

The presence of the latter symptom is of great interest. In keeping with the profound hemiplegia (from which there has been no recovery) and with the persistent hemianesthesia, hemianopsia and sensory aphasia, this symptom indicates a most extensive lesion—one involving the entire posterior limb of the internal capsule, adjacent structures, such as the thalamus and radiations of Gratiolet and probably a large portion of the substance of the left hemisphere as far as the cortex. The lesion, judging from the history of the case, was doubtless vascular, though whether hemorrhage or embolism, it is difficult to say.

Regarding the neuritis which developed in the right arm no inference can be drawn. It is a rare sequel of hemiplegia, but has been seen by the writer several times. No adequate explanation has ever been advanced as to its origin.

25. EPILEPSIE RETROPULSIVE (Retropulsive Epilepsy). M. Lannois (Lyon médical, No. 30, 1899).

The author calls attention to a rare manifestation of epilepsy, *i. e.*, the tendency to walk backwards before or during an epileptic attack.

The first case was that of a gardener, who had a large number of epileptic attacks, in many of which he retained consciousness, and in all of which he walked backwards ten or twelve steps. There were no grand convulsive seizures. The second case was in a tuberculous patient, who, before the convulsion occurred, would walk backwards three steps and then fall, either forwards or backwards. The autopsy revealed no cerebral lesion.

The third and most interesting case is that of a woman *act.* 46. She had convulsions in infancy, followed by a right external strabismus. The convulsions and various automatic movements began at 17. She has daily convulsions. Very often she has peculiar attacks in which she becomes pale, rises from a sitting posture and walks backwards until she comes to an obstacle, when she begins to take her clothes off. If interrupted in this procedure she becomes angry and utters piercing cries. There are no convulsive movements of the limbs, but some twitchings of the facial, jaw and eye muscles, and frothing at the mouth. The attack lasts usually from five to fifteen minutes. She remembers nothing of the attack.

MCCARTHY.