made on the vaso-motor system of nerves and in the changes produced in the composition of the blood whereby re-establishment of the activity of the circulation in the congested part is promoted. The arteries, from being at first constricted, become dilated, and the blood is rendered thinner by the increased absorption into it of serous fluid from the interstices of the tissues which, in the dilated condition of the arteries, more freely takes place.

In many low states of the system very stubborn inflammations occur, and though, in most such cases, it would be improper to take away blood by venesection, leeches are, nevertheless, often of the greatest service. Though it is evident also that the powers of the system require to be raised by nutritious diet and strengthening medicines, it is bad practice to have recourse at once to the indiscriminate and premature employment of this treatment. The nutritious diet, even if swallowed, fails to nourish, and the tonic medicines make matters worse. In a case of scrofulous ophthalmia, for example, it is a common thing to see port wine, quinine and iron, and the like forthwith prescribed, instead of first trying to check the inflammation in the only way in which that can be most easily and safely done—viz., by the application of a leech or two to relieve the congested vessels, and the exhibition of a few doses of grey powder to correct the morbid state of liver and stomach. When by such medication the inflammation with the intolerance of light is abating, then, but not sooner, tonics may be advantageously given, with liberty to the patient to eat as much as he has appetite for. Counter-irritation, if had recourse to at all, ought never to be ordered until this convalescent stage of the disease has set in. Blistering at first only tortures the patient, and does no good, if it does not aggravate the inflammatory symptoms.

Scarification of the palpebral conjunctiva we frequently practise in cases of purulent ophthalmia and chronic conjunctivitis. The vessels of the part being in such cases much engorged, a few superficial scratches with the lancet are sufficient to give issue to a very considerable quantity of blood, and the propriety of the operation may be estimated by the freedom with which the blood is evacuated. After the bleeding has ceased we apply to the scarified surface the nitrate of silver in substance or solution, the red precipitate ointment or the like according to circumstances. It is, however, to be remembered that it is not the mere quantity of blood abstracted by scarification that does good. By first emptying the engorged bloodvessels, the subsidence of the inflammatory congestion admits of being more effectually promoted by the action of the irritating applications which I have said we make to the conjunctiva for that purpose immediately after the scarification. Without previous scarification the irritating applications make little beneficial impression. Often they do ill. As to the *modus operandi* of irritating applications in curing conjunctivitis, you will find it explained in my *Ophthalmic Medicine and Surgery*.

LITHOLAPAXY.

By HENRY J. BIGELOW,

PROFESSOR OF SURGERY IN HARVARD UNIVERSITY, SURGEON OF THE MASSACHUSETTS GENERAL HOSPITAL.

THE object of this operation is to remove at once, by the urethra, the débris created by the lithotrite. It is based on the belief of a tolerance by the bladder of protracted manipulation, which has not been hitherto recognised. It is also based on a theory that a brief manipulation, in the bladder, of polished and blunted instruments is less irritating to that organ than is the protracted presence of fragments, especially in view of the fact that the bladder hugs the latter more or less sharply at each micturition. When lithotrity is properly done, the process of crushing is easy and safe. The real question relates to the subsequent disposal of the fragments. If the purpose of the surgeon is to get rid of them, and if their evacuation has hitherto been, for the want of instruments, slow and injurious to the bladder, it is fair to consider whether this object cannot be better attained with the aid of an instrument devised to evacuate the bladder more quickly, through the urethra,

while its walls are held apart at their full normal dimensions, such, for example, as this canal attains when the meatus is compressed during the passage of urine.

It is quite safe to leave this question to be settled by such further experiments as shall be deemed wholly satisfactory. In the meantime, in order that the operation of evacuation may be accomplished in the readiest and safest way, it may be well to offer a few explanatory suggestions in respect to it.

One difficulty at the outset may be illustrated by a simple experiment. If a small catheter, surmounted by a funnel, be tied in an ox's bladder and held vertically, water introduced by the funnel will distend the bladder very forcibly. Similarly, the hydrostatic pressure of a column of water but a few inches higher than the pubes will distend the human bladder beyond its safe capacity. So that it may be assumed that any catheter surmounted by a bulb is more liable to injure the bladder by over-distending it than when the bulb is lowered to the level of the bladder or below it. The weight of a full bulb in the former position also prevents accurate manipulation of the catheter. It is more easily handled in a stand.

The instrument now employed is a syphon, at one of whose extremities is a large metal tube that takes advantage of the full normal capacity of the urethra, while the other terminates in an elastic bulb and glass receptacle. The tube extremity is shaped to facilitate its introduction, and also so that its orifice may be placed, if necessary, at the lowest point in the bladder, at the apex of an inverted funnel, among the fragments, without being occluded by the usual valve-like action of the bladder-wall. A straight tube has advantages over a curved one; the surgeon can more readily direct it and regulate its pressure; it is less liable to obstruction, and better cleared by a rod. Again, a straight instrument, when of the size 29, 30, or 31 of Charrière, is more easy to introduce than a curved one. In order to carry it into the bladder with least resistance, it should be passed as far as possible in a nearly vertical direction towards the anus, before the handle is depressed; its extremity then lies not only in front of the aperture in the triangular ligament, but, what is of more importance, below it; and is afterwards tilted up to this orifice by depressing the handle of the instrument to the axis of the body. The tube should be next rotated through this aperture like a corkscrew; and during this process, its point, which is made a little excentric for this purpose, revolves in a small circle until it discovers the orifice. A common difficulty in introducing a catheter or lithotrite results from depressing the handle prematurely, by which the beak is arrested above the aperture in the ligament, instead of below it. This error should be carefully avoided in introducing the large straight tube, and which so fills the urethra that it needs every advantage in traversing its less yielding parts. Indeed it is not superfluous to say that even a free use of oil injected with a syringe both into the urethra and into the tube may sometimes determine the question of the passage of the larger sizes. And yet, with these precautions, the whole manœuvre is a very easy one. Before introducing the tube the calibre of the urethra is ascertained by a common olive-pointed elastic bougie, or a sound. I have not found much advantage in trying to dilate it, unless at the moment of the operation, by incising the meatus, which is the narrowest part, or "divulsing" a stricture when it exists. Such preliminary enlargement of the contracted canal, unless especially contra-indicated, suggests itself.

But there is another important measure that does not suggest itself, and yet deserves careful attention. In order to protect the bladder from over-distension the first movement of the aspirator should be to withdraw water from it, and not to inject water into it. This cannot be too strongly emphasised. As preliminary to this precaution the surgeon ascertains exactly the capacity of the bladder. For this purpose it should be emptied by a catheter, and tepid water slowly injected, until the fluid is forcibly expelled between the catheter and the loosely grasped urethral walls. This reaction of the organ, which anaesthesia does not prevent, may be relied on to furnish a sufficiently accurate indication of the size of its cavity. If the fluid be now drawn off and measured, the same amount may be safely reinjected through the large tube, which is next to be introduced, the injected fluid being retained in the bladder by a tape tied round the penis, with the thumb, or a cork, at the extremity of the tube. Here is, I think, a critical moment of the operation, because if the contents of the bulb are added to those

of the already distended bladder the organ may be over-stretched. It should not be forgotten, therefore, that the elastic bulb, before it is attached to the tube, should be compressed until the sides meet, *in order that its first action may be to withdraw from the bladder, by its expansion, such an amount of fluid as can be reinjected with safety.* The present instrument holds ten ounces. By compression it loses five ounces. This amount would then be aspirated from the bladder by its first expansion, after which the usual pumping action displaces only about two ounces of fluid at each stroke, and can hardly injure the bladder. In order to direct attention to the importance of compressing the elastic bulb, a clamp is provided with the instrument to compress the bulb before it is attached to the tube. Other details, relating especially to the signs and avoidance of obstruction, may be found elsewhere. Briefly, when fragments are still felt, and yet cease to fall into the trap, there is either obstruction, which usually occurs in the tube, and should be removed by compressing the bulb or introducing the rod, or the fragments are too large to pass the tube, and need further crushing. In respect to the time occupied by the whole operation, I would suggest that if the surgeon will crush the stone as usual, and then remove the fragments, he will probably find his patient to be still in so good a condition that he will be tempted again to introduce the lithotrite, and again the tube. At the end of half an hour or more he will have repeated this several times, and will then find that the stone, if of average dimensions, has been evacuated. He will also find, in a common case, that the patient is no worse for the operation.

As these remarks apply chiefly to the mechanical part of lithotripsy, an operation which, indeed, depends more than any other for its success upon a careful and economical adaptation of its instruments to their employment, a few observations may be here added respecting the lithotrite.¹

It is very desirable, during a long operation, to prevent the impaction of the lithotrite, which interferes with crushing and sometimes injures the neck of the bladder. This may be effected by raising the floor of the female blade, and by adding lateral notches to the male blade, which should be also provided with a central septum at the heel. The lateral grooves for the male blade should also extend through the heel of the female blade. I am persuaded that the simplest movement for locking a lithotrite is a quarter rotation of the right wrist, without displacing the fingers of either hand. Further, it seems not to be always remembered by surgeons that the "rapidity" of a lithotrite depends upon the inclination of its screw thread, and that while the slowest screw gives most power and requires the strongest blades, "rapidity" sacrifices power. In the longer and more rapid operation now contemplated, larger and stronger blades than have been commonly employed, and which also better protect the bladder than do the latter, seem to me desirable.

I have elsewhere said that the blades of a lithotrite should be as nearly at right angles with the shaft, and their floor as straight, as is compatible with their convenient introduction into the bladder. Many instruments are made with an oblique blade, which is also so rounded at the heel as to curve their floor. This is a mistake. A cubical stone, for instance, would exactly fit a right-angled lithotrite. But when the same blades are made oblique, at an angle, for example, of forty-five degrees with the shaft, then, in order to grasp the same stone, they must not only be opened wider, but they will touch the stone at points nearer the shaft than before. In other words, the size of their grasp rapidly diminishes with their obliquity. They must be opened wider, and they seize less of the stone. Their power also diminishes, because, if they are made longer with the view of retaining the size of their grasp, their increasing leverage increases friction in the slide. This is readily seen by increasing their obliquity until they reach the line of the stem of the instrument, when they tend merely to roll the fragment between them. The latter then acts only as a wedge to separate them, and the friction of the slide is then greatest. In other words, right-angled blades crush best and wedge least. Oblique blades, on the contrary, wedge more and crush less, while the depth of their grasp is also less. And what is here true of the whole blade is true of any part of it—the heel, for example, which should not be oblique nor much rounded, but as nearly at right angles

with the shaft and with as straight a floor as is compatible with its convenient introduction. It may be added that the tip of the female blade should be bevelled, so that (if we compare it to a bent finger) it may impinge against the upper wall of the prostate, while passing it, rather with its pulp than with its nail.

In conclusion, it may be said that a small stone does not usually entail a serious operation of any kind. On the other hand, a large stone does. It may be better to resort to lithotomy in the case of a large stone, with a diseased bladder or kidney, however fatal the former operation may be in such cases. But there has hardly been any material change of late years in the operation of lithotomy, unless it be in a greater attention to cleanliness and in the general treatment of the patient. It is certain that no *bond fide* Lister dressing can be applied to the wound. The danger also of hæmorrhage remains the same as heretofore. So that, I think, any very great increase in the percentage of recoveries from lithotomy would, perhaps, result chiefly from a resort to the operation in more favourable cases than formerly, by which I mean in certain cases of stone which have been of late relegated to lithotripsy, and which might as well be subjects for experiment with the operation now proposed, as for lithotomy.

The question then reduces itself to the following:—

1. Whether, when any stone, large or small, has been crushed, it is not better to distend the urethra to its natural limits, and remove the detritus at once, by a few strokes of the syphon, than to leave it to be expelled through the contracted urethra by the unaided and comparatively ineffectual efforts of the bladder.

2. Whether, by thus assisting the bladder, we may not so increase the amount evacuated, and so diminish the time of the operation, that it may be possible to treat with success, by the method now proposed, larger stones than heretofore by lithotripsy.

NOTES ON SAN REMO AND THE RIVIERA.

By ARTHUR HILL HASSALL, M.D. LOND.,

FOUNDER OF AND CONSULTING PHYSICIAN TO THE NATIONAL HOSPITAL FOR CONSUMPTION AT VENTNOR.

THE time having now arrived when the patients of many medical men will be preparing to start for San Remo or some of the other health resorts on the Riviera, the following brief observations may not be uninteresting.

First, I would make a few remarks as to the journey here, which is really far less formidable than persons who merely read of the Riviera would suppose. I started from Cannon-street station on Monday, the 14th inst., by the 7.45 A.M. train, and reached Paris at 6.5 P.M. By the aid of one of the officials, who kindly got my luggage passed quickly, I was just enabled to reach the station of the Paris, Lyons, and Mediterranean railway in time to catch the express train, which left at 7.15, and arrived at Marseilles the following morning at 10.40. Here I was detained till 1.20 P.M., the train then travelling slower than the slowest of English parliamentary trains, stopping at the most insignificant stations, where not a single passenger alighted, and I consequently did not get further than Mentone, and this not until nine o'clock at night. The distance from Marseilles to Mentone is only 155 miles, and from the latter place to San Remo but twenty-two miles. The time occupied in the journey to Mentone was eight hours and a quarter, and from thence to San Remo, in consequence of a long detention at the border town, Ventimiglia, two hours and three-quarters. Now, had it not been for the long and apparently unnecessary detention at Marseilles and Ventimiglia, it would have been the easiest thing possible to have reached San Remo in good time the same night, thus accomplishing the whole journey from London to San Remo in thirty-six hours, and this notwithstanding the slow rate at which the train travelled from Marseilles.

But some will say that the time occupied in the journey is not of so much consequence to invalids, and that it is not desirable for them to travel such a long distance so quickly. This, no doubt, would be quite true in some, but not in the majority of cases, as if one night only be passed on the road,

¹ The aspirating syphon and the lithotrite have been made by Weiss and Son, with great mechanical perfection.