Localities: Devonian. Pilton (Phillips), Castleton; Hurst, Richmond, Yorkshire; Scotland, very generally distributed; Northumberland.

There are still remaining a few doubtful species, in my own, and in Mr. Aitken's collection. I cannot work these into the present paper for they are without a local habitation, and the fragments are not in such a state of preservation as to allow me to do them justice.

Now that attention has been directed to the Polyzoa of Derbyshire and North Yorkshire it is to be hoped that in the coming season for geological excursions some little attention will be given to the presence of these as well as to other, probably, more attractive groups of fossil remains.

ON THE GEOLOGY OF PALESTINE. BY W. H. HUDLESTON, M.A.,
F.G.S., &c.

IN the original choice of the subject, two principal considerations influenced me. In the first place, there is no foreign country whose history and geography attract more attention from all classes than does the Holy Land. It seemed therefore certain that a previous geographical acquaintance with the region might induce some curiosity as to its geological structure. And this brings me to the second consideration, which influenced my choice of a subject, viz., that Palestine—using the term in its widest geographical sense—contains within its borders some of the most remarkable features on the surface of the earth; and, notably, that astounding fissure, the Dead Sea Basin, which probably has exercised the imagination of most of us for a long period, and which geology, of all the sciences, is the most competent to investigate.

To facilitate reference, the subject has been divided into eight sections, as follows:—

- 1.—HISTORY OF THE INVESTIGATION OF THE REGION.
- 2.—PHYSICAL GEOGRAPHY.
- 3.—GEOLOGICAL OUTLINE AND EXPLANATION OF THE MAP.
- 4.—BEDS OLDER THAN THE CRETACEOUS LIMESTONE.
- 5.—THE CRETACEOUS AND NUMMULITIC LIMESTONES.
- 6.—POST-NUMMULITIC (MARINE) BEDS.
- 7.—VOLCANIC ROCKS.
- 8.—DEPOSITS OF THE DEAD SEA BASIN.

1. *History of the investigation of the Region.*—This branch of the enquiry is by no means devoid of interest, and might almost form a subject of itself, independent of the others. Of course we could go back as far as the days of the Crusades, for even then the fossil fish of the Lebanon were not unknown, since Richard Cœur de Lion is said to have noticed them; but for our purpose it will suffice to begin with the present century.

What we must now term the first occupation of Egypt by the British forces, in the early part of the present century, drew particular attention to the East, but such was the disturbed state of affairs during the period of the revolutionary wars that travelling was for the most part suspended, though some of the Royal Engineers about this time made short surveys in portions of Palestine. It is evident, however, that the occupation of Egypt had drawn attention to the East, for there was a meeting held of the *Palestine Association*, as it was then called, on the 24th of April, 1805. What they did is not known to me, though there is a tradition that agents were sent out, who got no further than Malta. Anyhow no active steps were taken by this body after the year 1809, and the Association may be deemed to have led a suspended existence up to 28th January, 1834, when a meeting was called, Mr. Bartle Frere in the chair, which resulted in the balance of its funds, amounting to £135 9s. 8d., being handed over to the *Royal Geographical Society*, and the original *Palestine Association* was, I presume, thenceforth regarded as defunct. But this is anticipating: we must now revert to an earlier date. At the close of the great war there was an immense increase of travelling all over the world, and the East very naturally came in for its share. Some of you are perhaps acquainted with the interesting narrative of Irby and Mangles, and with the classical work of Burkhardt, which, together with others published before 1830, laid the foundation of our physical knowledge of Sinai and Palestine.

Geology in those days was quite in its infancy, yet Burkhardt had an eye in that direction. But the first really geological work was one by Botta, entitled *Observations on Lebanon and Anti-Lebanon*, published in the Memoirs of the Geological Society of France, in

the year 1833. So that France led the way to geological research in Syria, and that lead she has since fully maintained.

I must now say a word about an important discovery. This was no less than a recognition of the singular physical fact, that the Dead Sea Basin, for a length of 200 miles, lies at a considerable depth below the level of the sea. Not only were the writers of antiquity ignorant of this peculiar feature, but it had also escaped the notice of Burkhardt and the earlier travellers, so that five and forty years ago, the whole world, scientific and unscientific, were quite ignorant of it. Like many great discoveries, it seems to have dawned upon several people nearly about the same time. It is not easy to assign the credit of the first discovery, but in the year 1837, two Englishmen, Messrs. Moore and Beck, published an account in the journal of the *Royal Geographical Society*. There was a considerable difference then and subsequently, as to the actual amount of the depression,—but to anticipate, I may say that the Surveyors of the *Palestine Exploration Fund* have fixed the level of the Dead Sea surface at 1,292ft. That was a time of general activity in the way of Palestine research, as the well-known work of Dr. Robinson appeared in the following year,—viz. in 1838,—and it was at this period also, that the Austrian mining engineer, Russegger, travelled, though his work was not published until ten years afterwards.

Encouraged by the success of their learned fellow-countryman, Dr. Robinson, the Americans entered into the investigation with great zeal, and the decade between 1840 and 1850 witnessed some important contributions from them. American missionaries have generally been observant men; some of them in Syria had made certain notes and collections of specimens, and these were submitted to Professor Hitchcock, who wrote a most important paper, so valuable indeed, that M. Lartet says, that Hitchcock, although never there, had a far better idea of the geology of Palestine than the majority of travellers, whether before or since his time; but then Hitchcock is a geologist, whilst the majority of travellers are not.

The efforts of the Americans culminated in the expedition undertaken by Lt. Lynch, in 1848, with Dr. Anderson as geologist. The party launched a couple of metal boats on the Lake of Tiberias,

and after an exciting and hazardous voyage down the Jordan, reached the Dead Sea in safety. Considerable difficulties were experienced, owing to the shallows, and to the tortuous character of the river, which describes a course of 200 miles, when the actual distance is 65 in a straight line. The Jordan was probably then navigated for the first time, though the *Rob Roy* canoe afterwards, I believe, ventured upon its waters. A very efficient investigation of the Dead Sea was made, and the leading physical features of the great fissure, known as the Ghor, ascertained beyond question. It was at the same time proved that even the weird and unwholesome nature of the place, in addition to Arab troubles, could not prevent a well-led and devoted party from doing its work effectually. Lynch's men were, I believe, all total abstainers. A large collection of fossils was made by Dr. Anderson, and many of these were described and figured in the fine quarto which recorded the results of the expedition.

After this adventurous journey there was a lull for some time in Palestine exploration: the cream of the mystery had been solved, though as yet but little real progress had been made in geology. But in the next decade renewed attention was paid to the subject, and two works appeared, which may be said to have placed the geology of Palestine on a firm basis. These were *Aus dem Orient*, by Dr. Oscar Fraas, and *La Géologie de la Palestine* by M. Louis Lartet. These authors have published other works on the subject, but I mention the above, as being amongst their most important contributions. Lartet's book was the outcome of his experiences as geologist to the celebrated expedition of the *Duc de Luynes*, undertaken in 1864, and which has been more fruitful in geological information than any other. The party commenced their investigations in the Lebanon, and thence passed by way of Samaria and Galilee to Jerusalem; then on to the Dead Sea, where they embarked on board the iron vessel, *Ségor*. Their examination of the Dead Sea was most thorough, methods being adopted for obtaining the water from various depths for analysis, anticipating as it seems in some respects those used by the *Challenger*. The party then ascended the right bank of the Jordan, about half-

way to the Lake of Tiberias, returning by the left bank, and so along the east side of the Dead Sea, through the mountains of Ammon and of Moab, which they were the first to examine from a scientific point of view. A second visit was made to Jerusalem, whence they again started by way of the mountains of Judæa for the Dead Sea, and so up the Wady Arabah into the Petra district, and thence over the watershed between the Wadies Arabah and Akabah, and down to the head of the gulf. It may be safely said that no observer ever brought away with him such an important mass of geological information respecting these difficult countries as did M. Lartet. In fact, it would be needless to say more of this particular journey at present, as the results form the basis of this communication.

About the same period, some good work was done rather outside the Palestine area, and chiefly in the Sinai district, by some of our own fellow-countrymen. I would especially mention the papers by Messrs. Duncan, Carter and Holland, in the *Quarterly Journal of the Geological Society*, and very important papers by Mr. Bauerman and Professor J. Milne. Mr. Bauerman's paper is by far the most valuable contribution we possess relative to the mining districts of the Sinai Peninsula.

We now come to the *Palestine Exploration Fund*, which was founded in 1865, "for the purpose of elucidating and illustrating the Bible, by a systematic survey of the Holy Land, by excavations, by the collection of traditions, manners and customs of the people, and by researches into the natural history, meteorology, and geology of the country." In the *Quarterly Statement* for 1869, it is remarked "Of this (*i.e.* of geology) we are in ignorance in almost every detail; the valley of the Jordan (and basin of the Dead Sea) is geologically one of the most remarkable on the earth's surface."

It is impossible to speak too highly of the topographical work of the survey, but somehow, very little geological work has been done, though there are papers by Captain Conder, which show that the subject possessed considerable attraction for him. His Physical description of the Holy Land,* and his chapter on the Jordan

---

* Tent Work in Palestine.

GEOLOGICAL SKETCH MAP of SINAI & PALESTINE, (based chiefly upon *M. Lartet's maps.*)

N.B. The figures represent elevations above the sea level in English feet:
those with a minus mark represent depressions below sea level.

To illustrate *Mr Hudleston's paper.*

	Littoral Deposits.	a
A	Deposits of Dead Sea basin.	a'
	Miocene?	b
B	Nummulitic & Grelaccous.	c
	Nubian Sandstone.	d
C	Neocomian or Jurassic.	e
	Schists, Gneiss, Granite, &c.	f
	Ancient Porphyry of Mt Hor.	g
	Basalts, &c.	h

valley,* should be read by everyone desirous of becoming acquainted with the geognosy of these regions. The one-inch map of Western Palestine was published by the Committee of the *Exploration Fund*, in 1880, and there is a very useful reduction, together with sections executed by Mr. Trelauney Saunders. Fig. 1 is a reduction from a meridional section constructed by him. I ought to mention here also the Ordnance Survey of the Peninsula of Sinai, undertaken in 1868–9, of which a most interesting account is given by the late Professor Palmer, in the first volume of the *Desert of the Exodus*.

Though scarcely geological, the admirable volume of the late Dean Stanley, and the adventurous narratives of Canon Tristram contain much geognostic information, and as they are well illustrated and ably and pleasantly written, these works have always been in great favour with the English public. In concluding this part of the subject, I may remark that as far back as the year 1867, no less than eighteen hundred and sixty works were enumerated on the subject of Palestine, and they have increased portentously since then. It is true, that very few of these contain the smallest item of geological information, yet much has to be read in order to pick up the scanty morsels. The importance of M. Lartet's works cannot be over-estimated.

2. *Physical Geography*.—It will be necessary to consider the geography before attacking the geology. According to Captain Conder, the limits originally assigned by Moses to the Holy Land extended from Mount Hor on the *south*, to the water-parting of the two great Lebanon rivers, Orontes and Leontes, on the *north*; and from the Mediterranean on the *west*; in the direction of the Euphrates on the *east*. For our purposes we must exceed these limits on the south, so as to include the whole of the peninsula of Sinai. Without attempting to give it a general name, we shall take into consideration the whole country, from, say Baalbec, which is very near the aforesaid water-parting, to the most southerly part of the peninsula of Sinai; the western boundary will be the Gulf of Suez, and the line of the canal; the eastern boundary stretches towards

---

* Hand Book to the Bible.

Fig. 1. Meridional Section of Western Palestine. (Reduced from Mr. Trelawney Saunders' section, by W.H.H.)

the Arabian desert.

In order to give some idea of distances, and by consequence of relative size, let us commence at the foot of Mount Hermon, which must be regarded as the parent of the Jordan, and the great condenser of moisture in those regions.

	MILES.
Foot of Hermon to the Lake of Tiberias	28
Lake of Tiberias, length	12
Jordan Valley... ..	65
Dead Sea, length	46
Dead Sea to Gulf of Akabah	114
Head of Gulf of Akabah to Jebel Mousa	105
	370

Again, the width of lower Galilee, from Acre to the Lake of Tiberias is about thirty miles, whilst the width of Judæa from the Mediterranean to the Dead Sea is about fifty miles.

Palestine, as we usually understand it, is an upland plateau, split longitudinally by the *crevasse* of the Dead Sea Basin, and it lies between the great limestone ranges of Lebanon and Anti-Lebanon (including Hermon) on the north, and the almost equally lofty granitic ranges of Sinai on the south. Lebanon and Anti-Lebanon roll over in a series of convolutions, but nowhere, according to recent survey, is there so high a point as Hermon, which must be regarded as the monarch of the region, with an elevation of 9,200ft. Hermon is the southern termination of the range of Anti-Lebanon, and looks down upon the mountains of upper Galilee, which barely attain to half its elevation. The volcanic mass of Jebel Kuleib, the hill of Bashan, attains an elevation of 5,600ft., but south of this, there are not many points over 3,000ft. on either side of the Jordan, till we come to Mount Hor, 4,360ft. Between this and the gulf of Akabah, are some lofty hills, one of which is the Sinai of Dr. Beke. There is one feature to which I would particularly draw attention, and that is to the very considerable watershed which exists between the Wady Arabah and the Wady Akabah.

The country west of the Wady Arabah is usually known as the Negeb, or south country, and was part of the "Land of Promise;" south of this again is a most terrible country known as the Desert

of the Tih, a limestone plateau, whose central point is marked in the maps as 1,480ft. This is about half way between Suez and Akabah, and is the place for which the late Professor Palmer was making when he was murdered.

A remarkable limestone escarpment forms the termination of the desert of the Tih, wedging itself into the Sinai Peninsula. The Negeb or south country, wedges itself into the Tih in a similar manner. In the Sinai Peninsula the mountains again become very lofty, and there are three granite ranges which especially claim attention: on the west is Jebel Serbal; in the centre is the Sinai group, culminating in Jebel Katarina, the highest peak in the peninsula, 8,536ft; on the south is the peak of Um Shomer, for a long time regarded as the highest of all. These lofty mountains of the Sinai Peninsula constitute the central section of the tripartite granitic region, which is severed by the gulfs of Suez and of Akabah.

Let us now direct our attention to the topography of western Palestine, since this has been surveyed with great accuracy (see Figure 1) which affords a meridional section of the whole country, I might say, from Dan unto Beersheba. In this section the upper sinuous line represents the watershed of western Palestine, between the Mediterranean and the Jordan valley; the middle line represented the level of the Mediterranean, and the lower line the course of the river Jordan. In sections of this kind the difference between the vertical and horizontal scales produces a false estimate of the abruptness of the slopes. Bearing this in mind, the section is an instructive one, though perhaps it requires a little reflection to understand.

The great basaltic mountains near Safed attain an elevation, as I before said, about half the height of Hermon: north-east of this great mass, lies the basin of the Hûleh marshes, which may be regarded as the *first* section of the Jordan valley system. In this marshy hollow the four rivers which form the Jordan unite, and open out into the waters of Merom or Lake Hûleh, whose surface may be regarded as nearly on a level with the Mediterranean. The great drop in the chasm of the Jordan valley occurs between Lake Hûleh and the lake of Tiberias, the fall being at the rate of 68ft.

per mile. This fall takes place between the volcanic masses of Safed and the Jaulan.

The *second* section of the valley includes the basin of the Lake of Tiberias, twelve and a half miles long, eight miles wide, and one hundred and fifty feet deep. There is a very great lessening in the height of the watershed opposite this basin, as though Lower Galilee had been let down bodily. The hills, though famous in history are amongst the most moderate elevations of the country.

The *third* section includes the gorge of the Kaukab, opposite Mount Tabor and Little Hermon, where there is another sharp drop.

The *fourth* section includes the Plain of Beisan, which connects by way of the Vale of Jezreel with the Plain of Esdraelon over a watershed, which cannot exceed 300ft., and which is therefore considerably lower than the watershed between the Wadies Arabah and Akabah. Indeed if the whole country were lowered between 300 and 400ft., perhaps rather less, the waters of the Mediterranean would flow into the great depression of the Dead Sea basin and fill it up, leaving still 300ft. before those of the Red Sea could meet them. The Vale of Jezreel is probably one of the most desirable and well-watered portions of the country, and fertile, like the Vale of Esdraelon, on the other side of the watershed. It might be tedious to detail the remaining four sections into which Captain Conder divides the Jordan valley. I will merely point out that it receives a considerable affluent from the west in the perennial stream that flows down the Wady Fârah, which, there is reason to think, may follow the line of some important dislocation, and, at all events marks a great change in the character of the country. The Wady Kelt flows into the Jordan through the Plain of Jericho, and there are many salt springs throughout this part of its course.

The shape of the basin of the Dead Sea may be gathered from the section. (Fig. 1). An extreme depth of 1,300ft. is marked, though this is a little in excess of any on M. Lartet's map. The southern end is quite shallow for many miles, so that the actual basin of the Dead Sea now occupied by deep water is less than one would suppose.

I would direct your attention for a moment to the watershed

which, if you were to trace it on the map, is pretty nearly equidistant, sometimes inclining towards the Mediterranean and sometimes towards the Jordan. It is not a little singular, how many places of interest are situated on, or close to the watershed—Nazareth, Shechem, Bethel, Jerusalem, Bethlehem, Hebron, illustrate the fact. Leaving the plains and moderate hills of Lower Galilee, we rise through the hills of Samaria to the Mountains of Judæa, and as over 1,000ft. has to be added to their height above the sea, the Mountains of Judæa show a massive section on the east side, the brow of the cliff or escarpment, overhanging the Dead Sea, being something like 2,000ft. above the great depression.

Now this great mountain group of Judæa is a natural fortress, and has been held for ages by a proud and determined race of highlanders, who had great contempt and hatred for such lowlanders as the Philistines. In after times the plateau was held by Saladin, whilst Richard overran the lowland plains. The plateau of limestone resembles very much the limestone pavements of Craven.

The physical geography of western Palestine would be incomplete without a word about the maritime plains. There is a strip of low sandy land all along the Mediterranean shore. In Phenicia, this is very narrow, and sometimes cut off altogether, as where the ladder of Tyre protrudes into the sea, and also where the still more conspicuous Mount Carmel makes a solid headland, the highest part of the block being 1,700ft. The Plain of Esdraelon is an inland plain, averaging between 200 and 300ft. in elevation, and covered with a rich basaltic loam; we have already seen that this is on a line of sinking in connection with the Vale of Jezreel. By all accounts, it is the most fruitful district in Palestine.

South of Carmel, along the coast, the maritime plain gradually widens, until the plain of Sharon merges into the wide expanse of Philistia. This is continued all the way to El-Arish, on the River of Egypt, though the boundary of Palestine proper may perhaps be regarded as on the Beersheba stream. In this portion of the maritime plain, Nile mud mingles with the drift sand, and further inland much of this is covered by the *debris* from the mountains. The dimensions are, 80 miles long, and near Gaza, 20 miles wide; 100–200ft. high.

3. *Explanation of the Map and Geological Column.*—"Geological Sketch Map of Sinai and Palestine." (Plate VI).

We must now turn our attention to the geology of the district, and by way of preface, I will give a brief geological demonstration of the map and vertical column (Fig. 2).

FIG. 2.

The use of a vertical column in geology is very considerable, as it enables people to see at a glance what they have to expect. At the same time it must be understood that no such section as is exhibited by this generalized column is ever seen in one place. In the map the *crystalline rocks* are all placed under the symbol *f*, and

this includes the schists, gneiss and granite, which form the nucleus of three mountainous masses, viz.: the Arabian chain (in Egypt), the Sinai Peninsula, and the mainland of Arabia. The crystalline rocks are continued in the Mount Seir range, and probably re-appear in the ancient porphyry of Mount Hor. It is quite possible that these three granitic areas may have formed one continuous mass, as doubtless they still do beneath the surface. At present, as probably for many ages past, they are separated by that curious bifurcation of the Red Sea, which produces the broad and shallow Gulf of Suez, and the deep and narrow Gulf of Akabah.

The next formation, or rather group of formations, is the *Nubian Sandstone*, which has an extensive development in connection with the granitic centres, from whose degradation it has in times past been largely derived. It is most extensively developed in the north of the Sinai Peninsula, where the upland plain of Ramlah is interposed between the chalky desert of the Tih, and the rugged mountains of Sinai. Thence it may be traced all along the east side of the Arabah, through Petra, to the eastern cliffs of the Dead Sea, and even up to the Jordan valley, half way to the Jabbok, where it disappears and is not seen again until we arrive at the great hollow between Lebanon and Anti-Lebanon, which is a continuation of the Jordan depression.

The age of the Nubian Sandstone as a formation, is a difficulty which we shall appreciate in coming to study it in detail. Meantime, for us, it represents everything between the crystalline schists, and the Cretaceous beds. The meaning of these two wedge-shaped masses, *s* and *e*, in the vertical section, Fig. 2, I will explain presently. Now consider for a moment what a variety of rocks there are in our own country, between the crystalline schists and the base of the chalk. There is the Cambrian, Silurian, Devonian, Carboniferous, Permian, Triassic, Jurassic and Neocomian. This shows the fundamental difference between the geology of England and of Palestine. Can the Nubian Sandstone represent all these? Surely not,—yet it may well represent more than one of the great systems in time.

Next in order and far more extensive are the *Cretaceous* and *Nummulitic Limestones*, which occupy such a large area in the map,

and constitute indeed, the entire mass of the Palestine hills, of the Negeb, or south country, and of the terrible desert of the Tih. They are classed together, because of their physical unity, as one great calcareo-siliceous formation. Indeed the only test of difference is the presence of true nummulites. Now there can be no doubt that in Egypt, large portions of the area coloured blue on the map are nummulitic limestones; and also that a considerable portion of the western escarpment of the Tih is nummulitic chalk with flints; there is also a limited area near the head of Wady Akabah. The only localities known to yield nummulites in Palestine proper are the Mountains Ebal and Gerizim, in Samaria; and Mount Carmel, on the coast. There is also a small exposure of nummulitic limestone in Phenicia. Beds of Cretaceous age are largely developed in the Lebanon.

With respect to the portion of the map and section described as Miocene, there is some doubt. All the evidence goes to show that beds usually called "Miocene," were cut through in the southern portion of the Suez Canal, but how far these extend is by no means clear. Of course no great accuracy is claimed for the map in this respect. There is a quantity of flint breccia, the result of the breaking up of the Nummulo-cretaceous series, scattered about in different parts; some of which may be of Miocene age. In one place, near the Gulf of Suez, this was found to be associated with corals of a Miocene type.

The Deposits of the Dead Sea Basin include everything in the long fissure from the foot of Hermon to beyond the roots of Mount Hor. All we can say of them is that they are post-nummulitic, and the same may also be said as regards the *Lavas*, which are so abundant in the north.

The Coast deposits, marked *a*, are regarded for the most part as Quaternary and Recent; they are said to contain shell limestones, much used for building the cities on the Mediterranean; the shells being all those of recent species.

Since the Cretaceous limestones are by far the most important rocks in the country, they have a section to themselves, No. 5. The whole of the rocks below them are detailed under section 4, and those above them in section 6.

4. *Rocks older than the Cretaceous Limestones.*—We must of course begin with the crystalline rocks, of which granite is the most important, but with it are associated those porphyries, which in the Egyptian granitic region have yielded such fine building stones.

The mountainous core of the Sinai Peninsula, including all the great ranges before mentioned, consist of some variety of granite, and it is evident that red felspar granites are the most abundant, and characterize the region by the colour which they impart to the rocks. We obtain our best account of the physical aspect of this region, and of the peculiar features produced by granite and its associated rocks from the pen of the late Professor Palmer.*

This lamented author ascended most of the principal granite mountains, such as Jebel Mousa, Jebel Katherina, &c., and brought away specimens of the rocks; some of which by the kindness of Professor Bonney I have been able to inspect.

Rock Specimen 1.—Let us begin with the summit of Jebel Katherina. This is a *syenitic granite* of a peculiar nature: in a pinkish felspar base, which occupies the bulk of the mass, are small crystals of orthoclase, quartz, and hornblende. The quartz is not interstitial as in the more typical granites.

Rock Specimen 2.—Jebel Musa. This is from Mount Sinai itself, and is a mixture of coarse red felspar and grey vitreous quartz. There is very little trace of any third mineral, so that the rock nearly approaches a pegmatite.

Rock Specimen 3.—Summit of Serbal. This is a more thorough pegmatite than the other, and in better condition; the felspar is a bright red. The two last specimens are very typical of the granites of the Sinai Peninsula.

Serbal is especially remarkable for the number of diorite dykes which traverse the softer granite, and project like so many teeth, according to Dr. Fraas, from the ridge forming the summit of the mountain. The serrated aspect which one sees in pictures may be due to this cause.

We obtain a considerable amount of information relative to the granites of this region from Professor Milne, who accompanied the

---

* Desert of the Exodus, vol. I., p. 27.

late Dr. Beke in search of the true Sinai. They avoided Jebel Musa, and the whole region of the traditional Sinai, and proceeded to the head of the Gulf of Akabah, where Dr. Beke fully expected to find the volcano from which, according to him, the law was delivered. This supposed volcano is situated a few miles to the north-east of Fort Akabah, and turned out to be a granite mountain, of which Professor Milne gives a good description.¹ The rock was found to be very prone to disintegration, but the dykes which traverse it decay still more rapidly.* From Professor Milne also, we have testimony to the predominance of the felspathic element in the granites and the dykes by which they are traversed. Of seventy two specimens examined, only two approached syenite. It is probable that these observations apply mainly to rocks collected near the gulf of Akabah.

Leaving the crystalline rocks, we now find ourselves face to face with that difficult problem of the *Nubian Sandstone*. The relations of the Nubian Sandstone to the crystalline rocks are well shown in Mr. Bauerman's (Fig. 3) section. This may be regarded as the first stratigraphical aspect of the Nubian Sandstone. As we follow its long exposure on the east side of the Arabah, no base is seen,—this is the second stratigraphical aspect. Then it disappears, and its representative in the Lebanon rests upon the *Glandarius* limestone, and this probably upon the Jurassic rocks of Hermon. I have endeavoured to show this by inserting the wedge-shaped mass, *e*, into the stratigraphical column.

With the help of Mr. Bauerman, let us examine the Nubian Sandstone of the Wady Nasb, where its base is distinctly seen, so that we know that there is nothing but crystalline rocks underlying it. I need not detail the lithological characters, but the chief point of interest lies in the fact that a thin bed of limestone here occurs in it, which contains undoubted Carboniferous fossils. There was much discussion about this at first, but Mr. Holland subsequently sent better specimens, one of which, *Orthis Michelini*, was in very good

---

* The episode of Jebel Baghir, or Jebel-el-Nur (the mountain of light), the "true Mount Sinai," first a volcano, then a lightning-haunted peak, is very amusing.

FIG. 4.—VIEW OF THE ARABAH FROM MOUNT HOR.

condition, and Mr. Davidson had no difficulty in determining it. There are other indications that portions of the Nubian Sandstone of the Sinai Peninsula are of the Carboniferous age. There is for instance, in the Museum of the Geological Society, a cast of vegetable impression, marked *Lepidodendron Mosaicum*. Formerly a certain amount of suspicion attached to this fossil, and doubts were thrown upon its being an authentic specimen from the Sinai Desert. It was, I believe, brought to the late Sir Roderick Murchison, and probably received its name from the late Mr. Salter. Since the discoveries of undoubted carboniferous Brachiopoda there seems no reason to doubt the so-called *L. Mosaicum* is a genuine Sinai fossil.

At the base of the Tih escarpment is a fossiliferous deposit containing Cenomanian urchins, &c. There is a strong tendency on the part of both M. Lartet and Dr. Fraas to regard the whole or nearly the whole of the Nubian Sandstone as of Cenomanian age. Dr. Fraas, in his last publication is very decided on this point, and if his statements are to be relied upon, the Carboniferous exposure in the Wady Nasb must present a very exceptional phase.

Let me say a word about the turquoise mines. These occur at Serabit-el-Khadim and other places, in the Wady Nasb district, where they have been worked at intervals of time for ages. This was a great mining country for the ancient Egyptians. The turquoise occurs for the most part in ochreous balls that are situated in certain horizons of the Nubian Sandstone. The best stones are in the solid sandstone, a short distance from the joints. Here again, Mr. Bauerman is at variance with Dr. Fraas, who says, that the turquoise occur in gneiss. The mineral wealth of the sandstone region is very considerable, both iron and copper occurring in abundance. The Arabs were very recently working for turquoise, and have destroyed many inscriptions in these sandstones. Burton, a few years ago, made an expedition into the land of Midian, where the sandstone prevails, and brought away great quantities of minerals.

The last thing to consider is the effect which the Nubian Sandstone has on the scenery of the districts where it prevails. Professor Palmer gives us some capital descriptions.*

---

* Vol. cit., p. 17.

The *red* of Edom is notorious, but Dean Stanley says that it should be regarded as *dull crimson*. His account of Petra in "Sinai and Palestine," is the most vivid description of Nubian Sandstone known to me.

The summit of Mount Hor, where Aaron died, is composed of Nubian Sandstone. Here is a sketch (Fig.4), taken from "La Géologie de Palestine," which may serve to give an idea of the chaos of rocks in that mountainous region. The masses of porphyry form an important feature. These were formerly supposed to be eruptive, as they seem to come through both the sandstones and Cretaceous limestones. Thus it was held by some that the country had been lifted up by these rocks, and that in this way the watershed between the Wady Arabah and the Wady Akabah had been formed. M. Lartet supposed that these are old eruptive masses, belonging to the crystalline series, and that they must have been there before the deposition of the sedimentary beds. This has an important bearing on questions which will be discussed in section VIII.

The cliffs on the east side of the Dead Sea and the deep gorges of the Arnon and Calirrhöe presents splendid sections of the Nubian Sandstone, where they may be seen underlying the Cretaceous limestones of Moab. The sandstone here has fallen 5,000–6,000ft. below the elevated position it occupies on the summit of Mount Hor. The pages of M. Lartet and of Canon Tristram supply us with most graphic accounts of the extraordinary effect of these cliffs. The gorge of the Calirrhöe is especially remarkable, as there the sides of the deep "cânon" are moulded in black basalt, white limestone, and purple sandstone, the effect of contrast being most marked. Canon Tristram says, that the gorge leading up to Kerak is almost Alpine in its grandeur, and that author considers that there is a decided unconformity between the Nubian Sandstone of that district and the overlying Cretaceous limestones.

Throughout the upper part of the Jordan Valley the Nubian Sandstone is lost sight of, but re-appears in the hollow of Syria. Before noticing its character there, we must consider the beds known to occur in Hermon, which in the stratigraphical column are marked under the letter *e*. These are what are known to the Leb-

anon geologists as the *Glandarius* limestone, which by the testimony of all underlies the so-called Nubian Sandstone of that region. The spines of *Cidaris glandarius* are characteristic of this formation, and have long attracted attention. These were known formerly as "Olives of Sodom," or "Judah stones," though not really found either at Sodom or in Judæa. M. Lartet thought that some portion of the *Glandarius* limestone might be Jurassic, but the Lebanon geologists regard it as of the age of the Upper Greensand: *Syphonia pyriformis*, *Cyphosoma cenomanense*, *Terebratula biplicata* are quoted from it.

The only exposure of undoubted Jurassic rocks occurs on Hermon, where *Rhynchonella lacunosa* has been found on the very summit. There are palæontological indications both of the White and Brown Juras, whilst fossils not uncommon in our Coralline Oolite occur, such as *Ammonites plicatilis* and *Terberatula Maltonensis*, both of which are found in the neighbourhood of Malton. The Brown Jura is represented by Ammonites, such as occur in the Kelloway Rock of Scarborough Castle Hill.

The Lebanon Sandstone is described in considerable detail by Dr. Fraas. The fossils are numerous and leave little doubt as to the Cenomanian age of these beds. Trigonias are very characteristic, and especially a variety of the group of *Scabra*, one of which is figured in the accompanying plate. There is a considerable amount of contemporaneous igneous rock in these sandstones.

On reviewing the whole question of the Nubian Sandstone, we must admit that the identity of the Lebanon rock with that of the Dead Sea, &c., is not quite so clear as Fraas and Lartet maintain, but in our present state of knowledge we must be content to let the question rest.

5. *The Cretaceous and Nummulitic Limestones.*—The earlier geologists, especially those connected with the Lynch Expedition, believed that there was a considerable development of Jurassic beds, both in the Lebanon and in Palestine proper, and thus many of the fossils are described as Jurassic. There is very little doubt now-a-days that the whole of the calcareous system, with the slight exceptions already made, belongs to the Cretaceous epoch or to yet

newer beds. Hitherto we have met with few traces of fossils, but when we reach an horizon which is represented in N.W. Europe by the zone of *Pecten asper*, *Pygaster lampas*, *Scaphites equalis*, *Ammonites navicularis*, and *Exogyra columba*, there is a great change. This horizon is Cenomanian, and partly corresponds with our Upper Greensand, and it forms the base of the Cretaceous limestones over an immense track of country, extending from S.E. Arabia, and probably even further on the east, to Algeria on the west; so that Palestine forms but a fragment of this extensive area. The deposits of this old sea went on accumulating long after the period when chalk had ceased to be deposited in our own islands, and plainly continued even where Nummulites had become numerous, so that there is no physical break in this great calcareous system. An important change took place in middle Tertiary times, the Cretaceo-nummulitic sediments had become rock, and were subjected to a series of earth movements, such as those which formed the chains of Lebanon and Anti-Lebanon, where this group of beds are folded in undulations which have a general N.N.E. strike. At the same time, the Cretaceous limestones of Palestine were uplifted in the great anticlinal fold, of which Fig. 7 is a cross section. But the complications here are slight compared with those of the northern ranges. It was at this period that the main physical features of the country were written upon it, though the agents of denudation have been carving minor features ever since. We may possibly regard the period of the uplift as Miocene. No marine fossils have ever been deposited since in the area coloured blue on the map, but there are immense deposits of flint conglomerate, gravel and superficial rubbish, which have been accumulating for an immense period of time.

The Cretaceous geology of the Lebanon has been the subject of much study, and of several important memoirs. The first is the memoir by Botta, to which I have already alluded, as the earliest *geological* attempt made in the east; the latest work is the second part of *Aus dem Orient* by Dr. Fraas, published four years ago.

The Lebanon sandstone is credited with the Trigonias already mentioned. Above this comes the Gasteropod zone towards the base

of the Cretaceous Limestones. It contains many species of *Nerinaea*, two of which are figured in the accompanying plate; next succeeds the zone of *Ammonites syriacus*. Dr. Fraas regards this as Turonian, or what we should call lower chalk, the typical fossil clearly having affinities with *A. rhotomagensis*, and other ammonites found towards the base of the chalk at Lewes. The presence of Hippurites in abundance is another characteristic feature. The upper white chalk-with-flints of the Lebanon Dr. Fraas refers to the Senonian or upper chalk, so that he does not regard it in any way as Tertiary. The remains of the Nummulite beds consequently are rarer in the Lebanon even than in Palestine, but evidently increase in importance as we approach Egypt.

It is impossible to quit the subject of the Cretaceous Beds of Lebanon without saying a word about the fish deposits. These are on two horizons, of which that of Hakel is probably the lower. The fishes are all of those species whose bones are perfect, ganoids being absent. The beds therefore are either Cretaceous or Tertiary, almost certainly the former. There is an enormous collection of these Lebanon fishes at the British Museum, and along with these are some wonderfully well preserved remains of cuttle-fish, &c., one of which Dr. Woodward has lately described in the *Geological Magazine*. Other forms are equally well preserved. The Sâhil Alma deposit is thought to be on a higher horizon, near the white chalk-with-flints, and may possibly correspond with an horizon in Palestine, where M. Lartet detected many remains of fish.

We must now return to Palestine. Not the least valuable and original of M. Lartet's work, was that done in the Cretaceous limestones of Moab, a country always difficult and even dangerous to travel in. He gives us a great number of important sections. I have selected the section across the gorge of the Arnon (Wady Mojib), because it shows the whole body of the Cretaceous limestones from bottom to top, everything in fact, between the Nubian Sandstone and the Basalt.

The complete section is rather over 3,000 feet, and when we have deducted some hundreds for the Nubian sandstone at the bottom, and Basalt at the top, there still remains something like 2,500ft. for

FIG. 5.—SECTION ACROSS THE VALLEY OF THE ARNON
(W. MOJIB) EAST OF THE DEAD SEA.

a, Limestones with flint weathering red; *b*, marls with *Pholadomya*; *c*, grey, compact limestones; *d*, alternations of yellow and red chalky marls with tabular limestones containing *Am. Luynesii*; *e*, yellowish limestone with *Ostrea Mermeti*, var. *minor*, and *O. vesicularis*, var. *judaica*; *f*, grey marls with *Hemiaster Fourneli*, *O. olisiponensis*, *Plicatula*, *Pholadomya*, *Venus*; *g*, marly limestones with bivalves and gasteropods; *i*, yellow limestones with *Pterodontes* and other gasteropods; *j*, limestones with *Ostrea flabellata*, *O. africana*, var., *Holctypus Lartetii*, *Heterodiadema lybicum*, *Pterodonta elongata*; *k*, green saline marl; *l*, white sandstone; *m*, red sandstone.

the Cretaceous limestones. Three-fourths of this, comprising the beds, form *a* to *j* inclusive, represent the middle and lower chalks of Palestine, not necessarily corresponding with the middle and lower chalk of England, but about that horizon. There are beds of hard and soft alternations, containing a recognized and tolerably abundant fauna.

In western Palestine these middle and lower beds are for the most part seen, as here, in the gorges, though sometimes perhaps brought to the surface of the plateau by synclinal folds. These are the beds which it has so often been the fashion to speak of as Neocomian and Jurassic, but whose very lowest horizon contains a fauna which M. Lartet regards as Cenomanian. The gorge of the Wady Mojib also shows a fine section of the upper white chalk-with-flints,

from which, at another point near Kerak, a remarkable series of fossils was obtained.

In this one section then we have an epitome of the entire series of the Cretaceous limestones of Judæa and of Moab. These are the beds, and especially the upper chalky ones with flint, whose acquaintance the traveller in the Holy Land is most apt to make. Denudation, going on probably ever since the Miocene period, has carved them out in a variety of ways, and the soft upper beds have received much ill treatment. Sometimes everything has been washed away except the flints, which are left in long rows, or as circlets round banks and knolls of the harder rocks. Sometimes the material from these soft upper beds, fills cracks and crevices in the lower and harder ones. Generally one may say that they make a dry soil, and have a tendency to produce a wilderness, such as that of Judah. This kind of thing is probably repeated in the more terrible desert of the Tih, where perhaps there are considerable developments of Nummulitic limestone in addition. All these upper limestones are impregnated with saline matter, and the water from them is both bitter and unclean.

I ought to say a few words about the palæontology of the Cretaceous beds of Palestine. There are two or three noted localities for fossils. One is reported to be near Jerusalem. At any rate numerous Ammonites are found on a certain horizon there, such as French geologists would call Cenomanian.

They belong to a type not unfamiliar to collectors from the lower chalk or chalkmarl, such as *Am. Mantelli*, *Am. rhotomagensis*, *Am. varians*, and along with these are others not known in England, though for the most part related to the same group. The Nerinæan limestone of Jerusalem is celebrated; it will take a good polish, I believe, and I have heard say that the temple of Solomon was in part constructed of this stone. Its excavation has left great galleries in the rocks beneath. The abundance of Nerinæa is noteworthy. At least twelve species are quoted by Lartet, not of course all from the Jerusalem rock; and it is this abundance of Nerinæa which may have induced the older geologists to regard these limestones as Jurassic, since in our country no Nerinæa

is known in beds of later date than the Corallian,* but the presence of the Ammonites named, of Baculites, and especially of Hippurites, leaves no doubt as to the age of these beds.

Amongst the conchifera there are several specimens of *Pholadomya*. A small but very handsome species is one from the upper chalk of Judæa, figured in the plate. There can be no doubt that it has a tertiary aspect, and reminds one in its deeply chiselled ornaments, of a species figured by Sowerby from the Eocene, I think, of the Isle of Wight. This is probably from Marsaba or Nebi Musa, where there is a well-marked fossiliferous horizon. The most numerous of all the Cretaceous fossils are oysters; these undergo division into several species. The exogyroid forms are related to *Exogyra columba*. The one figured in the accompanying plate is pretty common, and has been identified by Lartet as the oyster of Lisbon (*Ost. Olisiponensis*), described by Sharpe from the Cretaceous beds of Portugal. Sea urchins, especially Hemiasters, are also plentiful.

It remains for me to say a few words about the bituminous deposits of Nebi Musa, which underlie the fossiliferous horizon already mentioned. There is an abundance of bituminous matter in many parts of the Cretaceous series of Palestine, the Lebanon, &c., but this particular place is well-known and often visited, and the black bituminous limestone which it yields is worked up into rosaries and other ornaments by the folks at Bethlehem and Jerusalem. It is known by the name of Dead Sea stone, as the idea of asphalt has always been associated with that lake. The bed of Nebi Musa, however, is a couple of thousand feet above the level of the Dead Sea, and several miles away from it. The limestone, according to Hitchcocks report, contains 25 p.c. of bitumen, and would be valuable as an article for producing asphalt. It is worthy of remark, that just below the chief deposit of bitumen, is a bed full of fish remains. Now, organic matter is the source of all bitumen, which is nothing more than a partially oxidized hydrocarbon with certain impurities;

---

* Mr. Walter Keeping, curator of the York Museum, has lately described and figured a very peculiar *Nerinea* from the Lower Greensand of Upware. If this is not a *remanié* fossil, it brings the genus down to a much later horizon in our English beds than previously recognized.

it seems to me not unlikely that the putrifactive decomposition of the animal matter of the fish bed, has yielded the principal part of the bitumen. I may here add, by way of parenthesis, that all fossiliferous limestones are more or less bituminous. When a hot sun falls upon certain portions of the Coralline Oolite of Weymouth, the odour of bitumen is strong, just as in the Wady Mohamet, for example, where, according to Lartet and Tristram, the bitumen actually exudes. The only circumstance which causes me to doubt the animal origin of the Nebi Musa bituminous limestone, is that its odour is described as fragrant, which rather points to a vegetable origin. No doubt most of the Lebanon bitumen is due to destructive distillation of vegetable remains in the Nubian sandstone, and much of the bitumen in former times found floating in the Dead Sea, may arise from bituminized wood in Nubian sandstone on the east side dislodged perhaps, by earthquakes. From this source also, M. Lartet would derive the bitumen of Nebi Musa, volatilized by heat through several thousand feet of beds.

As the subject of sulphur, of which we hear so much in connection with the history of the Dead Sea, is closely related to that of asphalt, reference may be made to a short paper by Dr. Fraas respecting some deposits of sulphur recently observed near Jericho.*

Everybody remembers Lot's wife, and all travellers to the Dead Sea have remarked upon the singular salt deposit in the neighbourhood of which the city of Sodom is supposed to have been located. Fig. 6 is a pictorial section, borrowed from M. Lartet, which may serve to give us some idea of the geological structure of the place. First let me say a few words as to its position. It is a hill, about 5 miles long from N. to S., and a mile broad, which occurs at the S.W. corner of the Dead Sea. This is the shallow south bay of the Dead Sea with the "Sègor" sailing on it; immediately over that vessel is seen the narrow channel between the cliffs of the west coast and the low promontory of the Lisan. These cliffs form part of the great Cretaceous escarpment 2,000 feet high, on which Masada is situated. Perspective causes them to appear so low in the section. On the left of the section is seen the north end of

---

* *Quarterly Statement, Palestine Exploration Fund*, October, 1880, p. 246.

FIG. 6.—JEBEL USDOM AND THE SOUTH END OF THE DEAD SEA.

Jebel Usdom.
Grotto.

1, Gypsum; 2, Saline and Gypseous Clay; 3, Rock Salt; 4, Cretaceous beds, constituting the cliffs on the west side of the Dead Sea, lowered by perspective; *m*, Old Lacustrine Deposits of the Dead Sea basin.

Jebel Usdom, close at hand and only 400 feet high. This is divided geologically into three portions, the lowest (Fig. 6. ³) is the rock salt, the thickness of which is variously estimated from 80 to 200 feet. It weathers into numerous pinnacles, some much larger than others, and these pinnacles are Lot's wives. By all accounts it must be rather like a crevassed glacier. Upon this come beds of saline and gypseous clay (Fig. 6. ²), and in this series are regular beds of gypsum (¹). The whole is supposed to be surmounted by chalk-with-flint, like any other of the higher Cretaceous beds in the district.

Such is the isolated ridge of Jebel Usdom, and many are the speculations as to its origin. Some have regarded it as an old deposit of the Dead Sea, or rather of its predecessors at a higher level like the deposits of the Lisan, but its structure is very different to these extremely thin-bedded deposits. I will merely say that M. Lartet concludes it to be part of the Cretaceous series, where the saline and gypsiferous element is in excess, and he points out certain horizons, especially in the upper chalky beds, where there is a considerable development of salt and gypsum, though nothing like what occurs here. Lartet says that the position of the hill from a stratigraphical point of view is also favourable to the notion, whilst we have already seen that the limestone series south of Jerusalem to the borders of the Sinai sandstone is throughout so saline that decently potable water is very scarce. Somehow the salt of the old Cretaceous-nummulitic sea has never been fairly washed out of it. We must conclude therefore, for the present, that Jebel Usdom is merely an exceptionally saline development of the Cretaceous rocks which form the principal part of the rim round the basin of the Dead Sea.

Canon Tristram, I may observe, who formerly advocated the notion of these being old Dead Sea deposits brought up by a swelling of the ground, has lately abandoned this idea, and now places Jebel Usdom at the top of the Nubian Sandstone, which he is disposed to regard as "New Red." This seems to me a most gratuitous conjecture, although there may be evidence in the background, which that author does not adduce. This concludes the section relative to the Cretaceous and Nummulitic limestones.

6. *Beds of marine origin newer than the Nummulitic limestones.*—

Beds with Miocene fossils I have already stated as occurring in the Chalouf cutting of the Suez canal, but whether the whole of the district marked *b* in the map, is to be regarded as Miocene is extremely doubtful. Yet the hollow or depression of the Isthmus of Suez, is, in part, filled in by Miocene beds, and it is probable that since Miocene times, there has been no communication between the Red Sea and the Mediterranean, notwithstanding the very slight water shed. It is said that not a single species is common to the Mediterranean and the Red Sea, whose fauna belongs to the Indian Ocean. The Red Sea fauna extends up to the basin in which the Bitter Lakes are situated; and the Mediterranean fauna is traced southward for some distance, but they never meet. A portion of the intervening ground between Lake Timsah and Serapeum is occupied by freshwater beds, which show that a branch of the Nile, most likely in Quaternary times, flowed through the hollow now occupied by the sweet water canal, past Tel-el-Kebir and Ismailia, where it most probably turned round into the Red Sea. The Etherias of the Nile, now confined to the cataract districts and above, are found in abundance in these deposits near Serapeum. Etheria, I may mention, is a peculiar genus of the family Unionidæ, very characteristic of some of the great African rivers.

Most of the coast deposits in the great bight which lies between Mount Carmel and the delta of the Nile, are regarded by Lartet as Quaternary and recent.

7. *The Volcanic Rocks.*—Under this head I would only include the Tertiary eruptions, interpreting the word Tertiary in its widest sense, as coming down to the present time. Mr. Bauerman concluded that the volcanic rocks of that portion of the Sinai peninsula visited by him were of two periods, of which the second was contemporary with the flint conglomerate.

The twenty-eight *harrahs* of Arabia, in alignment with the Red Sea, the Gulf of Akabah, and the Dead Sea hollow, are remarkable instances of the parallelism of volcanic centres to fissures filled with water. I should mention that a *harrah* means a fire district or volcanic region. One of these, not far from Medinah, erupted

during the reign of the Caliph Omar, and this is the last sign of activity of which we have historical notice.

As we travel northwards, small basaltic flows are noticed in Moab, and these prepare us for the great lava-covered districts which lie to the east and north east of Galilee, whilst many smaller ones are noted in Galilee itself. The upland plain of Damascus seems at one time to have been enveloped in a semicircle of fire, of which remains are to be seen in the three huge fire districts of the Safâh, the Ledja (including the Hauran), and the Jaulan. Jebel Kuleib (5,600 ft.) is the Hill of Bashan, and the country thence to the Jordan contains some barren scoriaceous districts, and other of extreme fertility, where, the basalt in decomposing, mixes with a peculiar loam. This probably nourished the vegetation on which the bulls of Bashan waxed fat. Enormous quantities of magnetic iron ore occur in the hill of Bashan, in fact, no part of the country calls for exploration more than this does.

Many of the basaltic blocks are covered with inscriptions and figures of beasts, &c. Basalt, in fact, seems to have been much in favour as a medium for inscriptions. Take the case of the Moabite stone for instance. Canon Tristram says that it was much in favour with the old Cyclopean builders. Most of the specimens from Moab and Galilee brought for analysis are olivine dolerites, extremely rich in iron, and rather poor in silica.

It might be asked, what is the age of these masses of Basalt? When did the eruptions begin, and when were they brought to a close? As a partial answer to the first question, M. Lartet observes that no basaltic fragments have been found in the older deposits of the Dead Sea basin, and, as these include pebbles towards their margin, it would seem as if the basalt had not reached the surface when the oldest Dead Sea deposits were made. On the other hand, there is no tradition which can plainly be identified with volcanic activity, in those regions where volcanic matter is now found. It is plain that the fate of the cities of the plain was in no way connected with volcanic eruptions, as this is just that portion of Palestine which is most free from anything of the sort. The age of the volcanic eruptions awaits further investigation.

8. *The Jordan Valley and Dead Sea Basin.*—This is the most important of all the sections, and I have been obliged to hurry over some of the others in order to arrive at it. The others being more or less necessary to a correct understanding of this one. By Dead Sea Basin I mean the whole of the longitudinal fissure from the slopes of Hermon to the water-shed near Akabah. The drainage of this finds its way into the Dead Sea, on the north by the Jordan valley, on the south by the Wady Arabah. From the surface of the Dead Sea it is dissipated by evaporation, whereby the level is maintained, and maintained so effectually, that a depth of nearly thirteen hundred feet intervenes between its surface and the level of the Mediterranean, so that the very clouds in this valley are often hundreds of feet below the level of the sea.

Broadly speaking, there can be no doubt of the general connection between the Dead Sea fissure, the Gulf of Akabah, and the Red Sea, right away to the Straits of Babel-Mandeb, and the line of depression is carried north in the hollow of Syria, between Lebanon and Anti-Lebanon, though there above the level of the sea. Such a singular place has been fruitful in theories, and the destruction of the cities of the plain for instance, has led travellers to endeavour to account for the facts of biblical history, by statements too often entirely devoid of truth. The craters of De Sauley, &c., have no existence. And it so happens that the volcanic eruptions are quite in another part of the country, as we have already seen.

There are two theories which claim our attention as not being obviously inconsistent at first sight with physical facts. One is, that previous to the disaster of the Pentapolis, the river Jordan flowed into the Gulf of Akabah; the other is, that the Red Sea came right up over the present water-shed, and filled the great fissure almost to the foot of Hermon. According to the first view, the whole system would be fresh water, according to the second it would be marine.

Before entering on this question generally, we should direct our attention to the geological structure of this fissure basin, in connection with the high grounds of Judæa and Moab. Figure 7 is a diagrammatic section of Palestine from the coast to the mountains of Moab. The Cretaceous limestones rising from beneath the waters

FIG. 7.—GENERALIZED GEOLOGICAL SECTION ACROSS PALESTINE.

o, Level of the Mediterranean; a, beds of the maritime plains; m, old lacustrine deposits of the Dead Sea basin; n, deposits now forming beneath the Dead Sea; p, tuffaceous deposits of hot springs; h, basal.

of the Mediterranean, and the recent deposits in the Vale of Sharon, ascend by a major anticlinal fold to the plateau of the Judæan highlands, whose physical geography we have already studied. The Mount of Olives and the country to the eastward, is occupied by the very highest beds of the Cretaceous series, the soft white chalk-with-flints. On this side of Jerusalem there occurs a preponderance of easterly dips in these limestones, which the various local sections well bear out. The consequence is that the beds come down with an increasing dip, and thus in all probability pass right under the Dead Sea in the manner shown in the section, where they are sharply faulted against the Nubian sandstone, which is seen only on the east side. Here a general westerly dip is said to prevail, at any rate near the Red Sea, in the manner shown, so that if there is no fault, there must be a synclinal curve with arms of very unequal steepness, which would have all the effect of a fault. Whichever of these views we adopt, the chasm must in the main be due to stratigraphical causes, and not to erosion, as in the case of a canon.

I will also note one or two points which this diagram suggests: The deposits marked *a*, are those referred to so briefly in section 6, the raised beaches, blown sands, &c., of the plain of Philistia and the Vale of Sharon; the deposits marked *m*, are old deposits of the previous Dead Sea, to which allusion will presently be made; the deposits marked *n*, are those now forming beneath the Dead Sea itself; the deposits marked *p*, are those produced by hot springs, of which Zara and Calirrhœe are the most remarkable on the east side of the Dead Sea. There is also a hot sulphur spring on the west side near Masada, which no doubt deposits slightly. Tufaceous limestone, sometimes aragonitic, is the principal substance deposited by these hot springs. There are small eruptions of basalt also on the east side of the Dead Sea. The lines which indicate the upcast pipes of the hot springs, are of course somewhat hypothetical, but there can be no doubt whatever that the thermal conditions of the Jordan valley have some connection with this line of fissure, be it actual fault or unequal synclinal.

Having thus far studied what I may term the geological accessories of the Jordan valley, we must now enquire into the nature and

probable age of its deposits, including of course those of the Wady Arabah, and see how far their history bears on the question of the sudden sinking of the Dead Sea basin, which has been supposed by some to have been the cause of the ruin of the five cities.

Now M. Lartet maintains that the great fissure has been an independent basin for a long period, indeed, at one time he was disposed to regard its origin as having dated from Miocene times, but in his latest work *La Mer Morte*, he does not seem disposed to carry it so far back in time. But that it has been an independent basin for an immense period, he considers can be shown in many ways. Firstly, as was noticed by Hitchcock, from the surveys of Robinson and Smith, all the tributaries of the Arabah flow in so as to form an acute angle with the main valley pointing to the north, a very strong physical proof that the drainage of that region has not been reversed. On the other side of the Dead Sea the lateral wadies of the Jordan system for the most part point to the south; all the evidence indicating that the drainage has been towards the Dead Sea in former times just as it is now.

But in spite of these facts, so ably put forward by Hitchcock and Lartet, it must be admitted that the zoological evidence does point to the probability of some connection, doubtless of very high antiquity, with an African continental river system. Canon Tristram, with his usual energy, brought to England numerous specimens of fish from the upper waters of the Jordan system, probably from the lake of Tiberias. Some of these were decidedly of African types. According to Dr. Gunther the following genera occur, viz.: *Chromis*, *Hemichromis*, and *Clarias*. Now *Clarias macracanthus* is a common fish of the upper Nile. But to show the remote nature of the connection, it is a curious circumstance that *Hemichromis* is not represented now-a-days in N.E. Africa, but chiefly in the central African lakes and even in the west coast rivers.

On the whole, Dr. Gunther, who had the examination of Tristram's fishes, demurs to M. Lartet's notion as to the complete independence of this valley system. Nevertheless, an examination of the nature and mode of occurrence of the Jordan valley deposits, as shown both by M. Lartet and Captain Conder, prove the very

considerable antiquity of the valley as an independent basin, and thus show how impossible is the theory which would date its origin from the catastrophe of the Pentapolis.

Captain Conder shows four very distinct margins of old lakes, which were predecessors of the present Dead Sea. The present north shore of the Dead Sea is a shingly beach with a ridge of pebbles at the top of a somewhat steep slope. About thirty feet above the top of this is a similar one. This marks the surface of extinct lake No. 1; a hundred feet higher is a third beach, which marks the surface of extinct lake No. 2. Above these beaches, and some 300ft. above the present Dead Sea, are flat shelves of marl, known as the Siddim level; this marks extinct lake No. 3. And there is yet another still higher. All these lakes would obviously fill up a considerable expanse of the valleys of the Jordan and the Arabah, and their deposits would serve to give us some idea of the history of the period. These lacustrine deposits M. Lartet calls the "deposits of the Lisan," from their great development in that peninsula, which wholly consists of them. They are, as a rule, very thin bedded marls with much salt and gypsum, both in beds and throughout the mass, and have considerable resemblance to deposits now going on at the bottom of the Dead Sea, where crystals of salt and selenite are brought up from the bottom, along with blueish marl. From the absence of organic remains, M. Lartet argues that the physical conditions must have been pretty much the same as now, except probably for the greater quantity of water, and this, M. Lartet considers, is due to the much greater humidity of the atmosphere during Quaternary times.

Anyhow, what we have before us in these Jordan valley deposits, is pretty fair evidence, both on the testimony of Conder and Lartet, that neither the waters of the Red Sea, nor of a fresh water system, have prevailed over that area where these saline and gypsiferous marls have been thrown down, and consequently the existence of these old saline lakes at much higher levels than now is proof of the considerable antiquity of the present depression. Indeed, as Captain Conder remarks, no less than four Dead Seas had dried up before the days of Abraham.

FIG. 8.—SECTION ACROSS THE JORDAN VALLEY.

Figure 8 serves to show the relation of the old lacustrine deposits, which are remarkable for their sterility, to more recent fluvial and torrential deposits, usually of a much coarser nature, brought down and again cut into by the Jordan and its tributary wadies. This secondary valley in which the Jordan runs is known as the Zor, whilst the larger or primary valley is known as the Ghor. These fluvial deposits are often just as fertile as the old saline lacustrine deposits are sterile.

There are many other subjects of interest connected with the Jordan valley, such as the evidence of climatal changes, &c., but we have no time for these. Before concluding I must just allude to the hot saline springs in the Jordan valley. Of these I will only mention three, although Dr. Salah Merrill has pointed out several others. It is clear from the abundance of these springs all along this great line of fault, that the forces of vulcanicity are not yet wholly quiescent, although lava no longer flows from the neighbouring craters of Galilee and Bashan. Indeed, so late as 1837, the inhabitants had a reminder of the fact, in the earthquake which destroyed the city of Tiberias and killed thousands of people.

That there is no necessary connection between Sulphur waters and vulcanicity the evidence of Harrogate will show, but when waters are thermal over a long line of country, we naturally regard hot water as proof of heat somewhere.

There is a remarkable hot spring at Gadara, near the junction of the Jabbok with the Jordan. I mention this particularly, because, according to Dr. Merrill, this place has natural advantages, which will cause it to become the Montpelier of Syria some day. I think he mentions five sulphur sources. There are magnificent views from this place, which was evidently a favourite one in the days of the

Roman empire, from the numerous remains of villas in the environs. It only requires that the detestable Bedoueen should be driven away, for the place to become one of the most charming in the east. Allusion has already been made to the hot springs of Calirrhœe, and to the marvellous scenery of the gorge in which they occur. These are hotter than the Gadara waters, being about 140° F. This also is about the temperature of the hot springs of Emmaus, on the west side of the Lake of Tiberias, which have been made the subject of analysis. There is considerable general analogy between these waters, and those of Harrogate. The water of Emmaus is a highly saline sulphur water, containing bromide of magnesium in very appreciable traces.

We cannot quit this part of the subject without a word about the waters of the Dead Sea.

Well may the Arabs express astonishment when they see the spring floods of the Jordan swallowed up year after year, without any material increase in the size of the lake. To them the notion of evaporation will hardly present itself,—an evaporation which makes this place the recipient of everything that is soluble throughout the valley.

The waters are so bitter and briny, that the fish of the Jordan are pickled at once, and cast up on its shores. Some persons have started the idea that the waters of the Dead Sea owe their salinity to the vicinity of Jebel Usdom, but when we compare the composition of the salts of the Dead Sea waters with those of the Salt Mountain, it is at once perceived that such can not be the case. The rock salt of Jebel Usdom is nearly pure Chloride of Sodium, so that nothing but common salt and gypsum could be obtained from this source. Let us now turn to the composition of the waters of the Dead Sea. The amount of saline matters vary much according to position and depth. Samples from considerable depths contain $\frac{1}{4}$ of their total weight as dissolved solid matter; the salts of magnesium exceed those of any other base, and there is a very considerable proportion of bromide. Sodium, calcium and potassium come next in the order named; all these exist principally as chlorides and sulphates; carbonates are very scarce. There can be no doubt

G. M. Woodward. lith.

West Newman & Co. imp.

Cretaceous fossils, from the Lebanon & Palestine

that the numerous hot springs in the Jordan valley have supplied a large quantity of the saline matter; and we know that the one at Emmaus contains bromine. During the long ages when the four extinct lakes were in existence, their waters were relieved by constant deposits of common salt and gypsum, but so very soluble are the haloid salts of magnesium, that these have not even yet been deposited in any appreciable amount, and, although the present lake yet continues to deposit cubes of salt and lenticular crystals of selenite in the marly mud which is forming at the bottom, it would seem that magnesian salts still remain for the most part undeposited. Thus the present Dead Sea contains the magnesian chlorides and bromides, not only of its modern affluents, but those also of its four progenitors.

The deductions which the chemical geologist is enabled to draw from these facts, are not the least valuable among the many marvels of the great Syrian fissure.

EXPLANATION OF PLATE VII

- Fig. 1.—*Ammonites syriacus*, Von Buch. Lebanon.
- Fig. 1a.—Portion of back of same specimen.
- Fig. 2.—*Nerinæa* cf. *gemmifera*, Coq. Abeih, in the Lebanon.
- Fig. 3.—*Nerinæa* cf. *Coquandiana*, d'Orb. Abeih, in the Lebanon.
- Fig. 4.—*Pholadomya Vignesi*, Lartet. Judæa.
- Fig. 5.—*Procardium*, species. Abeih, in the Lebanon.
- Fig. 6.—*Trigonia scabra*, Lamarek. Abeih, in the Lebanon.
- Fig. 7.—*Exogyra olisiponensis*, Sharpe. Judæa.
- Fig. 8.—*Cidaris glandarius*, Lang. Lebanon or Hermon.