these cases. Now then, a 3 per cent. solution of silver is applicable to eczema intertrigo when located at the following places, namely: Behind the ears, on the penis, and between fingers and toes. Ten to 20 per cent. in the axilla, bend of the arm, under pendulous breasts and labia majora, in the latter event especially when the trouble is due to diabetes (pruritus). Twenty-five to 50 per cent. on the inner surface of the thighs and pubis. On several occasions a .5 to 1 per cent. solution was applied with benefit, in the form of a wet dressing, to a scrotum which was the seat of chronic eczema for months, if not years. The appearance of such a scrotum is quite characteristic; its tissue and size are increased in thickness and extent, and the indurated, rough skin is thrown into itching folds. The application in all these cases can be repeated three or four times during the week; the only time to stop is when an unduly marked hyperemia occurs, as it sometimes does, after the excessive use of the medicinal agent.

In acute and chronic eczema of the face and head, of infants and children, a 3 per cent. solution of silver nitrate is at times very serviceable. But the application need not and ought not to be kept up too long, as the above mentioned resulting hyperemia in some cases will interfere with its speedy and good effects. "Occupation eczema" sometimes can only be cured by the application of a 3 to 5 per cent. solution of silver nitrate. In all these instances the treatment in the interval consists of powdering the parts with bland powders and in keeping them well separated with wadding to prevent further friction.

Parasitic eczema.—Eczema marginatum is a disease which very frequently defies all sorts of applications, but will yield to a strong solution of nitrate of silver. I never began the treatment of marginal eczema with a solution of a strength less than 20 per cent., and very frequently applied a 50 per cent. solution. The raving itching can be successfully combated by this drug, while a complete cure is not a remote possibility.

Herpes tonsurans is the only other parasitic skin disease which can be practicably treated with silver. The application of silver to ring worm is, however, limited. The discoloration the salt produces would be a strong objection to its extensive use on the face, for example. Yet in suitable cases nitrate of silver is a clean and effective remedy. Not very long ago a friend, a young counsellor, visited one of the Turkish baths of the city. A week after, at least this was his affidavit, the right side of his gluteal region became the seat of a good baker's dozen of ring worms. After a few applications of nitrate of silver (30 per cent.) he was completely cured. The discoloration in this case was no ground for objection.

I now come to consider the last condition in which nitrate of silver is in the highest degree the drug. I refer to pruritus ani. The difficulty in managing this symptom can only be appreciated by the multiplicity of suggested therapeutics. It is evident that the proper treatment of a symptom is the removal of the disease. Such, however, is not always the case with itching of the anus. It frequently happens that after curing the hemorrhoids, the fistula, the constipation, the pin worms, etc., the itching will still continue unabated. Sometimes it seems as if it becomes a habit with the anus to itch and the patient to scratch. Besides, there is no denying the fact that the pruritus is often present to an excruciating degree while all

efforts to find its cause end in failure—we then take refuge in that mysterious term, neurosis. While eczema may occur, either localized at the anal region or as an extension and continuation of such a condition existing at the genital organs, yet in a great majority of cases the eczema in the case of pruritus ani is secondary, the result purely of that constant scratching in which all patients take a fiendish delight. In the treatment of this distressing ailment there is nothing in the pharmacopeia better, more effective, more gratifying to the patient and his medical attendant, than the vigorous application of a strong solution of nitrate of silver. In all cases the cause should be sought and when found removed, but if the pruritus still continues, apply a 30 or 40 or 50 per cent. solution, depending on the severity and intensity of the itching, and your reward will be worth the effort. In instances where the pruritus is apparently idiopathic, nitrate of silver in the strength alluded to is a veritable specific. Even where there is a palpable cause, silver will insure rest at night and quiet by day while the cause is being removed. The application should be made once or twice a week; in the interval, if necessary, use zinc salve plain, or with cocain (1 per cent.) in it. The preliminary use of cocain (4 per cent.) before painting the anal region with the solution of silver should never be forgotten, for while it does not obviate the subsequent transitory burning produced by the silver, it still mitigates it while the solution is being applied.

I am at an end of the discussion. I am aware that something is still missing in my paper, and that is an explanation of the *modus operandi* of the nitrate of silver in each and all of the diseases mentioned. But as the *rationale* is not far to seek, and as the physiologic action of the salt is so well understood, it is hardly worth while entering into it in a purely clinical paper such as this.

As a final remark I must state that many of the observations herein recorded were made and treated in private and dispensary practice, and that my entire experience with this valuable drug was rendered possible by my long, useful and grateful assistantship to that distinguished dermatologist, Dr. S. Lustgarten, in his skin clinic of the Mt. Sinai Hospital Dispensary.

156 Clinton Street.

THE PATHOGENESIS OF GOUT.

BY HANS FROELICH, A.M., M.D.

CHICAGO, ILL.

Garrod demonstrated in 1848 with his thread test the presence of uric acid in the blood in gout, and basing himself upon this fact, said that in gout there was reduced capacity of the kidneys to pass uric acid; and second, that the alkalescence of the humors in gout was decreased. Charcot believed in an increased production of uric acid, caused by a functional disturbance of the liver, and Cantani thinks that the uric acid originates in the cartilages themselves, and believes that the uric acid diathesis is the consequence of disturbance in the nutrition of the cartilages, the ligaments and the tendons near the bone. The scientific researches in gout and kindred diseases had for forty years been under the ban of Garrod's discovery; all the immense work done within this span of time has been in vain, of no practical value, because now we know that not only the methods which had been

employed for the discovery of uric acid were so inexact as to render the results almost valueless, but it has been proven that uric acid and the substance which generates it, is found in many other diseases without causing gouty symptoms. Von Jacksch found that in the blood of the normal muscle there is no uric acid, but that it is present in pathologic conditions, as in dyspnetic diseases like pneumonia, nephritis, and constantly in anemia. More and more evidence of this fact has accumulated. It was proven that diet had less influence upon the generation of uric acid than was supposed; it has been demonstrated that uric acid is constantly and considerably increased in leukemia, where it reaches the amount of 4 to 5 grams within twenty-four hours; even where the uric acid has been found not to be increased as such, it is always increased in its proportion to urea. Yet this increase of uric acid in leukemia never produces gout; there is in fact only one case known (Priebram's case) where gout complicated leukemia.

The first great step in discovery of the pathologic facts in this disease was taken by Ebstein, who proved (1882) that the precipitation of urates always is preceded by a necrosis of tissue, a purely local process, thus indorsing Cantani's views. Ebstein found characteristic changes in the kidneys, a shrinking caused by interstitial inflammation and accompanying deposits of urates, mostly biurate of sodium. Litten found that urates sometimes not only fill up the tubuli, but are interspersed within the interstitial tissue, and Dickinson joins him by declaring that precipitation of urates takes place solely within the intertubular fibrous tissue and that this part of the organ, thickened by a kind of inflammation, shrinks, compresses the tubuli, causing thereby the granulated kidneys. Wagner demonstrated that lead poisoning can be at the basis of this inflammation, and give rise to consecutive gout or deposition of urates. Cornil and Ranvier are convinced that in the gouty process of the cartilage the cells take part in the inflammation, and Rindfleisch calls them the very centers of the starlike crystals of urates. Ebstein found that the inflammation and precipitation of urates follow the cartilage fibrils, that is, their connective substance. The necrosis of tissue, according to his view, is secondary to the disturbance of the nutrition, which is the primary phenomenon of gout and must have reached the most advanced degree, must present the symptoms of complete decay before urates precipitate. Concerning the cause of this necrosis Ebstein clings to the view of all others, that it must have been caused by the gouty matter, the fluid uric acid, but he is not able to explain why the uric acid should cause tissue necrosis here and not in leukemia or other diseases which are accompanied by presence of uric acid. The question as it stands today is that we have a tissue necrosis in certain points of predilection, preceding precipitation of urates into these necrotic localities.

The questions we have to solve are:

1. What causes the pathologic appearance of uric acid in the tissue and blood?
2. What causes this necrosis?
3. What is the physiologic treatment of gout, after we have answered these questions?

The first question has been fully answered by Horbaczewsky (1892). His discovery brought us near the solution. He found that in maceration of nuclein-containing tissue outside of the body, which is pro-

duced by a certain degree of putrefaction or through boiling water, there are products split off, which produce uric acid. It is furthermore a fact, that nuclein-containing substances decay within the living body too; one is entitled, therefore, to suppose that the uric acid within the living organism originates from leucocytes.

The fact that in leukemia, where leucocytes decay *en masse*, the production of uric acid is constantly and considerably increased, confirms this supposition.

The further fact, that quinin and atropin, which arrest the formation and decay of leucocytes, reduce the production of uric acid, is another proof.

W. Weintrand (1895) in his investigations upon the influence of the nuclein contained in food upon the formation of uric acid, comes to the following conclusions:

The recent investigations with reliable methods concerning the secretion of uric acid, demonstrate that it is dependent in a high degree upon the metabolism of albumin. It has been found that animal or vegetable food had not the energetic influence which formerly it was supposed to have, even the quantity of the secreted uric acid did not vary according to the increased or decreased quantity of fed albumin.

It became evident that the absolute quantity of the secreted uric acid is individually somewhat varying, but represents a pretty constant quantity eagerly retained by each individual.

This agrees with Horbaczewsky's hypothesis, that uric acid originates from the metabolism, that is, from the decay of leucocytes, and that there is a constant proportion between the secretion of uric acid and the number of leucocytes. The increasing recognition of Horbaczewsky's theory removed the influence of food upon production of uric acid still more to the background. It was thought necessary to adopt this view only so far as the introduction of food produces digestive leucocytosis.

Weintrand demonstrated to a certainty that nuclein-containing food, thymus, brain, etc., is perfectly absorbed and causes a considerable increase of uric acid and its excretion.

The resorbed nuclein influences merely the formation and decay of leucocytes and by their increase indirectly the production of uric acid.

Horbaczewsky says that the number of leucocytes in fasting condition is small (hypoleucocytosis) and increases at once after introduction of food (leucocytosis). A similar parallelism exists if we compare the number of leucocytes of persons of various age, sex, nutrition, etc., with the secretion of uric acid of such persons. Children have more leucocytes than adults and secrete more uric acid. Women have less leucocytes in the blood than men and secrete less uric acid; also well nourished people have more leucocytes in the blood and secrete more uric acid than others do.

Not each rise of temperature causes increased secretion of uric acid, only the character of the fever is of influence (Bartel) as we see in the epicritic increase of uric acid after acute feverish diseases. This inflammatory leucocytosis takes place in croupous pneumonia, when the secretion of uric acid shows a hypernormal value during the fever, still increasing with falling temperature, and reaches its height when temperature is normal, and it increases in the same measure as the number of leucocytes decreases. T. F. Richter also says it is certainly true that this con-

nection between uric acid and leucocytes exists, but we do not yet know the degree of it; we know how much uric acid is secreted, but not how much has been formed in the organism.

According to F. Levison (*Zeitschrift für klin. Med.* 1894, p. 292) uric acid can not, like urea, be considered a metabolic product of albumin introduced through food. The secretion of uric acid in the body can not be considerably increased through introduction of large quantities of easily digested albuminates, neither decreased by hunger diet. It is generated through oxidation of the organized albumin in the body especially the nuclein, and its secretion is increased or decreased by all factors which produce a faster or slower destruction of the cellular elements of the body, especially the leucocytes, like disease, medicines, poisons, etc.

The production of uric acid in the body is in direct proportion to the number of leucocytes used up within that given time. So it is with the well known increase of uric acid of infants, it is directly proportional to the increase of leucocytes. Fleasburg found six hours after birth infarcts of uric acid in 14.3 per cent. of infants; twelve hours after birth in 56.2 per cent.; after twenty hours 94.7 per cent. These oscillations in uric acid are in direct proportion to the amount of leucocytes, which rises the first days after birth and then decreases until the fifth day, when it reaches the point where it has been the last month of fetal life.

Number of leucocytes in the last month of fetus 8,053; immediately after birth 19,600; twenty-four hours after birth 23,000; forty-eight hours after birth 17,500; fifth day after birth 8,500.

Gundoline examined the number of young, ripe (neutrophile) and over-ripe leucocytes and found:

	Leucocytes persq.cm.	Young.	Ripe.	Over- ripe.
Last month of fetal life	8,053	1,013	556	6,458
Right after birth	19,600	4,000	2,352	12,348
Twenty-four hours after birth	23,000	5,520	1,840	15,640
Forty-eight hours after birth	17,500	3,695	1,575	12,250
Fifth day after birth	8,500	2,720	680	5,100
Nursing-age	12,908	7,598	833	4,477

This table shows the rapid transition of leucocytes into the over-ripe form. They can increase in pneumonia to 40-50,000 (Monti, Bergrun), though never in typhus.

Levison doubts the existence of leucocytosis in gout and ascribes the production of uric acid to retention by the kidneys (Garrod, Roberts).

The renewed activity in this whole question brought out a few publications about the metabolic changes in leucemia, which has so much in common with gout and which I report here only with the purpose to more fully exhaust the metabolic percentage of uric acid.

Kuehnau relates statistics of two cases. The leucocytosis in these cases was about constant at the beginning; the daily secretion of uric acid vacillated between 1.5 and 2 grams. This was succeeded by a period of increase of leucocytes which was followed by an increase of the daily secretion of uric acid. After this a rapid decrease of leucocytes set in, while uric acid even increased to 3.39 grams. This certainly gives basis to the apprehension that uric acid is formed through destruction of leucocytes.

The most exhaustive publication about leucemia appeared lately in the *Berl. klin. Woch.* by A. Fränkel. He found the marrow of the bone always diseased, the

spleen only sometimes enlarged. Fränkel's investigations of the blood of acute leucemia are the only ones thus far published. He found increase of the white and decrease of the red corpuscles, the leucemic character of the blood is based exclusively upon an increase of mono-nuclear cells of the character of lymphocytes, which had lost the ameboid motion, the multi-nucleated cells were present only in minimal quantity. The cartilages in some cases swell and are painful, the pain extends over the bone. Two cases which died from septic infection, showed with the appearance of the infection an immense decrease of leucocytes, which Fränkel attributes to leucolysis; with this leucolysis the general condition of the patient grew worse, which was partly due to this leucolysis, partly to sepsis. With the decrease of leucocytes the uric acid increased considerably, not only relatively to the present urea, but absolutely. The secretion of uric acid in leucemia can increase to five or six times the amount of the normal secretion. This publication again demonstrates conclusively that destruction of leucocytes is essential as regards the amount of uric acid secreted.

There are only a few micro-chemic investigations published, but those confirm the facts gained by experiment and investigation on pathologic specimens in regard to the origin of uric acid from leucocytes.

Neusser (*Wiener klin. Woch.*, 1894) found the number of leucocytes always decidedly increased, contradicting Levison, who denies any leucocytosis in uric diathesis. He observed the appearance of particular granules and formations of lumps within the protoplasmatic zone, mostly in the immediate surroundings of the leucocyte nucleus which color black with the basic component of Ehrlich's triacid. They form perinuclear granulations roughly dentated, especially within the mononuclear large and small leucocytes, sometimes in polynuclear neutrophile and even in eosinophile cells, probably a forerunner of uric acid.

Neusser found these changes in gout, muscular rheumatism, asthma nervosum, chronic dermatitis, diabetes, neuralgias, migraine, vertigo and neurasthenia.

L. Heine (*Zeitsch. für phys. Chemie*, xxi, i) says nucleinic acid, unpaired, possesses very peculiar qualities, which disappear more or less in the paired combinations with albumin, and one is justified in thinking that formation and destruction of these combinations is a physiologically important process, which is connected with important functions of the nucleus. Liberfeld's experiments with Ehrlich's coloring demonstrates that in those conditions where a morphologic partition goes on, also a chemic dissolution of the nucleus constituents takes place, a splitting off of nucleic acid from albumin. Heine believes himself to have found that the essence of mitosis does not consist in a chemic dissociation of the nuclein bodies; these substances seem more to preserve their peculiarity as albumin combinations of the nucleic acid, only their physical grouping is modified in mitosis for the purpose of exact bisection.

In the above we have been able to demonstrate that the uric acid in the human body is generated by the decay of leucocytes, which have not reached their full physiologic development, that it is not a weaker oxidation of the albumin of the food, which latter always is oxidized, to urea. This destruction of leucocytes, for which Loevitt gave the name of leucolysis, is always preceded by a leucocytosis. Leu-

colysis is a chemic process which separates nucleinic acid from the albumin.

We proceed now to answer the second question.

What causes the necrosis in gout? The solution of the question as to the generation of uric acid has great practical importance, inasmuch as it tells us that without leucocytosis there is no leucolysis, without leucolysis there is no formation of uric acid, and that only food rich in nuclein increases uric acid. We know furthermore that in many other diseases formation of uric acid takes place within the blood without accompanying necrosis and gouty symptoms, and still the belief in uric acid as *causa peccans* is paramount; no discoveries, no reasoning is potent enough to unsaddle the uric-acid diathesis in the mind. Philosophy used to, and still, plays too big a rôle in our science, as long as it could not lay claim to be an exact science; but the practical results of our physiologic researches upon the recognition of morbid conditions and their treatment, have changed this also. There was so much we did not know that we had to make up for this apparent deficiency with some theory, which used to be labeled with a big name, sufficient to embrace a controlling part of the things which we do not know or know but little. This comprehension of a whole mass of vague ideas and fallacious conceptions, I might say a world of negatives within a seemingly positive name, for the time covered up this chasm in our knowledge, the habit to classify nothingness within something became second nature and we thought we were talking science when we generalized a whole lot of negatives in groups as hysteria, functional nervous diseases, etc. When I, a few years ago (*Med. Mirror*, January, 1893), demonstrated that these functional nervous diseases were built up on the same physiologic and pathologic basis as chronic rheumatism, a few called it ingenious, the great army of authorities and their blind followers shrugged at last their shoulders and ignored it. Yet I had the satisfaction to see that chorea was taken away from the functional nerve diseases and referred to the rheumatic diseases, and only lately Leyden declared that tabes was of peripheral origin, caused probably by the same influences which generate usually rheumatic disorders. The same conditions hold good with that devotion to the uric diathesis, it serves as a covering of the most incongruous morbid states and affiliated diseases which have a pathologic or physiologic relation whatever. Rheumatism and gout are—especially by the English school—thrown together in all possible modification of Greco-Latin nomenclature, rheumatic arthritis, rheumatoid arthritis, gouty rheumatism, goutic rheumatoid, meaning to say something which does not really exist. How grateful am I to Roentgen and his rays, which will establish in most of the cases the absolute absence of any urates! Rheumatism can be following gout as it follows any disintegration of tissue or necrosis through the disturbance of the lymph current (lymphostasis), but that is their whole affinity, with the sole exception of one etiologic fact, the want of exercise and indulgence, too much and too often, in all good palatable things. There the relationship stops. Rheumatism is essentially of delicate reaction upon atmospheric changes, vacillations in the humidity of the air; gout is not. In the long run of the disease we can have atrophy in rheumatism, in gout we have necrosis from the start. In rheumatism the destruction of leucocytes plays hardly any part, in gout it does

most conspicuously. In rheumatism we have agencies, probably ferments, of purely peripheral origin, like cold or auto-intoxication from lymphostasis with their tendency upon the connective tissue to produce fibrinous masses; in gout we have to deal with direct poisons, which destroy the leucocytes and liberate nucleinic acid. This is a fact proven by experience and experiment. Lead is demonstrated by experience to cause the same tissue necrosis with following precipitation of urates as gout. Ebstein produced the same phenomena by injection of chromic acid. We have only one pathologic process, which can rank with that of gout, I mean atheroma of the intima of the blood vessels. There we meet with the same tissue necrosis and following atrophy, only we find instead of urates calcium precipitated into the rough necrotic surface, probably caused by the surplus of calcium in the blood, which in the age when atheroma occurs is no more needed for the building up of the osseous skeleton of the body. Never did a scientist and prominent authority himself utter a more truthful word than Sternberg did when he said that nothing impeded the progress of science more than blind thoughtless following of so-called authorities. Bouchard injected 64 centigrams of uric acid per kilogram of living weight into rabbits, to investigate its toxicity and comes to the conclusion that we can never introduce into the veins of an animal more uric acid than in his first experiment (250 centigrams to 0.641 kilogram), since the doses of uric acid would saturate a quantity of water which of itself is toxic. Yet we meet with recent publications where the toxicity of uric acid in the uric diathesis is the cause of all kinds of auto-intoxications. If asked the cause of this intoxication we get again a big word for an answer, which is meant to say much, says nothing, the leucomains. We have harmless as well as noxious leucomains, but the main excrementitious poison of our intratissue respiration, the carbonic acid, which through strain in the parenchymatous lymph current stays too long in contact with the tissue cell, killing its ameboid function and provoking from the connective tissue in defense of the cell proliferations, never is alluded to.

We have seen in Fraenkel's exquisite observations that in acute leucemia the marrow of the bones is diseased, the epiphyses swollen and painful. The leucocytes do not reach their full ripe stage and fall to destruction without causing tissue necrosis. In gout we do not notice these changes in marrow and epiphyses. We have an acute feverish though a septic inflammation localized within certain places of predilection, cartilages, tendons, etc., where circulation is naturally slow. We must deduct from this phenomenon that in leukemia the undeveloped mononucleated leucocytes, which lost the ameboid motion, and with it their ability of development, decay in a place where the alkalescence of the tissue is so powerful as to neutralize the tissue-destroying ability of those agencies at once, which are liberated through the chemic act of decay. That is after they have passed their breeding places and reached the free current of the blood.

In gout it is quite different. There the destruction takes place in certain places, which show the effect of the chemic action of the destruction which has taken place there in tissue necrosis. As these places have a naturally slower circulation and the plasma there accumulated is limited, we have certainly to accept the consequences of the fact, that

the noxious agencies, which are liberated through the decay of leucocytes, remain in longer contact with the tissue, and as the alkalescence of the present plasma does not suffice to neutralize these agencies, they as a matter of course corrode the tissue, cause necrosis, and the uric acid, which originates at the same time, precipitates upon the rough necrotic surface. What is this noxious agent? It must be an acid, and as only one acid appears in this chemic process, the nuclein acid, it must be this acid. Nuclein acid dissolves living tissue if this is exposed to it. It remains to explain what causes the destruction of leucocytes in gout, why does this destruction take place in those places of predilection and cause that destruction and not within the blood, as it does in leukemia? In leukemia the destruction takes place owing to the innate weakness, the undevelopment of the leucocytes. In gout we meet the same mononucleated leucocytes, though not to the same amount. We can explain their presence through the etiologic fact that the habits of those persons, their irregular meals, taken too often and too much of it at the time, are bound to produce a more or less continuing leucocytosis, the hemopoietic organs are subject to permanent activity, the leucocytes are not given the time for full development and the organs of their production put them out in a decrepid, mononucleated condition. But nevertheless they would not decay before they reach the general current of circulation if they did not meet on their way there an agent which provokes their destruction already within the cartilage, etc. In lead poisoning they come into contact with this mineral poison; in those cases of gout, which are not due to lead poisoning, there must be another poison. What might that be? There is one powerful poison, produced through the intratissue respiration of our body. If the circulation is impeded and accordingly this poison instead of being rushed away is left in longer contact with the tissues and may and probably does cause destruction of leucocytes in gout on their way to the general circulation, that is probably carbonic acid.

Carbonic acid, produced in intratissue respiration, is ejected with certain leucomains from the parenchymatous cell to the surrounding lymph spaces and from there through diffusion or action of endothelium, or both, through the venous capillaries. A stasis within these lymph spaces of course impedes this action, the carbonic acid stays there longer and is allowed to exercise its toxic influence upon everything it comes into contact with, especially decrepid leucocytes. That in goutic patients we have lymphostasis owing to their want of exercise and overfeeding is certain, and can be demonstrated if the lymph current is accelerated through a laxative, which causes at once a mitigation of all symptoms of the gouty attack, and that the gouty attack takes place where there is already a natural disposition to lymphostasis is another affirming factor.

Intratissue respiration is without any doubt the most potent factor for preservation of the physiologic integrity of all our organs. A. Poehl (*Zeitschrift für klin. Med.*, 1894) of late made it the object of his investigations in connection with the effect of organic extracts, which he says, especially spermin, cause in dissolved condition, intratissue respiration, acting like a ferment. The products of regressive metabolism, the leucomains, he continues, are those nitrogen-holding products of regressive metamorphosis, which are oxidized into so-called intermediary products

instead of urea. The accumulation of these leucomains causes autointoxication, and the most common kind of autointoxication he declares is caused by reduced tissue breathing. Spermin, a nuclein found in testicles, ovaries, etc., has oxidizing power, but only in dissolved form; in presence of phosphates it turns insoluble, inactive. (Charcot-Leyden crystals.) In tissue which is passed by normal alkaline blood no insoluble phosphates can form, reduced alkalescence of blood is the essential moment of inactivity of spermin and its formation can be explained as follows:

Any nerve or muscle tissue if irritated decreases its alkalescence and can be brought to acid reaction, and as phosphates always are present the moments of spermin-phosphate formation with acid reaction are given. Under normal conditions no inactive spermin phosphates are found, because the passing normal alkaline blood prevents it. Reduction of the alkalescence of the blood favors its formation and continuing irritation of tissues can reduce the alkalinity of the passing blood, which again causes formation of inactive spermin. He thinks urea is a special product of aerobic tissue life and goes parallel with intra-organic oxidation and explains the vacillations of the production of uric acid, that it depends upon the circumstance whether the destruction of the nucleins proceeds in a more or less alkaline or acid medium. If this destruction takes place in alkaline medium the spermin will remain active and the oxidation to urea goes on normally, but if it takes place in an acid or reduced alkaline medium, then through splitting off of phosphoric acid insoluble spermin phosphate is found, which reduces the oxidation process and uric acid accumulates.

This is a very captivating and ingenious theory, only it does not hold good in gout. It may be the cause of neurasthenic troubles, but even then it would not explain the long duration of such chronic ailments, even if the formation of insoluble spermin were the cause the current of the blood would in less than no time remove it and replace alkaline reaction, and even if lymphostasis should be present enough alkaline plasma would pass through to reestablish the normal. Uric acid is not in gout increased through retention, but through hypoleucocytosis, which followed by leucocytosis produces first decrease, afterward excess of uric acid, not only relatively to the present urea, but absolutely. Uric acid, furthermore, is generated through decay of leucocytes and not a lower oxidation of food, which always is oxidized to urea. Another very important factor against Poehl's theory is the circumstance that the curative effect of spermin in those ailments, which he praises so highly because of intense and rapid effectiveness, fail to give other investigators the desired and promised results. We therefore sum up our answer to question No. 2 as follows:

The tissue necrosis in gout is caused by a poison, probably nuclein acid, sometimes lead; the uric acid in gout is generated by decay of leucocytes, which destruction is brought on probably through carbonic acid. The reduced amount of uric acid in the beginning of the gout attack is parallel with hypoleucocytosis, the following increase with leucocytosis.

Treatment.—If ever we succeed in establishing the true physiologico-pathologic basis of a disease, it will naturally be followed by the indications of a real physiologic treatment, and the adoption of the latter will culminate in its success in praxis. So it is here. The

treatment which I have built up upon the above facts has proved superior to any other and a success throughout.

Prevention.—We have to convince the patient of the absolute necessity to regulate his habits. Two light meals and one substantial one every day, almost to the minute, at the same hour. Never eat so much that you could not again sit down to the table and eat more. No cheese or eggs, no brains, sweetbreads, liver or kidneys; light wine, if any alcoholics. This will reestablish regularity in the periodic leucocytosis, and regularity in this act will again strengthen the hemopoetic organs and enable them to turn out vital leucocytes. Passive and active exercise sufficient to agreeably fatigue the body, but never so much as to provoke exhaustion, this is especially to be avoided. Massage daily, not by a so-called professional masseur, but by a physician who makes this treatment a specialty. The professional masseur gives everybody the same general, more or less superficial, massage. This is good enough for a business man who substitutes this treatment for exercise, but it will never do where intratissue respiration has to be influenced. I do not mention this from personal motives, *pro domo*, because I limit my practice to this kind of work; I had to do it because I did not find satisfaction with the work of these masseurs. Especially the powerful muscle layers of the body have to be worked and emptied of their lymph down to the periosteum of the underlying skeleton, and in gout especially, those parts have to be well taken care of, which, on account of their retarded circulation are predisposed to localization of gout.

Alkaline mineral waters, if recommended, ought to be taken hot. The only effect of the lithia waters consists in their being substituted for alcoholic drinks, which still is a fact in their favor. I only mention here an experience I had with a patient whom I sent to Hot Springs, Ark. She suffered from rheumatic hemiplegia, but no one suspected her of being subject to lithiasis. Whenever she went there and drank the hot water, for the first few days she used to evacuate immense quantities of gravel. She drank for months lithia waters, *ut aliquid fiat*, without any effect. I ascribe this effect to the small proportion of alkali in those springs, which, especially if taken hot, approach more the real physiologic conditions. I recommend douches alternately with hot or cold water (Scotch douches).

Treatment of the attack.—We have to treat the attack locally, because gout is a local disturbance caused by poisons suspended in the blood, and generally to eliminate these poisons from the blood.

The general treatment has to be directed toward the change of the reduced alkalinescence of the blood and strengthening of the blood corpuscles. Adherents of the old school of course would forbid anything acid, fruits, salad, etc., on account of the alkalinescence of the blood. I am only glad in the interest of the welfare of these patients and of the promulgation of the principles of digestion, that they will never succeed in keeping the stomach from producing hydrochloric acid, they may envelop humanity in as much sweetness as the please. To strengthen the blood corpuscles we have in diluted hydrochloric acid a very good medium, it is an agreeable drink and can be taken oftener during the day. To remove the poison from the blood, we have only one way, that is a good old-fashioned bleeding, which may be repeated according

to the condition of the patient, especially if lead poisoning is apparent. Bloodletting should not be limited to the pocketbooks of our patients, it is a harmless remedy, as the removed blood is restored in a few days, and is the most powerful means of changing the condition and composition of the blood by its enormous influence upon the hematopoetic organs to more energetic and more extensive activity. The experience of our grandfathers, which praised this proceeding even in those persistent cases of chlorosis in apparently well nourished individuals, we know now to be based upon the fact that the bloodletting removes the numerous decrepid mononucleated cells and replaces them with more vigorous corpuscles. I can not make a plea strong enough for rehabilitation of this remedy, the interdiction against which I contemplate as one of those modern windmills, which have to be fought. Of course its use has to be left to the discretion of the physician.

To increase the alkalinescence of the blood we might apply the lymphagoga, which Heidenhain introduced. The extracts of crab-muscles and leech heads seem to be varying in their efficacy.

Of all the lymphagogas cinnamon acid is the most reliable, the most rapid and the most harmless, but as the physiologic effect of all these medicines is connected with and based upon leucocytosis, I preferred to do without it, as just in gout too often repeated leucocytosis seems to be the main cause of the deficient development of leucocytes. Still, in cases where the kidneys or other organs are effected, which we can not reach by local treatment, lymphagogas could be the only way to increase the blood's alkalinescence at once.

The local treatment is the most important, and in its effects the most effective. We are accustomed to regard a surgical treatment of the local precipitation of urates, the tophi almost as sacrilege. They play the part of foreign bodies in our system, and each foreign body, if accessible, ought to be removed and we should thank Heaven in the interest of our patients, if there are any surgical means to treat their ailments. There is so much talked and written about dissolving these urates with lithium salts, piperazin and of late lysidri. They work very nicely in the reagent tube with water, but as soon as we take urine, the dissolving effect stops. How it works in plasma, I don't know; if these agencies are locally injected in a concentration strong enough to take effect, that the resulting chemic process causes a painful inflammation, which lasts almost two days. The only way to remove these tophi is curetting under asepsis, it ought certainly to be done in each case during the time no acute attack is apparent.

Another way to cut off an attack is the neutralization of the nucleic acid with a very diluted solution of bicarbonate of sodium. It is injected peripherally from the inflamed section hypodermically, and tenderly massaged over the whole part. It can be preceded by an injection of cocain. My experience is that this local treatment is really abortion, it cuts off the attack within a day or two. Connected with these locally intended alkaline injections the same solution might be injected in larger quantities underneath the skin, certainly the best way to influence the alkalinescence of the blood.

As a matter of course the free action of the bowels is the first thing to be attended to in an acute attack of gout, it has a sudden effect upon the lymph motion

and through this accelerates the access of alkaline blood to the inflamed districts.

Our views of disease at present are in a stage of transition, the inherited slab bones are found to not fit anymore and if several diseases are discovered to present one identical pathologic phenomenon, they are thrown together heterogeneously as they might be. Loewitt observed what he calls leukopenia in rabbits with rapid sinking of temperature as soon as they were exposed to shock; another repeats the same experiment on turkeys, with the same results, finds leukopenia in neurasthenia (a totally arbitrary classification of morbid symptoms, which accompany many entirely different diseases) and says that leukopenia is the cause of neurasthenia, because we notice the same morbid fears there. Since excessive formation of uric acid has been discovered in many other diseases besides gout, Haig and others subject almost every chronic ailment to the uric diathesis, it has seemed to me therefore necessary to show that uric acid has no toxin effect and that its presence has directly nothing to do with the apparent disease. We have to go back to the generation of uric acid, to the poisonous noxes, which create uric acid, and there we find that gout is a disease *per se*, it has nothing to do with rheumatism or other diseases, we are able to distinguish it clearly from that whole mass of confused conception and substitute for its treatment a really physiologic basis.

544 N. Wells Street.

THE USE OF THE CURETTE IN THE TREATMENT OF ENDOMETRITIS.

BY C. L. BONNIFIELD, M.D.

CINCINNATI, OHIO.

Endometritis, for the purpose of discussing the use of the curette in its treatment, is best divided into three varieties—catarrhal, septic and gonorrheal.

Catarrhal endometritis is the form of pelvic disease most frequently induced by improper habits of life, consequently the one most amenable to constitutional or medical treatment.

A large percentage of cases occur in unmarried women. In the society girl it is produced by the nightly exposure of the arms and a large portion of the chest, whereby the blood is driven into the interior of the body, congesting the uterus along with other organs; the tight corset, displacing the abdominal organs and interfering with the action of the heart and lungs; the numerous meals at irregular hours of stimulating and indigestible food, the position assumed in dancing and the exposure and lack of rest during menstruation.

In her less fortunate sisters, the school teacher and the shop girl, the constant standing serves as an efficient exciting cause.

Another cause, I deem of importance, is unsatisfied sexual desire. Genito-urinary surgeons tell us this is an important factor in the production of hypertrophy of the prostate, which is a condition in the male somewhat analogous to fibroid tumors of the uterus in the female. We all recognize the fact that this latter affection is a penalty frequently paid for celibacy. If it produce this graver result of persistent uterine hyperemia it is fair to suppose that it will also produce catarrhal endometritis.

The cause of the disease having been recognized, the proper treatment suggests itself. The faulty habits must be, as far as possible, corrected. If the patient

is anemic, iron, quinin and strychnin between, hydrastis and hyoscyamus or cannibis indica during the menstrual periods. An occasional mercurial purge is always indicated. While the symptoms are acute and exercise gives rise to pain, rest in bed should be enforced; later the patient's mind should be diverted from herself and her disease and healthful exercise in the open air, horseback riding, bicycling, etc., are of the greatest value. A change of climate, particularly a residence for some time at mineral springs, is beneficial. These measures faithfully used will cure many cases, and they should always be employed before resorting to any local treatment in the unmarried. I wish to protest against the growing tendency to curette every girl that has a little leucorrhea and menstruates a day too long or with some pain.

If these measures fail the curette is indicated. It is much to be preferred to prolonged office treatment.

In married women the objection to local treatment does not exist, and in the majority of well-marked cases curettement is the appropriate measure.

It is true that a large number of this class of cases can be cured by office treatment. Drainage of the uterus secured by gauze, application of carbolic acid and iodine to the endometrium, bleeding the cervix, correction of mal-positions, etc., will in time effect cure, and there are cases where it is advisable to proceed in this way. Where existing disease of the heart or kidneys makes it unsafe to use an anesthetic, or where the patient will not readily submit to an operation, the gynecologist may prefer this method of treatment. To the general practitioner who desires to treat his own cases it will usually commend itself, for lack of operative experience makes an aseptic technique well nigh impossible for him to attain, and without there is some danger in the operation.

The severer forms of the disease can not be cured without curettement. Nor will curettement alone cure all of them, but curettement combined with such other local and constitutional measures as are indicated will.

It may be necessary to repair a lacerated cervix, or perform an anterior or posterior colporrhaphy, or both. Or an Alexander operation or suspension of the uterus may have to be done, for unless the circulation in the uterus can be restored to its normal condition, no relief of the symptoms will be permanent.

Septic endometritis usually follows abortion or labor at term, and this phase of the subject will be discussed by my friend, Dr. Mitchell; but it may follow the introduction of an unclean sound or operation upon the cervix; it may also develop during menstruation by infection from germs existing in the vagina. The curette with antiseptic irrigation should be used as soon as the diagnosis is made. Where there are stitches in the cervix they should be removed. If this treatment is instituted sufficiently early, it will generally give relief and obviate the necessity of later resorting to more dangerous surgery, that, even if successful, leaves the patient mutilated.

Gonorrheal endometritis when seen in its incipency can usually be cured by intrauterine irrigation and gauze packing, provided the disease be vigorously attacked at the same time, wherever else it may have found a lodging place, in the vagina, the urethra, the glands of Bartholin.

But if it has been in progress some time when it