loss, nor even to defect, of sight, irritation by tumour in any part of either may lead to inflammation of the optic nerves, on which inflammation blindness usually, not always, follow.—*Brit. Med. Journ.*, Oct. 14, 1870.

60. *Parenchymatous Keratitis associated with Acute Rheumatism.*—Mr. SPENCER WATSON read before the Clinical Society of London (Oct. 14) the reports of four cases of this. The patients were all young adults, all under 19 years of age. Both corneæ were affected, and the symptoms and appearances were not distinguishable from those of parenchymatous keratitis due to inherited syphilis. Out of a series of cases, extending over the last ten years, Mr. Watson had only observed acute rheumatism in association with parenchymatous keratitis in these four; but he suggested that in some others of the series this complication may have itself cut short the observations, by preventing the attendance of the patients at the Ophthalmic Hospital. From these cases it was conjectured that the interstitial keratitis of inherited syphilis may be allied to rheumatic inflammation, and may possibly yield to treatment such as is beneficial in the latter disease. Two of the patients, whose cases were related, were exhibited to the Society. In one, the attack of keratitis had occurred eight years ago. It was found in this case that the young woman had become myopic since, and most probably in consequence of the attack.—*Brit. Med. Journ.*, Oct. 22, 1870.

MIDWIFERY.

61. *Application of Pressure to the Uterus in Cases of Lingering Labour.*—Von Ritgen, in 1856, suggested the employment of external pressure on the uterus as an adjuvant in cases of powerless labour, and the suggestion was put into practice by Kristeller, in 1867, and he published a number of cases in which he had advantageously resorted to it.

Dr. W. S. PLAYFAIR in an instructive article (*Lancet*, Oct. 1, 1870) observes "It is certain that the advantages to be derived from external pressure are not yet widely known or recognized; and as I have now received very material assistance from it in many cases of lingering and powerless labour, I believe it may not be without interest to state briefly the result of my experience on this point, especially as I do not know of any published cases in this country in which its use has been described.

"The class of cases in which external pressure is likely to prove serviceable is of very frequent occurrence, viz., in which the presentation is natural, and the pelvis roomy, but in which delivery is retarded simply from deficiency or absence of uterine contraction. These are the cases in which resort to the forceps is so often essential, in which the head has passed well into the pelvis, possibly descended as low as the perineum, and in which apparently but one or two good pains are required to complete the delivery.

"Firm pressure, applied under such circumstances, may act in two ways: First, and most commonly, it may merely stimulate the sluggish uterus to increased exertion, just as firm pressure after delivery will cause a relaxed uterus to contract. In this way, pains that are feeble and ineffective may be rendered strong and useful, and a natural termination may result when artificial assistance might otherwise be required. I have of late been frequently in the habit of thus stimulating the uterus, and I feel certain that I have in many instances greatly shortened the progress of a labour that threatened to be long and tedious. It is, indeed, often curious to observe how rapidly the pains increase in force and duration, under the stimulation of gentle and steady pressure at the commencement of each pain." * * * "I need hardly add, by way of caution, that gentle but firm pressure in a proper direction is to be used and that all rough handling of the uterus is to be avoided. The pressure can be most readily applied with the patient lying on her back, but this is by