The persistent false statements in regard to Tuffier's deaths would be amusing if it were not that they so outrageously falsified his experience.

OPERATOR.	ANALGESIA.			
	No of Cases.	Successful.	Partial Success.	Failures.
Anderson, C. H., Mississippi Valley Med. Association, Oct. 9-11, 1900.....	6	6		
Bogart, J. B., N. Y. Academy Medicine.....	25	18		7
Boldt, H. J., N. Y. Med. Jour., Nov. 3, 1900.....	1	1		1
Bainbridge, Medical Record, Dec. 22, 1900.....	12	11		1
Cartledge, A. M., Southern Surgical and Gynecological Association, Nov. 13, 1900.....	8	8		
Coe, H. C., N. Y. Acad. Med., Oct. 25, 1900.....	1	1		
Dudley, A. P., N. Y. Acad. Med., Oct. 25, 1900.....	6	6		
Fowler, Medical Record, Dec. 22, 1900.....	81	73	8	
Goldan, S. O., N. Y. Acad. Med., Oct. 25, 1900; N. Y. Med. Jour., Dec. 22, 1900.....	31	25	2	4
Grandin, E. H., Medical Record, Dec. 22, 1900.....	3			3
Gibney, V. P., N. Y. Med. Rec., Dec. 22, 1900.....	5	5		
Hartmann, F. S., Cook Co. Hosp., Chicago.....	1	*1		
Lee, E. W., Phila. Med. Jour., Nov. 3, 1900.....	7	7		
Lyman, J. V. R., Inter County Med. Society, Nov., 1900—Eau Claire, Wis.....	10	10		
La Place E., Phila. Med. Jour., Nov. 3, 1900.....	1	1		
Marx, S., Medical Record, Oct. 6, 1900.....	23	23		
Matas, R., Phila. Med. Jour., Nov. 3, 1900.....	9	9		
McDonald, W. G., Southern Surgical and Gynecological Assn., Atlanta, Nov. 3, 1900.....	1		1	
MacMonagle, B., Southern Surgical and Gynecological Association, Atlanta, Nov. 3, 1900.....	2	**2		
Mallet, E. P., Southern Surgical and Gynecological Association, Atlanta, Nov. 3, 1900.....	2	1	1	
Morton, A. W., Pacific Med. Jour., Nov. and Dec., 1900.....	31	31		
Murphy, J. B., Chicago Clinic, Nov., 1900.....	40	38	1	1
Neff, J. M., Personal Correspondence.....	2	2		
Kammerer, N. Y. Med. Record, Dec. 22, 1900.....	40	40		
Pitesci Racoviceanu, JOURNAL A. M. A., Oct. 20, 1900.....	125	121		4
Podesta, B., Semana Medica, Buena Ayres, Aug. 23, 1900.....	21	21		
Rodman, W. L., Southern Surgical and Gyn. Assn., Atlanta, Nov. 13, 1900.....	3	3		
Weber, C., Cook County Hosp., Chicago.....	3	3		
Tuffier, Semaine Medicale, May 16, 1900.....	130	130		
Total.....	631	596	14	21

* Eucaïn.

** Dr. Barbat.

RHEUMATIC DISEASES OF THE EYE.

H. W. WOODRUFF, M.D.

Professor of Ophthalmology in the Chicago Eye, Ear, Nose and Throat College, and Surgeon at the Illinois Charitable Eye and Ear Infirmary.

JOLIET, ILL.

That the eye should be subject to rheumatic disease does not seem strange when we remember that its principal protective coat is a dense fibrous structure, the sclerotic; that it contains within it the delicate muscular fibers of the iris and ciliary body, and externally the muscular bands which move it about; that it is very vascular, one coat alone, the choroid, being largely made up of blood-vessels. The wonder is that the eye does not suffer more from a disease as prevalent as rheumatism and one which shows such a predilection for fibrous, muscular, and vascular structures.

It is not the object of this paper to discuss the subject of rheumatism in general. Suffice it to say that many observers believe it to be, like gout, caused by uric acid, a product of inefficient proteid metabolism; others by lactic acid, a fermentation product of the hydrocarbons. The researches of Singer, Achalmé, Thirolox and Schüler would lead us to consider it to be a microbic disease. Reinhard believes the throat to be the common infection atrium.

Neither the eyeball nor its appendages are often affected by articular rheumatism. Iritis rarely occurs with it, but cases of conjunctival congestion without mucopurulent discharge have been noticed. We are

compelled, however, to recognize the use of the term "rheumatism" as applied to an extensive group of symptoms which are probably dependent on the same causes with articular rheumatism. Sudden exposure to cold and overmuscular exertion are the principal exciting causes. Some of the diseases of the eye ascribed to the chronic type of rheumatism are iritis, episcleritis, scleritis, keratitis, orbital cellulitis, optic neuritis, chorioiditis, ocular palsy, glaucoma, and opacity of the vitreous. The most important of these in point of frequency is iritis.

It has been computed by various writers of authority that iritis furnishes from 2.3 to 4 per cent. of all ophthalmic cases, and that syphilis and rheumatism are causative factors of this disease in 90 per cent. in the proportion of syphilis 60 and rheumatism 30. It was formerly taught that the etiology of iritis could be determined by simple inspection, and it is true that in many cases the papules or condylomata characteristic of syphilitic iritis can be made out.

In the absence of these masses in the iris, however, the diagnosis depends on the history of the individual case. In the absence of a syphilitic history or signs, we may find a personal or family history of acute or chronic articular rheumatism, symptoms of lumbago, neuralgia, sciatica, torticollis or pains in the articulations or fascia or sensitiveness to changes in the weather. Many cases are subject to relapses of iritis in spring and winter, seasons of the year when rheumatic affections are most prevalent. Some cases of iritis alternate with rheumatism in other parts and some recur with the swelling of the joints.

I will briefly report three cases representing, respectively, iritis, episcleritis, and scleritis of undoubted rheumatic origin.

A woman, aged 25, came to the clinic at the Illinois Eye and Ear Infirmary, in February, 1899. She gave a history of chronic articular rheumatism. Her father was a chronic rheumatic. A diagnosis of rheumatic iritis was made. There was present the typical sign of iritis: ciliary injection, pain in the temple, slight adhesions of the iris to the lens capsule, which yielded to atropin. No nodules were present in the iris. No history nor signs of syphilis, either inherited or acquired were evident.

The patient made a rapid recovery under salicylates internally and atropin in the eye and hot applications. She was free from trouble until last April, when she presented herself again after having suffered pain in the eye for two weeks. The same treatment was continued and improvement again occurred until the early part of this month, then, with a sudden change in the weather, came a relapse of the rheumatism and also a violent pain in the eye. This gradually yielded to hot baths and pilocarpin and leeches applied to the temple; but the exudate from the iris was more abundant and plastic, and adhesions more numerous and permanent.

Next in frequency occurs a superficial form of scleritis, episcleritis, which consists of a circumscribed inflammatory nodule, generally on the temporal side. It is not movable over the sclera, but firmly attached and of a reddish-violet color. This condition can never be looked on as merely a local disease, but must be regarded as the manifestation in the eye of some systemic derangement. The large percentage, if not all of them, are associated with the rheumatic diathesis.

A case illustrating this condition was recently under treatment at the Chicago Eye, Ear, Nose and Throat College. The patient, a man aged 40, had suffered from

articular rheumatism for many years. He had had at least five attacks of episcleritis, and each had left small scleral scars with conjunctiva attached. There was a general chronic nodular condition of the sclera a few millimeters from the corneal margin. Atropin and massage had been used, and salicylates internally, but local blood-letting with the artificial leech gave most relief.

The most serious, but fortunately rare, rheumatic affection of the eye is deep scleritis. Here the circumcorneal congestion is more general, showing extensive bluish-red discoloration. It is a chronic disease and destructive to vision by the formation of deposits in the cornea, and staphyloma of the sclera. I showed a case of this type before the Chicago Ophthalmological Society last December. It was that of a young woman of 29, with a history of hip-joint disease at 3 years of age. Since that time her general health had always been good with the exception of rheumatic pains in the joints, always aggravated during changes in the weather. The family history was negative, and there were no evidences of syphilis, except the interstitial character of the corneal deposit. There was violent scleral congestion with triangular corneal opacities, which the magnified cornea showed to be quite vascular. This condition had existed over a year. A period of five weeks was the longest without a relapse. She was treated with potassium iodid, salicylates, salines, and mercury by inunction; locally with atropin, and hot applications and massage. The iris was slightly involved. In this case again leeches gave instant relief during the acute exacerbations.

I have selected these three cases because they represent, in the first two, the most common rheumatic eye disturbances, and the third the most serious in its consequences. Note particularly the history of many relapses; in all a positive rheumatic condition, in one also a family history of it. The case of iritis yielded best to treatment, but so far is not permanent. The two scleral cases are hardly influenced at all except by heat and local blood-letting.

OVULATION AND MENSTRUATION NOT INTERDEPENDENT FUNCTIONS.*

C. C. THAYER, M.D.

CLIFTON SPRINGS, N. Y.

From one of the most popular text-books¹ on midwifery in our medical colleges, I quote the following: "The intimate connection between ovulation and menstruation is now admitted by most physiologists, and it is held that the determining cause of the discharge is the periodic maturation of the Graafian follicles.

and when the ovaries are removed by operation, menstruation does not take place." One of our latest physiologies, also popular as a text-book² in our medical colleges, states the following: "Women do not menstruate when both ovaries have been removed by operation. Some instances have been recently recorded, indeed, of a sanguineous discharge, occurring periodically from the vagina after both ovaries have been removed for disease; and it has been inferred from this that menstruation is a function independent from the ovaries, but this evidence is not conclusive."

Most of us have been educated along the line of the ovula theory of menstruation, and probably most phy-

sicians to-day believe that ovulation and menstruation are one and the same thing, or at least, duplicate features of a single function. Another view, claiming a respectable and growing minority, divides the thought of the medical profession, and these views are fairly set forth in Gould's "American Year-Book." In an article by Hirst and Dorland, we read: "The divergent views are, first, that ovulation occurs at each menstrual period, and that the two processes are due to the same active cause. The other view is, that ovulation and menstruation are independent of each other, and are due to independent stimuli, and are coincident only by accident."

Since the days of Alexander's "Philosophy of Menstruation," Hamburg, 1841, followed by Grafft, Berlin, 1842, revised and improved by Raciborski, "Habit of Menstruation," Paris, 1868, many others have duplicated their noble efforts, and a clearer and truer understanding and comprehension of the functions of ovulation and menstruation have been gradually introduced into medical thought and literature, so that now many physiologists and gynecologists are concurring in the theory that ovulation and menstruation are naturally coincident, usually concurrent, exceptionally, independent functions. Let us consider this subject from three different standpoints.

1. *Ovulation without Menstruation.*—In Kirke's "Physiology," we read: "Cases are known where ova have been discharged in amenorrhoeic women. It must, therefore, be admitted that menstruation is not dependent on the maturation and discharge of ova." This statement is in harmony with the frequent observations of many gynecologists. Frequently mothers consult their physicians in regard to their daughters, in age ranging from 14 to 18, well-developed girls, who are having regular periodic ovarian disturbances, as regular and complete, minus the menstrual flow, as their mothers did. Perhaps this does not prove that these ovarian disturbances were ovulation; yet had these symptoms been attended with the customary discharge, ovulation would have no longer been considered hypothetical. While we have in the young, substantial proof as to when the menstrual flow began, we have no substantial proof as to when ovulation began.

We have had many patients who have menstruated for years and suddenly became amenorrhoeic, yet monthly had their regular periodic ovarian disturbances, less the external manifestations, and afterward when the flow reappeared it was when they were having these periodic attacks, suggesting at least that the ovaries had been performing their functions, though the uterus had failed to respond with its associated function.

In the year 1881, a young lady came under my care, who at the age of 21 attended the National Centennial, at Philadelphia, and was menstruating at that time. She took cold; the catamenia ceased and she saw no more of it for five successive years, though she experienced for about a year her regular, periodic ovarian disturbances, until gradually they ceased with her failing health. During the fifth year and after about six months' treatment, she noticed the return of the regular ovulation symptoms, and after three or four months more the regular menstrual flow appeared.

During a recent visit to Florida I was consulted by a well-developed, hearty young lady of 18 years, who gave the following history of herself: "About five years ago I began to have bad feelings in my head, back and sides, reaching through the bowels, which I could not account for. They seemed to come on about every four

*Presented to the Section on Obstetrics and Diseases of Women, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.