latter act as any alkaline medicament in emulsifying fatty foods and rendering them easy of digestion. From internal antiseptics most beneficial results have been obtained, the remedy most commended being benzoyl-guaiacol given in capsules, commencing with a moderate dose of a grain or two and gradually increasing it to ten or fifteen grains thrice daily. This drug apparently stimulates the nerve-centres; it increases the appetite; it lowers the specific gravity of the urine; it controls the excretion of urine; and, lastly, it modifies the sugar-formation, thus lessening the amount excreted. In addition, the use of lithium carbonate and Fowler's solution gives the best results.—*American Medico-Surgical Bulletin*, 1895, No. 23, p. 1413.

CONCERNING DIAPHORETIC METHODS.

DR. KARL DEHIO prefers the hot-air bath, administered in bed, in the manner familiar in American hospitals, because it brings about: (1) Extensive change in the circulation and distribution of the blood with lowering of blood-pressure. (2) There is an energetic abstraction of water through increase of perspiration. (3) There is a marked increase of tissue-change and increased expenditure of body-substance. The indications for its use are anasarca from renal disease, either post-scarlatinal or acute, acute anæmia, and chlorosis.—*St. Petersburger medicinische Wochenschrift*, 1895, No. 44, S. 373.

PHENACETIN-POISONING.

DR. G. KRÖNIG reports a fatal case. For occipital headache a seventeen-year-old printer's assistant received five fifteen-grain phenacetin-powders, not more than two to be used in a day. After an evening-dose vomiting commenced, then were noticed great weakness and a bluish-gray coloration of face and lips. The temperature was 102.2° F., the pupils were of medium size, the pulse weak, and the patient complained of headache, vomiting, and diarrhœa. The urine was of chocolate color, the conjunctivæ slightly jaundiced. General icterus followed, and cyanosis of lips, ears, hands, and feet. Urine obtained by catheter was thick, dark reddish-brown in color, containing masses of almost pure blood. Death followed two days after the ingestion of the remedy. As the patient was septic from a purulent otitis, a necropsy was necessary to show that death was due to the drug. The diagnosis reached by this means was stated as universal methæmoglobinæmia, although the patient within the preceding three weeks had taken four similar doses with no apparent ill effects.—*Berliner klinische Wochenschrift*, 1895, No. 46, S. 998.

THE HÆMOSTATIC PROPERTIES OF ANTIPYRIN.

DR. H. HUCHARD notes that as early as 1884 these properties had been noted, and it was suggested by Arduin that the drug might be used for the same purposes as iron perchloride and ergot. Its hæmostatic effects are powerful and rapid. In epistaxis a simple tampon moistened in a solution of the drug (1 to 5 or 1 to 2) is usually sufficient. In more extensive hæmorrhages, hot irrigations with a 4 per cent. solution or even insufflations of the powdered drug are useful. For hæmorrhages after ablation of polypi or ex-