This article was downloaded by: [UQ Library]

On: 15 March 2015, At: 02:34

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK

Annals and Magazine of Natural History: Series 4

Publication details, including instructions
for authors and subscription information:

<http://www.tandfonline.com/loi/tnah10>

On the growth of the stem of *Fontinalis antipyretica*

Professor H. Leitgeb

Published online: 16 Oct 2009.

To cite this article: Professor H. Leitgeb (1868) On the growth of the stem of
Fontinalis antipyretica, *Annals and Magazine of Natural History: Series 4*, 1:5,
392-392, DOI: [10.1080/00222936808695722](https://doi.org/10.1080/00222936808695722)

To link to this article: <http://dx.doi.org/10.1080/00222936808695722>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly

or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Leucodore calcarea.

To the Editors of the Annals and Magazine of Natural History.

GENTLEMEN,—Allow me to draw your attention, and that of your readers, to the figure of *Leucodore calcarea* appended to my paper last month. I have to apologize very greatly for its erroneous and sketchy appearance. The setæ are by no means correctly indicated, whilst the great branchial cirri, which curve over the back, are omitted altogether. This is owing to illness, which prevented me from seeing to the proofs of the plate. The figure given by Dr. Johnston originally, I believe, in this Magazine, and republished in the 'B. M. Catalogue of Worms,' is a very fair representation of his *Leucodore ciliata*, which I must refer to as a correction of the erroneous one in my plate. Since my paper did not deal with the morphological peculiarities of *Leucodore*, the figure was only of secondary importance.

I am, Gentlemen, truly yours,
E. RAY LANKESTER.

On the Growth of the Stem of Fontinalis antipyretica.

By Professor H. LEITGEB.

The apical growth of this moss takes place by repeated divisions of a three-sided apical cell. The divisional walls are parallel to the lateral surfaces of the apical cell. The spiral of division is as often directed to the right as to the left. The segments cut off from the apical cell by the divisional walls are arranged, in accordance with their origin, in three longitudinal series, and at first incline towards each other at an angle of about 70°. Each segment is divided by a longitudinal wall into an outer and an inner part. The inner part of the segment, which subsequently becomes horizontal (the stem-part of the segment), displays in general the same development as the segments in the roots of many vascular Cryptogamia and in the stem of *Equisetum*. It is divided by the sextant-wall into sextants, in the larger of which an inner cell is cut off by a tangential wall. From the stem-part of the segment is formed the widely cellular, axile tissue of the stem.

The outer part of the segment (the leaf-part) partially retains its inclined position. It divides by a horizontal wall into the *acroscopic* and the *basiscopic basilar* portion. The former grows out into the free leaf-surface, a two-edged apical cell being formed in it. From the basiscopic basilar pieces the buds are developed. Hence *each bud and the leaf standing above it belong to the same segment*. One wall of the apical cell of the bud is always turned towards the apex of the parent shoot. The segmental spiral of the bud is always antidromous to the segmental spiral of the parent shoot. The tangential growth of the basiscopic basilar piece always remains much behind that of the acroscopic portion.—*Anzeiger der Akad. der Wiss. in Wien*, February 13, 1868, pp. 43–44.