THE BOSTON MEDICAL AND SURGICAL JOURNAL.

VOL. LII.

THURSDAY, MAY 24, 1855.

No. 16.

POLYPI OF THE UTERUS.

[Communicated for the Boston Medical and Surgical Journal.]

REPORTS of cases have, usually, a practical value. In presenting the following sketch, the principal object is to show the importance of making a correct diagnosis. A very instructive case is reported in the American Journal of Medical Sciences, October, 1854, page 397, among the "Extracts from the Records of the Boston Society for Medical Improvement." It is a cause of regret that an operation so feasible had not stayed the ebbing tide of life. On the other hand, a conservative principle should guard against unnecessarily heroic treatment, even though its result may place upon record the *only* successful case of a formidable operation.

CASE I.—February, 1833. Widow B., of York, Me.; aged 48; not sterile; has been treated for several years past, by different physicians, for simple menorrhagia. She is perfectly blanched and anasarous. The rational symptoms of polypus have existed strongly enough, yet no sensible sign has hitherto been sought. The touch reveals a large polypus wholly within the uterus, the os being dilated to the size of a half dollar. Slight hemorrhage was caused by the examination.

Considering the almost bloodless condition of the system, I deemed it the most safe course to prescribe a generous diet, tonics and absolute rest; and to defer any attempt to remove the tumor until it descended naturally into the vagina. I did not see her again until the next August, six months afterwards, when I found the pelvis completely filled by the polypus, which had descended without pain. The neck of the uterus and of the tumor could only be reached at the anterior part. The peduncle was as large as a man's wrist. I applied a strong silk ligature, after Dr. Clarke's method. The time required for its separation was four weeks, the ligature being tightened once a week. The fetid discharge and the absence of hemorrhage showed that complete strangulation was maintained. The day after the canula came away, I found her with regular expulsive pains, which she had had through the night. The polypus was soon removed with midwifery forceps. Its weight was two pounds; its texture, fibrous. With no unfavourable

vorable symptoms, she soon recovered her strength and color, and resumed her labors as nurse.

CASE II.—This case affords an instance of another species of polypus (the mucous) occurring in the same individual.

July, 1840.—She has had some uneasiness in the pelvis since January last, but no hemorrhage. Her general health is good. A soft, non-pedicated, smooth tumor, nearly an inch in diameter, is growing from the vaginal surface of the neck of the uterus, near the os tincæ. No treatment was adopted. In November following she came with her abdomen enlarged as at the ninth month of pregnancy. Diagnosed it to be flatus, and prescribed pil. rhei comp. ter in die, p. r. n.—The tumor had increased but little, and there had been no hemorrhage. I did not see her again till July, 1842, when she requested the removal of the tumor, as it escaped through the vulva in making much exertion. She gave me the following brief history. The pills relieved her entirely of the abdominal swelling. In December she commenced menstruating apparently, and continued "regular" till November, 1841, when she was "unwell" for five weeks, and has had no colored discharge since. The polypus is about the size and shape of a hen's egg, attached by a comparatively broad base, partially covering the os uteri. It was ligated, with facility, by a loop of silver wire through a single canula, which came away on the fifth day. The polypus, like all of this and the vascular species, decayed entirely, and "like the baseless fabric of a vision," left not a trace behind. Six years afterwards, I learned that she continued in good health.

CASE III.—An old lady, of 60 years, from D——, N. H.; voids her urine with difficulty. Rational signs of mechanical obstruction.

Without making a sensible diagnosis, her physician had directed her to sit over smoking cobs; which, however, did not cure her. It is a case of polypus of the urethra, size of a large pea, partially concealed by the mouth of the canal. Ligated it with a horse-hair, and the patient returned home.

CASE IV.—October, 1838. Widow S., between 35 and 40 years of age, has slight leucorrhœa, and, on evacuating the bowels, often discharges a few drops of blood from the vagina. A soft, vascular polypus, the size of an acorn, is attached, by a small pedicle half an inch in length, to the upper and back part of the vagina. It was torn away with pocket-case forceps, although this was not a very convenient instrument for the purpose. The silver-wire ligature, which I afterwards procured, is the most simple and convenient.

CASE V.—November, 1839. Mrs. G. has a large family of children; age, 50 years; has had hemorrhage for the last five years, yet she, being a *bon vivante*, is not so much affected by the loss of blood as by distressing pains which are more severe at each periodical discharge. She has wearied the patience of her experienced physician, and is waiting impatiently, through much

tribulation, for a crisis at the "turn of life." She consulted me informally, at the house of a neighbor. I told her the probable nature of her complaint, and the mode of relief. She seemed surprised, and said she had never heard of polypi except in the nose. In her next paroxysm I was summoned to attend her. Found her weltering in blood, and groaning with pains which she said were as severe as those of labor. A pyriform polypus, the size of the fist, occupied the vagina, and was attached to the whole circumference of the os uteri, by a base an inch in diameter. After giving an anodyne, I applied a loop of silver wire with a single canula. The ligature caused no unpleasant symptoms, and the polypus came off in a week. Laid open, it looked like bloody meat, and was sarcomatous. The long wished-for "crisis" was experienced, with joy.

CASE VI.—May, 1841. Widow M., mother of several children, having reached the critical period of life, has been treated by me, occasionally, during the last three years, for supposed idiopathic menorrhagia, as no organic disease of the uterus could be detected by vaginal examinations made at sundry times. At this time the os partially admitted the finger, by which a small polypus was detected occupying the cervical canal. Gave her ergot, and at my next visit, a few days afterwards, the polypus was pendant from the os tinæ, and not larger than the end of the little finger from the last joint. The wire loop was readily applied, and the canula came away in four days. The polypus, being of the vascular species, disappeared. She had no hemorrhage nor catamenia afterwards.

CASE VII.—June, 1842. Miss A., aged 46 years, has a son 24 years old; has flowed at irregular intervals for four or five years past. She supposed it to be the "turn of life," and has not consulted any physician for it till at this time. There is nothing remarkable in her appearance. A vascular polypus an inch in diameter is pendant, by a small pedicle, from the os tinæ. Ligated it with the wire loop, which came away in three days. *Polypus non est inventus*. She had no more hemorrhage, nor catamenia.

CASE VIII.—August, 1845. Mrs. T. was confined with her first child six weeks since, and attended by a neighboring physician, nothing unusual occurring. She has flowed a little ever since. An examination disclosed what I supposed to be a small portion of the placenta lodged in the neck of the womb, and which I attempted to remove, but found adherent, causing pain and a flow of blood. Regarding it, therefore, as an originally intra-uterine polypus which had sustained some injury in the labor, I applied the loop of silver wire. The canula and tumor came away separately, in three days. There was no hemorrhage after the ligature was applied. The decaying mass was an inch and a half in diameter, and of finer texture than placenta.

CASE IX.—1846. Mrs. C., aged 51; barren. While attending her for asthma, she remarked that she had thought she had got

to be an old woman, as she had not had her monthly turns for two years, till now, and did not know what to make of it.

A fibrous tumor is growing from the internal and left side of the neck of the uterus, which easily admits the finger. The tumor is of an oval form, three inches in length, and an inch and a half broad at its middle. The upper extremity is imbedded in the wall of the uterus, and the lower projects an inch below the neck. The discharge of blood is not great, and she says it does not trouble her in any way, therefore I waited further development.

January, 1848, two years afterwards. She says she is as well as ever she was, that she has not bled much nor often, and none since June. The vagina has adhered completely around the projecting portion of the tumor, which much resembled and is substituted for the neck of the uterus.

CASE X.—On the 27th of Jan., 1850, I was called to remove a polypus uteri from Mrs. J., of Greenland, N. H. The late Dr. Cheever, of Portsmouth, had declined to operate, as he had never had a case of the kind before. The patient was 35 years old, and had never been pregnant. She was blanched by long-continued hemorrhage. The polypus occupied and filled the vagina, having its origin from the cavity of the uterus. Applied the silver-wire loop with the single canula, carrying the wire beyond the reach of the finger, by means of a probe bent so as to form a hook looking towards the end, and fastened to a piece of whalebone for a handle. There was no pain upon ligating, and she had no more hemorrhage. The wire was drawn on the 30th, and February 3d. The canula came away on the eleventh day after the application. The polypus was removed by the aid of a tenaculum. It was of the fibrous species, and would, probably, weigh a pound. No unfavorable symptom occurred, and she convalesced rapidly.

CASE XI.—In January, 1855, Mr. G., of Maine, requested me to remove a small polypus from his wife, who was 40 years old, and barren. It was a pale, soft, pediculated, cellular growth from the inside of the left labium pudendi, not larger than the common bean, and was more inconvenient than dangerous. It was removed with scissors, with the loss of only a few drops of blood.

CASE XII.—April 12, 1855. Mrs. D., of B., N. H., aged 46, has three children, the youngest 16 years old. Has not been well since the birth of the last, and for six years past has had almost constant hemorrhage, the longest interval being one week. She has had no pain; some weakness of the back, with pelvic weight. About eighteen months since, her family physician made a vaginal examination, and discovered a tumor growing from the neck of the uterus. With this he did not like to interfere without counsel, which was had. As the tumor was irregular in form, it was feared to be cancerous, and the conclusion was to let it alone. The patient flowed on, abandoned all hope of recovery, and discontinued medical attendance as useless.

A female friend of hers who had heard of the Greenland case

(Case X.) advised her to send for me, which advice, after consulting me, was followed in April. The patient is of spare habit, pale and weak; appetite good, bowels constipated. The vaginal discharge varies in color from a florid to a pale hue, and is without offensive odor. The tumor originates from the anterior surface of the cervical canal, by a peduncle an inch and a half long and an inch in diameter. It is of a lobulated form, or, more strictly, tuberos; the main tumor being two inches in diameter, of an oval form, and turned at nearly a right angle with the peduncle, having two spheroid tubers, each about an inch in diameter, growing from it at its junction with the peduncle. A tubercle, the size of a small pea, is detected on the peduncle midway between its origin, and the tumor. The whole is covered with that perfectly smooth membrane so characteristic of polypus. "Though the shape of the pediculated fibrous tumor is commonly ovoidal or pyriform, they are sometimes botryoidal, flattened, angular, cylindrical, strangulated, and divided into lobes." (Colombat.)

The os uteri admits the finger beyond the attachment of the peduncle, and the cervix is free from scirrhus. Hemorrhage from the polypus attends the examination, as usual. Applied the silver-wire noose with the single canula, without difficulty. The ligature encircled the peduncle just above the tubercle before noticed, and caused no pain.

On the 17th I found her comfortable, having had no pain. The hemorrhage continued slightly for two days, owing, most probably, to the partial division of the peduncle by the ligature. The discharge now is that muddy, offensive slime which always attends the process of separation by the ligature. This, as in all cases, is ordered to be washed away daily by injecting warm water. The neck of the uterus is patulous, indicating the yielding of the whole morbid growth. The ligature, on being drawn, yielded but little. Further attendance is deemed unnecessary.

In a letter dated May 2d, a member of the family says—"The instrument came off the next day after you was here, and the tumor immediately followed; which was about the size of a hen's egg. Her health is now fast improving. She is much better now than she has been for a number of years, and we feel very grateful that your labors have been so successfully crowned."

EZRA BARTLETT.

16 Henry st., East Boston, May 14, 1855.