PATHOLOGICAL.

Medullary Lesions in a Case of Acute Ascending Paralysis. Ballet and Dutil reported at the Société Médicale des Hopitaux, on the 15th of December, 1895, a case of polyneuritis of ordinary character, in which lesions in the cord were shown by three methods; first, by Nissl's plan, second by eosin and hemotoxylin, and third by picro-carmin. By Nissl's method the cells of the anterior cornua were found to have lost their prolongations, to show sometimes a partial swelling of the protoplasm, and sometimes to have lost their granular character. With the eosin and hemotoxylin it was noted that the nucleus was pushed to one side in some cases, and in others was star-shaped. No satisfactory results were derived from the picro-carmin stainings. M. Ballet thought that it was impossible to decide whether the primary alteration was in the cord or in the nerve, but was of opinion that, in spite of Waller's law, there were many modern observations proving the possibility of lesions of the cord consecutive to neural changes. There is nothing in the cellular modifications which would allow one to say whether the alterations were primary or secondary.

MITCHELL.

Hysterical Meningitis. *Le Med. Mod.*, 18th December, 1895, M. Huchard reports a case of hysterical meningitis. The patient, a syphilitic, had convergent strabismus, and later complete blindness, fever and delirium. At the apex of the right lung there was a slight loss of resonance. The diagnosis of hysteria was considered established by characteristic hysterical attacks, and M. Huchard also thought that it was common to find a similar dullness over portions of the lung in hysterical patients, which he believed to be due to contraction of the muscles. It is often mobile, and provoked by the slightest touch, even of the finger in percussing. It is important to distinguish, as it might create a suspicion of a pulmonary lesion of a tubercular character, and lead to a false diagnosis of tubercular meningitis, when it occurs with such a combination of symptoms as was present in this case.

MITCHELL.

Abnormal Development of the Nervous System as a Stigma Sign of Degeneration. By Prof. H. M. Popow (*Neurologisches Wiestnik*, Vol. III. No. 3.) The report concerns the case of a man, aged forty-two, who at the time of his admittance to the hospital of Karan, complained of pain in the lower extremities, accompanied by disturbances of gait. These symptoms were found to be due to multiple neuromata and fibromata, which