Fourthly, may it not sometimes be well to combine drainage with castration, thus affording immediate relief to the cystitis and pyelitis while the prostate has time to shrink at its leisure?

Fifthly, is the removal of the testes followed by any profound nervous shock, or is any deterioration or loss of balance of the nervous system likely to follow it?

In the above hasty remarks I have taken it for granted that, for the present at least, the operation would be confined to cases so seriously affected that some operation is imperatively demanded. It is conceivable that, if subsequent experience proves castration to be free from remote disadvantages, patients may sometimes elect to have it done early in the disease, before the bladder and kidneys have suffered secondary changes, and when it might be more efficacious in arresting subsequent morbid growth than afterwards in removing it.

Such possibilities are so alluring that we should approach them with cautious and careful scrutiny.

THE OPERATIVE TREATMENT OF PROSTATIC HYPERTROPHY.¹

BY FRANCIS S. WATSON, M.D., BOSTON.

In the paper which we have just heard from Dr. Cabot, that part of the subject which deals with the palliative measures has been discussed; they suffice in the majority of cases to avert or postpone the dangers of the disease, and to make the patient wholly or tolerably comfortable. It remains to be considered in what way we are to deal with that unfortunate minority of sufferers who, in spite of them, go on from bad to worse. For them there are, fortunately, other means which may relieve. These are the operative ones.

They may be divided into two classes: (1) Palliative, (2) Radical. The former consist of the various ways of draining the bladder, and are equally applicable to temporary, or to long continued, or permanent drainage. The two ways of securing bladder drainage are through the perineum by a perineal urethrotomy, and from above the symphysis pubis by a suprapubic cystotomy.

The technique of these operations is so familiar that I will only refer to some of its more important points when speaking of drainage.

When drainage is conducted through the perineum, it is desirable to provide for it in a manner which will thoroughly drain the bladder, and, if the treatment is to be continued for a long time, will allow the patient to move about comfortably. The arrangement which I think best fulfils these indications is that which I devised some three years ago. It consists of a hard-rubber button with a hole in the centre through which a catheter is passed, and four smaller holes, two on the upper and two on the lower margin, to which are attached four stout round rubber cords, which, when fixed to a waistband properly adjusted, hold the button firmly against the perineum.

The central hole in the button is to be one size smaller than the catheter it carries; if the catheter is stretched its calibre is of course diminished, and the button can then slide along it, but becomes fixed at whatever point may be desired by letting the catheter take its natural shape.

Dr. Tilden Browne of New York has contrived a similar arrangement, of which he speaks as follows: "The holder is of light construction except the two steel springs; these have sharp teeth which hold but do not puncture the rubber tube. The springs release the tube by turning the screw collar. The cylinder has a calibre of thirty-five, French, and the spring teeth have play sufficient to hold any tube of from thirty-five to twenty-two, French. Tapes or narrow bandages slipped under the horns of the plate, then brought up radiating and secured to a waistband in front and behind hold the apparatus in place. A con-

FIG. 1. The author's perineal drainage-tube, and hard-rubber button.

tinuous tape for each side is preferable, as it permits the plate to remain stationary while they play through or under the horns, as the changed position of the body in sitting or reclining may necessitate. The plate is narrow enough not to meet the opposing surfaces of the thighs. Nor does it interfere with defecation. I have the patient wear a fresh pad of iodoform gauze between the perineal opening and the plate after each movement of the bowels." (Fig. 2.)

FIG. 2. F. Tilden Browne's perineal tube-holder.

Two catheters should be provided, one for the day and one for the night, the one not in use being kept in a weak solution of carbolic acid.

With regard to suprapubic cystotomy, I think the practice of puncturing above the symphysis with a large trocar and leaving in the canula is a dangerous one, and should be condemned. It is much safer to do a true cystotomy. The incision into the bladder, when made for drainage alone, should be a short one.

Several modifications of the suprapubic operation for drainage have been introduced, the most notable being that of Hunter McGuire, who several years ago devised a method for making an artificial suprapubic

¹ Read before the Massachusetts Medical Society, June 11, 1895, and recommended for publication by the Society.

urethra, which he does as follows: The opening into the bladder is made at the lowest available point, and only the upper end of the abdominal wound is left open, the rest of it being sutured. This leaves a fistulous tract about three inches long, the walls of which are held together by pressure of the abdominal viscera and the action of the recti muscles. It is claimed that in this way the patients are given voluntary control of the bladder. Morris of New York has modified the operation by turning skin flaps down into the channel.

FIG. 3.
Buckstone Browne's
suprapubic drainage-
tube.

A variety of contrivances for carrying out suprapubic drainage have been introduced; one of them will suffice as an illustration of those which are suitable for permanent drainage; it is that of Mr. Buckstone Browne (Fig. 3). While the patient is still confined to bed, the double tubes of Perier, led through the dressing, I have found to be the most useful of those that have been proposed. They are shown in Fig. 4. If properly applied, the patient is kept perfectly dry. Dr. Fowler, of Brooklyn, and Dr. Farrar Cobb, of Boston, have suggested capillary drainage, the latter by means of an iodoform wick led through a glass tube. This method, also, is applicable only while the patient remains in bed.

The relative merits of the high and low operations as performed for drainage alone are as follows: The perineal is safer, is more rapidly done—an important point in these cases in which the danger is greatly increased by the effect of prolonged etherization on the lungs and kidneys. In performing it the urethra can be laid open with one cut of the knife under primary anesthesia, and as soon as this is done the ether can be removed. The disadvantages of the low operation are that some patients cannot support the presence of a catheter in the deep urethra, and that it sometimes causes epididymitis. The high operation avoids these

FIG. 4.

drawbacks, and offers the special advantage of allowing exploration of the bladder; the tube also is generally worn more comfortably above the symphysis than through the perineum; on the other hand, the high operation is rather more serious, takes more time, and if there is a small, contracted bladder, or if the patient has a large belly, the high cut may be very difficult to execute. Occasionally the perineal operation is embarrassed by ankylosis of the hip. From which it will be seen that the choice will be deter-

FIG. 5.
The author's method for tying
in a soft-rubber catheter
into the bladder.

FIG. 6.

mined by the condition of the patient, the degree of irritability of the deep urethra, etc. If, for example, the patient has chronic bronchitis, threatening uremia, and is exhausted, one would naturally drain his bladder in the easiest and quickest manner possible, and that would be through the perineum; or if it is thought best to avoid even the slight shock of that operation, a catheter can at least be tied in through the urethra. A very simple and satisfactory way of doing this occurred to me while operating one day on a case of stricture; it is that shown in Fig. 5, and consists in slipping over the outer end of the soft-rubber catheter that is to be passed into the bladder a piece of drainage-tube about five inches long, which has been cut in halves longitudinally, except for an inch and a half, through which the catheter passes, and by which it is firmly held so that it cannot slip in or out; all that remains to be done is to lay one of these halves along either side of the penis, and to retain them there by two circular strips of sticking plaster surrounding the organ. The catheter should be a little larger than the inner calibre of the drainage-tube, so that the latter will bind it sufficiently to prevent its slipping in or out.

The benefit of permanent drainage is twofold; it relieves cystitis, and what is more important it often averts or lessens the dangers of pyelonephritis, which is so frequent an accompaniment of this disease. I do not think that sufficient attention has been given to this method of treating the pyelonephritis in this

class of cases. Reginald Harrison advocated it several years since, and I have found it give strikingly beneficial results in a number of cases. The surest sign of improvement of the renal condition is an increase in the specific gravity of the urine; with it there is usually a marked gain in the general condition of the patient.

Whatever may be the manner of securing bladder drainage, its essential feature is, I believe, that it be maintained, or at the least that the channel for drainage shall be kept open, so that it may be resumed at any time, without a subsequent operation to re-establish it.

FIG. 7.

FIG. 8. Bottini's instrument.

lish it. This, because the conditions for which drainage was originally undertaken, even if temporarily relieved, are, practically speaking, sure to return in this class of cases, if drainage is wholly suspended and the channel allowed to close.

RADICAL OPERATIONS.

The defect of the operations for drainage alone is that they do not remove the cause of the trouble. It is not surprising, therefore, that surgeons should have tried to do this; and dating from the time of John

Hunter we find records of various methods by which removal of the obstructing portions of the gland was attempted. But the first systematic effort was made by Mercier, in 1857, who divided small median enlargements by means of specially constructed instruments, which he passed through the urethra (Figs. 6 and 7), and reported successful results. Mercier had no followers, however, until Gouley in New York and Bottini in Italy revived his operations about twenty years later; the latter performing them with a galvano-cautery instrument of the same form as that of Mercier's (Fig. 8); the former making the first step in advance by doing them through an external perineal urethrotomy.

In 1884 Harrison reported successful results from tunnelling the prostate with a trocar, and instituting drainage by leaving the canula in the channel so made; then in 1886 we have the introduction of the suprapubic method by Belfield of Chicago, and its establishment a little later through the successful cases and advocacy of McGill and Atkinson of Leeds in England, under the name of suprapubic prostatectomy. This was in 1886 and 1887. I did my first suprapubic prostatectomy in December, 1886, and my first perineal prostatotomy in the spring of 1887, but these cases were not published until 1888, when they were reported at the annual meeting of this Society, in June.

At that time the subject was in a very chaotic state; there was no agreement among the few surgeons who had done the radical operations as to the best methods to employ, and all such operations had recently been condemned by the authoritative statements of Sir Henry Thompson, Guyon and Socin.

The monograph which I published in 1888 had for its objects, to disprove the conclusions of Sir Henry Thompson and the others, to formulate the subject, and to place it upon a rational basis. In short, to show that these radical operations were feasible, in what class of cases they were appropriate, to demonstrate what the conditions were that should determine the choice of method, how they were to be best performed, and finally what prospect of success they offered. The conclusions were based upon anatomical investigations that had extended over a number of years, and upon such clinical data as were then available; the latter consisting of 21 cases of prostatotomy and 26 of prostatectomy. The conclusion reached was that a majority of cases could be successfully operated upon through the perineal route. It must be remembered that the subject was in its infancy then, and that in spite of the improvements in the high operation recently introduced, it was still a much more serious one than at present; my opinions have been since greatly modified in favor of the high operation, and still more recent methods to be mentioned presently.

From this time there was a rapid advance, and the number of cases reported multiplied fast. Opposition arose, however, especially from the French school, and was expressed in an interesting essay by one of Guyon's former pupils, Vignard, who in 1890 tried to prove that the loss of expulsive power of the bladder must be in the great majority of cases a factor fatal to the success of radical operations, even though, as had been demonstrated, the obstructing portions of the gland could be removed with moderate safety. He was met by Belfield and others with a mass of clinical

observations which had within a short time afterward accumulated to the number of 150 cases of radical operations, in which it appeared that two-thirds of the patients who survived had regained most of or the entire power of natural evacuation of their bladders after the operations, and were in good general condition. Since 1892 the operations have been modified and their scope extended, and to-day the operative treatment of prostatic hypertrophy is a well recognized and established procedure.

Such is, in outline, a sketch of the evolution of this part of the treatment of the disease, up to the most recent steps in advance. The latter I will describe, so far as the very short time at my disposal permits, when speaking of the technique of the radical operations, which I will now take up in sequence, as follows:

(1) Perineal prostatotomy. (2) Perineal prostatectomy. (3) Dittel's operation. (4) Tunnelling the prostate. (5) Zuckerlandl's operation for total extirpation of the prostate.

(1) Perineal prostatotomy consists, as you know, in a division of a median enlargement through an external perineal urethrotomy; it is efficient in but a limited number of cases, those in which there is a bar at the neck of the bladder, and being but little less dangerous than (2) a perineal prostatectomy, which is more thorough, it has given place to that operation for the most part; the latter is applicable to such cases as have but little or no lateral enlargement, and in which the median enlargement is a bar simply or a small lobe. The condition of the prostate can ordinarily be determined beforehand by rectal examination, and exploration of the bladder with a short beaked sound, more or less accurately, and the method of operation selected accordingly. If there is a small pedunculated middle lobe it can sometimes be surrounded by a wire snare passed beside the left index finger and held in position by it while the wire is drawn tight by an assistant; if there is a bar at the neck of the bladder, the excision of a V- or U-shaped piece from its central part will usually open a clear road into the bladder; these incisions can be made with a probe pointed bistoury, or with a galvano-cautery, or with instruments of special forms called prostatectatomes, of which there are a variety; the after-treatment consists in drainage and cleansing of the bladder.

Next comes Dittel's operation (proposed by him and executed by Küster). It consists in the excision of a wedge-shaped piece from each lateral lobe, at their anterior ends, after exposing the prostate and separating it from the rectum through an incision beginning at the tip of the coccyx, embracing the sphincter ani in the form of an ellipse and terminating in the middle of the perineum, the object of the excision of these portions of the gland being to cause widening of the prostatic urethra by the subsequent cicatrization of these wounds in a lateral direction, with the idea that the part of the lateral enlargement bordering the urethra would thus be drawn outward to either side. Küster's cases were not sufficiently successful to encourage further trial of this method, nor does it seem to me to be founded on a sound principle, and finally, there are other means at our disposal which promise better results.

Tunnelling the median lobe through a median perineal urethrotomy, as practised by Reginald Harrison, has already been spoken of when discussing

drainage. I do not think that this surgeon now employs it, and there is no difficulty in draining the bladder satisfactorily through the posterior urethra. The distance to be traversed by a trocar in tunnelling is often long, and the opening up of the tissues is by no means free from danger; the operation therefore does not seem to me to be a good one.

The last of the perineal operations is that proposed by Zuckerlandl for total extirpation of the prostate; he suggests that this be done by separating it from the rectum, and then dissecting it from the bladder through a long transverse crescentic incision extending across the perineum from the tuberosity of the ischium on one side to that on the other; this operation has not, so far as I know, been actually performed.

SUPRAPUBIC PROSTATECTOMY.

In the first place let us glance at the history of suprapubic cystotomy apart from its application to the prostatic removals. It is a time-honored one, and has passed through many vicissitudes. The suprapubic operation, properly speaking, had its origin in the hands of Pierre Franco, who performed it at Lausanne in 1560, for the removal of stone; we find already with him the suggestion of one of the modern steps of the operation, that of lifting the bladder upward, and making it more accessible by raising it by pressure from the rectum; this Franco did by two fingers of the left hand pushed well within the rectum. The case of Jean Doot of Amsterdam, a bootmaker, who, assisted by an apprentice, cut a stone out of his own bladder with a cobbler's knife from above the symphysis, and lived to tell the tale for many years afterwards, is one of the most interesting of the early ones. Another of the modern steps was forestalled by more than three centuries by Rousset (1581), who tried to raise the bladder by filling it with water, and about one hundred years later (1685) Piétre accomplished the same end by introducing into the bladder a long beaked sound armed with an arrow-like point, which could be projected from its tip. After the preliminary incisions, the tip of the instrument pushed up the anterior wall of the bladder above the symphysis and pierced it with the dart, which then acted as a guide to the incision into the organ.

The operation was frequently practised by laymen, as were the perineal cuts for stone, especially by priests. These men were known as "cutters for stone," and if we may believe the records, some of them had as wide an experience and almost as great a success as the surgeons of to-day. The operation was not much heard of between 1650 and 1750, at about which time it was revived by Douglas and Cheselden in England, after which it became relatively popular and extended to France and Germany, being practised in the former most successfully by the priest Frère Cosme, who in 1799 had performed the operation 100 times with but 19 deaths, not a bad record when we reckon with the absence of asepsis. In our own century, we find it in the hands of numerous surgeons: Dupuytren, Souberbielle, Sir Everard Home, Leroy d'Etiolles, Nelaton, Civiale, etc., etc. But between 1840 and 1880 the operation steadily lost ground on account of the introduction of lithotripsy by Civiale, and of the much smaller mortality attending this and the perineal operations for stone.

In 1880 it was once more resuscitated by Petersen, who proposed to avoid wounding of the peritoneal fold

covering the anterior surface of the bladder, which injury had been one of the chief dangers of the operation, by lifting the bladder well above the symphysis by means of the combination of filling the organ with water, and distending a rubber bag previously inserted into the rectum. This manœuvre, when practised in connection with the modern aseptic methods, lowered the mortality so much that the suprapubic operation once more took its place among the favorite surgical procedures, and its use was very soon further extended by its application to cases of bladder tumors and prostatic hypertrophy.

In Petersen's method about ten ounces of fluid were injected into both the bladder and the rectal bag. Presently reports of cases in which the bladder had been ruptured or the rectum injured were published, the procedure was thus shown not to be free from danger. This accident, however, has happened very rarely, and if in the adult not more than eight ounces are used for each injection it practically never will happen. Many surgeons have discarded Petersen's method, employing instead the Trendelenburg position, (Fig. 9), which was introduced by that surgeon

FIG. 9. Trendelenburg's position.

about this time, and which has become since so familiar to all. The elevation of the pelvis in this position lets the abdominal viscera fall toward the diaphragm, in doing which they draw the bladder after them out of the pelvis above the symphysis.

Trendelenburg's position is of much service, but the patient should be kept in it for as short a time as possible, especially if he is old and enfeebled. On this account I prefer to make the preliminary incisions before so placing the patient.

As to the incisions themselves we have the choice between the old vertical one and the transverse; the former extends upward from the middle of the upper margin of the symphysis for about three inches, and is carried down to the prevesical space through the linea alba and between the pyramidales muscles. These first cuts can be made rapidly until the prevesical space is exposed. This is easily recognized by the fat tissue which fills it. Immediately beneath the fat lies the bladder and within the fat layer the peritoneal reflection of its anterior surface. The bladder is now exposed in one of two ways. The fat tissue can be raised up by the tip of the finger inserted below the symphysis, or it can be divided cautiously with blunt-pointed scissors. I do not like the former method, because there is necessarily more or less disturbance of the prevesical tissues. When the fat is abundant a great quantity of it has to be drawn up before the

bladder surface is laid bare. A considerable cavity is formed in consequence, which gives lodgment to the urine, and may be the starting point of serious suppuration, even in our aseptic days, while the only objection to dividing the fat with scissors is the chance of wounding the peritoneal fold, which lies in it, but with ordinary care this is easily avoided. The bladder presents itself as soon as the fat is divided. It is readily recognized by its bluish red color and by the large veins upon its surface. I think it best to place the patient in the Trendelenburg posture when the prevesical fat is reached and before it is divided. In doing this as the pelvis is raised, if the tip of the finger is laid in the wound the bladder can often be felt as it emerges from the pelvis. With Trendelenburg's as with Petersen's method the peritoneal attachment is usually carried far enough above the symphysis to readily escape being wounded. If using the former it is well before opening the bladder to protect the peritoneal cavity from the urine by packing sterilized pads around the part of the bladder to be incised. If the patient is not in the Trendelenburg position a tenaculum should be passed through the bladder wall below the peritoneal attachment and held by an assistant in order to keep the bladder raised after it has been opened. The same result may be obtained by passing a suture through the bladder wall on either side of the proposed incision and through the margins of the abdominal wound; this can be done with a short curved needle. The tip of a long beaked catheter which has been previously passed into the bladder and pressed against its anterior wall above the symphysis, though not necessary, serves as a good guide in opening it. The transverse cut gives more room, and the bladder can be exposed more rapidly by it. Its defects are the risk of hernia and more or less impairment of function of the muscles. It is made in the form of a crescent, with its centre opposite and a little above the middle of the symphysis, the deeper cuts being in the same line, and dividing the whole or a part of the recti and pyramidales. On turning up this flap the prevesical space and its contents are freely opened to view. Personally I make a compromise between the two, dividing the skin by the transverse cut, and turning back a good-sized flap. The underlying fascia and muscles are thus fully exposed, and it is very easy to pick out the line of division between them, which I then lay open by the vertical cut, and if more room is needed later in the operation it can readily be gained by snipping with scissors the inner borders of the muscles on either side. It is very rare that more room will be required than is given by this means.

In opening the bladder the vertical or the transverse cut can be used. I see no advantage to be gained by the latter. The vertical cut into the bladder is to be made between the two sutures already described, with one stroke of the knife, avoiding the veins on the surface of the bladder, if convenient; no harm of consequence is done if they are wounded, for the congestion of the vessels is so much diminished when the organ is opened that hemorrhage ceases spontaneously as a rule. Up to this point the steps are alike for intravesical operations of whatever nature. In attacking the prostate there are special measures to be taken.

In the first place, much help may sometimes be gained by inspection as well as palpation of the gland within the bladder; to effect this several contrivances

are used. Hurry Fenwick has introduced an instrument, which he terms a caisson. Figs. 10, 11 and 12 show this device. By it is sought the opportunity to work at the bottom of a dry well, so to speak, on a diminutive scale, as is done on a large one in laying the foundations for the piers of bridges, etc. I have never used it and so cannot speak of it from experience.

Perhaps because of the partiality of the inventor for his own inventions, I like my own instrument best for this purpose; it requires a little practice before it

FIG. 10.

FIG. 11.

FIG. 12. Hurry Fenwick's "Caisson" for inspecting the interior of the bladder.

can be used with advantage, and I think it is rather more suited to cases of bladder tumors than to the prostatic ones, but it may be equally serviceable in many of the latter. It is represented in Fig. 13.

The removal of the obstructing portions is effected according to their form and size; formerly, only the parts projecting into the bladder were removed, but it was found that it was necessary to do more if a clear road for the exit of the urine was to be re-established; a prolongation of the middle lobe into the prostatic urethra is, for instance, pretty constant, and this when present should always be removed as well as the salient part of that enlargement within the bladder. Again it is often requisite to take away more of the lateral lobes than is represented by those parts which

project within the bladder. This was at first done by splitting the mucous membrane and capsule over these parts, and shelling out the obstructing lateral portions, if, as often happens, they consist of distinct enucleatable masses (fibro-adenomata); if not, by removing them with curette, or hitting them off piecemeal with suitable instruments. Later, the same end was accomplished better by the combined operation, which will be described presently. If there is a salient middle lobe, it can be taken away with scissors, or galvano-cautery, or by the snare.

The combined operation offers greater facility in effecting the removal of the lateral parts of the prostate than either the perineal or suprapubic, alone; it consists in doing first the perineal, and then the supra-

FIG. 13. The author's bladder speculum.

pubic, or vice versa, according to the conditions presented by the individual case. The mucous membrane and capsule of the gland are then incised over the portions projecting into the bladder, the finger of the left hand pushes the prostate up from below while the enucleation of one or both lateral lobes, in turn, is accomplished by the index finger or a blunt periosteum elevator; in doing this there is sometimes very free hemorrhage. The removal will have to be accomplished in some cases with a curette, in others it is not difficult to shell out the growth whole.

Nicoll has modified, and as it seems to me very advantageously, this operation, by not opening either the mucous membrane of the urethra, or of the bladder over the prostate. By his method a sound is passed into the deep urethra to act as a guide, an incision is

made into the perineum down to the apex of the prostate, but avoiding the urethra, the capsule of the prostate is then incised sufficiently to admit the tip of the index finger easily. In doing this, some part of the prostatic plexus of veins is apt to be cut and a more or less severe hemorrhage to result; this may be controlled by packing, while the operator proceeds to open the bladder by the suprapubic cut. Through the latter incision, the prostate is pressed firmly downward by the finger from above, and as much of the lateral enlargements as may be required to give a free exit from the bladder are removed through the lower wound in the manner already described. Hemorrhage is not usually severe; when it is, it is best controlled by means of the tampon of Keyes as modified by Cabot (Fig. 14). The tampon consists of a long bit

FIG. 14. Cabot's hemostatic tampon.

of sterilized gauze folded in squares upon itself, as shown in the accompanying figure. It is placed in the bladder and the thread which runs through all its layers (A) is carried out through the perineal wound, and upon being drawn tight and secured outside the perineal wound presses upon the bleeding area within the bladder. The second thread (B) is fastened to the upper layer only of the gauze and is led out through the suprapubic wound. When it is desired to withdraw the tampon the lower thread is cut, and the upper one being pulled the gauze is unwound layer by layer without difficulty. This may be done usually at the end of forty-eight hours.

In cases in which there is great enlargement of the gland it might on rare occasions be desirable to use Helferich's modification of the suprapubic operation; this consists in resecting the middle portion of the upper margin of the symphysis pubis, the intention of the operation being to give more room to carry out the subsequent steps.

There is finally one other method of entering the bladder, which I think is especially interesting. It is that proposed by Rydygier, and performed in three instances in my own knowledge, once by Dr. Frank Harrington and twice by Dr. Abner Post, of Boston.

I refer to making the bladder incision through its peritoneal investment instead of avoiding it. In the three cases referred to above the result was entirely satisfactory, and no harm whatever resulted. The advantages, of course, are that no time is lost in trying to avoid the peritoneum, that more room is afforded, and finally, that when it is desired, there is much better chance of securing first intention of the wound by suture. The bladder should be carefully irrigated beforehand, and a catheter *à demeure* is to be worn for about two weeks afterward.

Two questions remain to be answered. What determines the choice of operation, and what is the chance of success.

To answer them we must first know what has been done, and this is set forth approximately in the first table below, together with the mortality attending each method; and next, what is the percentage of cases in which there is restoration of the function of the bladder. This appears in the second table.

	Cases.	Mortality.
Suprapubic prostatectomy, and the combined operation	109	20%
Perineal prostatotomy	29	10%
Perineal prostatectomy	16	20%
Dittel's operation	3	00%

The percentage of restoration of the bladder function after different operations is as follows:

Suprapubic prostatectomy	nearly 80%
Perineal prostatotomy	" 50%
Perineal prostatectomy	" 50% plus.
Combined operation	50%
Dittel's operation	30%

From the above it will be seen that the suprapubic and the perineal prostatectomies have the highest mortality, but that the former has 30 per cent. more of recoveries of bladder function than any of the other methods.

This is what we should expect, since the suprapubic and the combined operations are more thorough and more serious than the perineal ones. This too must be remembered when reckoning the mortality.

Then again, as Mansell Moullin points out in his last work upon the subject (*Enlargement of the Prostate*), published in 1894, the mortality of the first half of this series of suprapubic operations was 25 per cent., whereas that of the last half was but 15 per cent.; and Mr. Mayo Robson, in 1894, published twelve consecutive cases with but one death.

It must be further remembered that this is the least favorable view of the operative results that will be shown at any time in their history, should the practice be extended in the future, because this series includes all the operations performed in what may be called the experimental stage of the undertaking, in which it must be remembered they have been applied to those cases only which have already passed into the worst forms of the disease.

Could we collect an equal number in which this operation had been performed earlier, in anticipation of the graver stages, we should no doubt find a greatly lessened mortality.

Then too, what of the dangers of not operating! We know very well that they are grave, but how great as compared with the operative risk, we know not at all. No one has ever, so far as I know, formulated any extensive observations as to the number of cases of prostatic hypertrophy which had already reached the graver stages of the malady (in which condition, be it remembered, almost all the patients

FIG. 15. From the author's operative treatment of the hypertrophied prostate, 1888.

FIG. 16. From the author's operative treatment of the hypertrophied prostate.

FIG. 17. From the author's operative treatment of the hypertrophied prostate.

FIG. 18. From the author's operative treatment of the hypertrophied prostate.

have been, upon whom the radical operations have been performed), and shown the results in the absence of operative, and while continuing under palliative treatment. If such compilation were made, it is my own impression that the comparison would prove favorable to the operative measures, in that particular class of cases.

The choice of operation depends, I think, upon the following factors. First, the age and general condition of the patient. Thus, if we have to do with a very feeble or a very old man, whose term of life is in all probability but a short one at best, we should naturally prefer to operate for drainage of the bladder alone, as described in the first part of this paper.

If the patient, on the other hand, is in good general condition, has not yet shown evidence of serious renal trouble, and might, if radically relieved of the obstructing prostate, live for a good many years, then it is right to try the more radical measures; and of these, for the reasons already given, I prefer the combined operation as modified by Nicoll for all cases in which it is necessary to remove considerable portions of the lateral lobes, as in Fig. 15, and the simple suprapubic method, where the median portion alone is involved, and is large (Fig. 16). When there is a bar at the neck of the bladder, or a small median enlargement (Figs. 17 and 18), I think perineal prostatectomy is to be preferred. It will usually be successful, and is a safer operation than the high one. This is the view of the operative treatment of prostatic hypertrophy that I should have given one year ago.

In the last year three new methods have come upon the scene.

The first is that of ligature of the internal iliac arteries for the purpose of producing atrophy of the prostate by limiting the blood supply. Of this I have only to say that it seems to me a thoroughly irrational procedure.

The second, suggested by Dr. Mears, of Philadelphia, is ligature of the spermatic cords, with the hope of producing atrophy of the prostate, by a similar effect to that of castration, but by a lesser operation. Of this method there is not sufficient experience to base a judgment upon at present.

The third seems likely to revolutionize the treatment, and may perhaps replace and make obsolete all the operations which I have been describing to you. It is that of double castration, proposed by Prof. William White, of Philadelphia, in June, 1893. This part of the subject will be presented by Dr. Cabot, and I will therefore not enter upon it.

STATISTICS CONCERNING INSANITY.¹

BY EDWARD B. LANE, M.D.

THOSE of us who are working in hospitals for the insane are asked daily by some one, What is the cause of insanity? or perhaps more often of late, What is the cause of the terrible increase of insanity, as shown by the census?

Strictly speaking, we who see the insane in hospitals cannot answer this question, which is one of the problems of social statics, with the authority that is expected of us. Questioning the patient will not give us the truth as to the cause of his trouble. The friends

¹ Read before the Boston Society for Medical Improvement, April 29, 1895.

are very loth to admit certain causes that would injure their family pride, and they deceive themselves into believing that overwork was the cause or ill health, whereas the family physician, if we perchance meet him, lets a flood of light upon the subject. He tells us perhaps of dissipation or of questionable habits which no one likes to confess of himself or his brother.

But for all this, asylum physicians think more upon this subject than do others, and people have a right to ask of us, What is the cause of insanity? and, What cause or causes are operating to make our hospitals so crowded faster than liberal States and cities can build new and expensive hospitals?

Many, years ago it occurred to some former commissioner of lunacy, perhaps of the Roman Empire, to demonstrate the causes of insanity. Commissions of lunacy ever since have not ventured to depart from the time-honored custom, namely, to request the superintendents of hospitals to forward to such commission or publish in his annual report the causes of insanity. If a superintendent reports 847 admissions, he is to fill out the table of causes; and he is forced by unalterable custom to report 847 causes, no more, no less.

A few men try conscientiously to make this table as near the truth as possible. But the request is an absurd one. Even more absurd than when as physicians we are expected to give the one cause of death in the return to the board of health when the pathologist reports no less than twenty lesions discovered at the autopsy. I confess I have been embarrassed more than once to select from the pathologist's list of from twelve to fifteen lesions the one cause to be put on the death-certificate. Who of us will not be influenced, unconsciously perhaps, to consider appearances and professional pride in selecting the "principal cause," for official record? How much more are friends or family physicians influenced in filing the cause of insanity to go on the commitment papers to be filed in the Probate Court!

It is the exception, I believe, that there should be a single cause for an outbreak of insanity; and I would respectfully suggest that our statisticians at the State-houses revise their method of getting at the causes. This has been attempted in one point the past few years in Massachusetts. I think all asylum men who have given this subject consideration are thoroughly dissatisfied with the cause-table in our reports. I have long felt that the compiling of it was not only a waste of time but that the result was a scientific lie.

As in so many things in our human life there are exciting causes and predisposing causes; and what we want to see tabulated is, In how many of the 847 cases was there evidence of heredity? in how many cases evidence of excess? in how many marital or family trouble? Such tables should be the result of careful study and not compilations from the commitment papers, the authority of which is generally grossly unreliable in the matter of cause — and I know many tables of causes in our hospitals are merely such compilations.

But if, then, our tables of causes are so unreliable, fortunately most of our other tables are not so, and one who will study them can learn much. Knowing that much time has been given to the compilation of these facts, and that in Massachusetts especially they are far above the average in accuracy and fulness, I have