CONSTIPATION AS A FACTOR IN DIARRHŒA.

BY C. P. PUTNAM, M.D.

THE object of this paper is to emphasize the important rôle played in diarrhœa by constipation, or, we might rather say, by retention of fæces. This is not a new subject, yet, I think the importance of it is not enough recognized.

Diarrhœa differs in its tendency from constipation, in that it is, almost always, an attempt at a cure. It is a successful attempt in many cases, and, though more often unsuccessful, is still a move in the right direction.

Diarrhœa occurs in most cases because there is something in the intestines which ought to come out. In case of constipation, however, we can by no means equally say that it is an attempt to keep in the intestines what ought not to come out. It is true that in certain classes of cases, no one hesitates to cure diarrhœa by increasing it. Green apples at once suggest castor oil, yet, I think that neither this drug nor calomel, nor, in their proper place, the many other laxative medicines and diets, are appreciated to their full value.

If I need to make any apology for bringing before the Society an old and simple subject, I think one can be found in the following cases, which are a few of the many that come to my mind:

(a) A girl little over a year old, she lived in a country town; had been nursed by her mother. She was weak at birth, and bled at the navel for a short time. When about three weeks old she had some kind of stomatitis and from about that time on, suffered occasionally with constipation. From time to time, however, she had diarrhœa; and for this, for the most part, she had been under treatment.

The diarrhœa had been worse when I saw her. For a week she had from six to nine dejections daily, which were loose, greenish, with whitish lumps, and very offensive.

In addition to the breast, she had, from time to time, for six months, been fed with potatoes and a little bread and milk, but she had again been put on the breast alone. The patient was extremely restless, especially at night.

The physician despaired of her recovery; and when sent for, a short time before, came unwillingly, saying he thought nothing more could be done for the child, that the digestive organs were unequal to their work.

The mother was advised to feed the child after each breast meal, with milk (warm from the cow, if possible), together with well-baked bread and indian-meal porridge. One-half grain of calomel was to be given every four hours until the bowels were freely moved, or until the offensive odor was gone.

Two days later the mother came, saying the child had taken about a pint of milk daily. The dejections were less green, but still very offensive. The child was even now very restless at night. Ipecac and calomel (one-quarter of a grain of each), were ordered to be given every four hours until the offensive odor disappeared, even if the bowels were very freely moved in the process.

One month later the mother came again to say the child had gained greatly in weight, and seemed perfectly well, the discharges having become natural and not offensive. Its health has never been so good.

(b) Miss B., about thirty years old, had suffered from diarrhœa for several years, seldom being free

from it except for short periods. The discharges were dark-brown, and offensive. She had received treatment for this trouble at various hands, but without any lasting success.

For a short time I advised hot-water enemata. This gave a certain degree of alleviation. Then a strong cathartic pill was administered, and from that time she improved rapidly, and the diarrhœa practically disappeared.

(c) A boy of fifteen months, seen in consultation. He had been attacked suddenly, a week before, with diarrhœa of unknown origin, and had been treated with bismuth and opium, with the result that he almost immediately improved, and seemed quite well. But soon after the treatment had been suspended, the bowels had begun to move, and the discharges became loose, green and offensive.

The same treatment again relieved the symptoms, only to have them reappear when the treatment was again given up.

Calomel, to be used as in the first case, was recommended; and from that time the child improved steadily.

The last case is that of a gentleman, about thirty-five, who was attacked with diarrhœa while travelling in Europe, and was treated in various cities for it. He took a number of sedatives, including a considerable quantity of opium, and finally came home almost in desperation. He was very weak and moderately emaciated when he arrived here, and was able to eat almost nothing.

He was ordered to take large hot-water enemata, which were increased by degrees to about two and a half quarts, with the result that before long his diarrhœa ceased and his usual health and flesh returned.

In reporting these cases, I have dealt with symptoms, not diseases, in order to keep the subject as simple as possible.

The practice of giving opiates for diarrhœa, irrespective of its cause, seems to me objectionable, and, in many cases, decidedly dangerous.

It is my practice to tell patients that the number of discharges is a very unimportant point, that the main question should always be, as to their quality. Any decomposing material in the intestines should be taken out of them as rapidly as possible. It can only be a matter of regret that we cannot more readily remove such material. Even in typhoid fever, I am inclined to think that the general fear of giving enemata and laxatives is an exaggerated one.

Clinical Department.

FRACTURE OF THE ANTERIOR SUPERIOR SPINOUS PROCESS OF THE ILIUM FROM MUSCULAR ACTION.

BY N. NICKERSON, M.D., MERIDEN, CONN.

As a contribution to the literature of this rare form of accident, I will report the following:

John Coe, aged seventeen, while running a foot-race of one hundred yards, was suddenly brought to a stop by feeling something give way in his hip, with a sensation of bones grating together. He found that he could not take another step, and was carried home where I saw him within an hour after the accident. On examination I found distinct crepitus, with motion

¹ Read before the Boston Society for Medical Observation January 6, 1890.

of fragment, on pressure upon the anterior superior spinous process which elicited considerable evidence of pain.

The leg was placed in a flexed position, resting on a fracture apparatus, but without bandage or other appliance to keep the parts in apposition, and the point of injury kept bathed in an evaporating lotion for several days.

At the end of two weeks he began to walk cautiously by the aid of crutches, and in three weeks dispensed with them altogether, though using care in going up and down stairs, or over rough ground.

The diagnosis in this case was confirmed by Drs. E. T. Bradstreet and E. W. Smith, who saw the case with me.

The following extracts may be of value to those physicians who have not access to large medical libraries, and contain all I have been able to find bearing on this point.

In the *New York Medical Journal*, Volume XII, page 184, and the *Canada Medical Journal*, Volume VII, page 97, Drs. S. Joy and J. W. McWhinnie report this case: Augustus S., age seventeen, a medical student, was engaged in a foot-race where a certain distance had to be run, then to turn and run back. In the exertion of turning he felt something snap in his right hip, walked a few steps and fell. On examination, distinct motion and crepitus could be felt, by the pressure over the process, also by placing the thumb over the origin of the sartorius and rotating the thigh.

The patient was placed in bed, with the thigh flexed, and the shoulder raised, a bandage being applied to aid in steadying the fracture. This position and adduction of the leg were maintained by bands attached to the posts of the bed. In two weeks the patient made a good recovery without displacement. In commenting on this case, the doctors make the following observation: "In looking up the literature of fractures of the anterior superior spinous process of the ilium, we read of their occurrence from direct violence, but not one case is recorded as having taken place from muscular action."

In the *British Medical Journal*, 1872, page 549, this case is reported by Dr. George J. Seeley: A young gentleman, age seventeen, of good muscular development, whilst running vigorously in a foot-ball match, was suddenly brought to a standstill by a snap in his side, followed by a feeling of "coming in two." Being quite unable to take another step he was carried to the house, and, on my examining him about an hour afterwards, I found considerable tenderness along the upper head of the rectus femoris, and a piece of bone, as large as the top of a finger, detached from the ilium; crepitation was very distinct.

In the same journal for 1872, page 513, Dr. W. E. Hyde reports another case, as follows: A young man eighteen years of age, whilst running over some uneven ground suddenly felt as if a stone had been thrown at him, striking him on the hip, and then felt that he could not move again. He was carried home, and upon my arrival an hour afterwards I found that he was unable to draw his leg forwards, and upon examination I found one head of the rectus femoris torn from its attachment to the ilium with a piece of the bone, about the size of a half damson, perfectly movable, so much so that distinct crepitus could be felt by moving the surfaces of the fractured parts together. I flexed the thigh and applied a pad with a figure-of-

eight bandage round the pelvis and thigh. At the end of two weeks union had taken place, with formation of callus, and in another week he could walk about with some difficulty.

Dr. Hamilton, in his classical "Treatise on Fractures," etc., gives this as the only case coming under his observation. William Alexander, aged seventy, on the 5th of September, 1869, after riding in a railroad car about half an hour, arose to leave his seat when he felt "something wrong" in his right groin, and found himself unable to walk without great pain. He was admitted to Bellevue Hospital on the same day, and I found a fracture involving about three inches of the ilium, including the anterior superior spinous process. It was inclined to fall outwards, but was easily replaced, with a distinct crepitus.

The following remarks are taken from the "Treatise on Fractures," by Lewis A. Stimson, M.D., as the last word on the subject. "Dr. Hamilton has reported a case of fracture of the anterior superior spinous process of the ilium by muscular action, and the same agency appears not improbable in other fractures of the same part. Riedinger asserts the same cause in many fractures of the crest. Considering the strength of the muscles attached to the ilium, and the occasional correspondence of the fragments to the insertion of the muscles, the theory does not seem unreasonable."

Reports of Societies.

BOSTON SOCIETY FOR MEDICAL OBSERVATION.

T. F. SHERMAN, M.D., SECRETARY.

REGULAR meeting, Monday evening, January 6th, Dr. C. F. FOLSON in the chair.

Dr. C. P. PUTNAM read a paper on

CONSTIPATION AS A FACTOR IN DIARRHŒA.¹

Dr. MINOT said that most of his cases where diarrhœa had been caused by retention of fœces were those where opium had been given for relief of pain, as in cases of cancer of the rectum. In young children he thought that many cases of diarrhœa were best treated by a cathartic, — and that the domestic practice of giving a dose of oil often gave immediate relief. Retention of fœces was not uncommon in old people, especially in women, and he was often surprised at the large amount of fecal matter which could be retained in the colon and which often caused a catarrh or even ulceration of the intestines if not relieved.

Dr. PUTNAM said that the object of his paper was to bring out the fact that many cases of diarrhœa could only be satisfactorily treated by the use of laxatives and not by giving astringents or opium which was too apt to be the routine practice.

Dr. G. K. SABINE reported

A CASE OF URÆMIC CONVULSIONS DURING PREGNANCY.²

Dr. MINOT said he saw no objection to bleeding except in pale, thin, anæmic women. All remedies fail in some cases and are successful in others. He thought that he had obtained the best results with pilocarpine given in one-quarter to one-half grain doses. Pilocarpine had, however, a depressing effect on the

¹ See page 224 of the Journal.

² See page 220 of the Journal.