The university authorities have determined to depend no longer on the supply of raw water from Fall Creek, and are installing a small mechanical filter which will be completed in a few weeks, and which is expected to ward off the possibilities of danger from this source. During the epidemic the university has supplied artesian well water for the drinking tanks in the campus buildings, and to its prompt action in this respect a considerable degree of immunity among the students during the later stages of the epidemic must be attributed.

Among the many persons in Ithaca to whom I am under obligations for friendly courtesies, I wish to express my especial indebtedness to Prof. V. A. Moore of Cornell University, who has generously placed at my disposal much information concerning the epidemic, and also to Dr. George A. Soper, who has kindly aided me in the course of my inquiries in many ways.

POSTSCRIPT.

The epidemic is now, March 30, to all appearances, at an end. Only fourteen new cases have been reported since March 11, and during the last week there have been only four. The revised and corrected tabulation of cases on the best information obtainable is as follows:

	Students.	Citizens.	Total.
Cases in Ithaca.....	124	660	784
Cases outside of Ithaca.....	106	22	128
Total	230	682	912

The corrected list of deaths to the present date is as follows:

January 23 to January 29.....	1
January 30 to February 5.....	4
February 6 to February 12.....	7
February 12 to February 19.....	12
February 20 to February 26.....	11
February 26 to March 5.....	10
March 6 to March 12.....	5
March 13 to March 19.....	5
March 20 to March 26.....	2
March 28.....	1
Total	58

Twenty-six of this number were students in Cornell University.

New Instruments.

A NEW NASAL SPECULUM.

O. B. MONOSMITH, M.D.
LORAIN, OHIO.

Realizing that the bivalve speculum, opening transversely, did not meet all of the requirements of a nasal speculum, I was led to devise one for my own use that would overcome these difficulties. That I have succeeded, at least to my own satisfaction, is the excuse for presenting it to the profession. Its evolution extended over a period of about three years, during which time I made many modifications. The model, a cut of which I present, I have used about a year, and, in the majority of nares, it meets all of the requirements of an ideal speculum, both diagnostic and operative. It has merits that will appeal to the specialist as well as the general practitioner.

The instrument is to be held in the left hand of the operator for both nares, with the ball of the thumb on the screw head, thus giving a firm hold of the instrument without necessarily forcing the blades apart. By holding the instrument in this

manner the blades open vertically. The advantage of this is that it raises the tip of the nose, giving a more advantageous angle of illumination and a better view of the nasal cavities. Another advantage is that the blades do not touch the mucous membrane. It will be seen that the upper blade is much shorter, and is placed somewhat back of the lower blade. It is also hooked slightly upward. This is, perhaps, the most important feature of the instrument. Being shorter and placed back of the lower blade, it conforms perfectly to the

anatomy of the nose, allowing both blades to enter the nose without leaving anything outside that is superfluous. The blades being attached to the shanks at right angles, prevent their being forced inward against the mucous membrane, and the upward curve of the upper blade prevents the speculum

slipping out at inopportune times, especially while operating. It may be noted, in passing, that with any bivalve speculum it is occasionally necessary to clip out the hair that is sometimes found in the anterior nares, in order to get a perfect view of the nasal cavities. A patent has been secured on the instrument, with the sole purpose of giving the control of the

manufacture to one instrument house that can be depended on to make it right, otherwise the merits of the instrument would be spoiled. As to the wisdom of this, I refer to an article in *THE JOURNAL of the American Medical Association*, vol. xxxvii, p. 1336, by Dr. Frank Allport, in which he writes: "As it is now, the moment he publishes an instrument all manufacturers have a right to make it, and, before long, the instrument on the market bearing the inventor's name may be so far removed from his original idea as to make him ashamed that he ever gave it to the profession."

THE THIERY CLAMP.

A SUBSTITUTE FOR EXTERNAL SUTURES.

ANDREW B. GLONINGER, M.D.

Surgeon to Good Samaritan Hospital.
LEBANON, PA.

I would like to give your readers the benefit of an idea that I picked up in Paris last summer at Professor Thiery's clinic. It is a little clamp, invented by Thiery to take the

Fig. 1.—1. Forceps for applying clamps. 2. Forceps for removing clamps. 3. Clamp before application. 4. Position of clamp when in use.

place of superficial sutures, which saves time in closing incisions after operations and prevents troublesome stitch abscesses.

The clamp itself is a narrow strip of soft non-corrosive metal $1\frac{1}{4}$ cm. in length, the ends curled and armed with little spurs.

Fig. 2.—1. Showing the clamps in place after an abdominal section. 2. Showing the clamps holding a flap in place, reinforced by silkworm-gut suture at points where tension is greatest.

The two skin surfaces to be united are held in apposition between thumb and forefinger and the clamp, held in the grasp of a heavy claw-tipped dressing forceps, is pressed lightly against the adapted edges and pinched on with slight pressure. The little instrument bends easily, and the points enter the skin, holding the edges firmly together. They should be removed on the fourth day. For this an ordinary small hemostatic forceps will do, but the forceps specially designed for the purpose has a prominence on one jaw and a depression in the other; the grooved jaw is slipped under the clamp, and when the other is closed on it it bends back readily into its original

shape with very little annoyance to the patient, leaving only a small cicatrix along the line of incision.

The outfit, including the two pairs of forceps and 500 clamps, cost me 20 francs in Paris. They can be used in all sorts of incised wounds where the tension is not great.

In abdominal operations they are the ideal suture. In fifteen sections I had complete union in four days in all but one, in which I had used some deep sutures on account of an unusually thick layer of fat. Following an amputation of the breast, I use two or three silkworm-gut sutures where there is tension and finish with clamps. I have had good results in closing incisions after phlebotomy for transfusion or for the ligation of varicose veins, and I tried them in two cases of amputation of the leg, but the grasp did not seem to be strong enough to hold the flap, and the result was not very satisfactory. They are specially well adapted for incisions of face, scalp and neck on account of the small cicatrix.

The misconception of the Code of Ethics which enjoins physicians against the methods of advertising like the quacks, i. e., by offering to give public advice to the poor gratis and by private cards or public handbills proclaiming one's skill to cure special diseases, has deterred capable physicians from imparting the kind of information they ought to impart on all proper occasions to the public. Up to date the public has obtained most of its knowledge indirectly from published accounts of epidemics, etc., and what health boards are doing and from occasional newspaper extracts from medical society proceedings selected on the judgment of non-professional and of inexperienced newspaper reporters and from the alarming, overdrawn and often perverted statements made in quack advertisements. Often the first and only knowledge of what the medical profession is doing gets to the public through the last-named mercenary channels, who often pay tribute to the great work of the medical profession by indirect reference to the merits of its dead masters and by naming their quack establishments and more or less fake remedies after our great names. For we know, though the public does not, how many of the patent medicine men of our time have been simply drug store clerks or bottle washers or porters sometimes in physicians' offices, evolved and transformed into patent medicine proprietors and kings of gullibility and Napoleons of fake finance founded on humbug—folly in matters medical and fake healing with the people, because we who have the light to shed give it not, but leave the popular medical world in darkness.—"Alienist and Neurologist."

The proportion of accidents and disease is brought out by an analysis of a recent census report made by the "Insurance Press," from which we learn that deaths due to accidents and injuries were highest among those whose mothers were born in Italy (119.5 per 100,000 of white population); lowest among those whose mothers were born in the United States (62.7). The death rate due to accidents and injuries is highest among persons 45 years of age and over. The average age at death from accidents is about 33.5. The proportions of death from accidents and injuries were greatest in the Cordilleran regions, the Pacific coast region and the Western plains; they are least in the North Atlantic coast region, the Middle Atlantic coast region and the Northeastern hills and plateaus. The warmer months show a larger proportion of fatal accidents than the cooler months. A person is more liable to meet death by accidental injuries than by any other single cause, except tuberculosis, pneumonia or heart disease. One is twice as liable to die from accident as from old age. Typhoid fever, cancer, apoplexy, inflammation of the brain and meningitis, paralysis, diphtheria, diseases of the stomach, liver and brain, all these are slight menaces to life compared with the accident hazard. If, as seems not unreasonable, a person is liable to die of any disabling accident that happens to him, what is to be said concerning the 11,000,000 accidents that happen every year. It seems to be true that a person is 100 times more liable to "catch" an accident than a fatal case of tuberculosis or pneumonia, and about 1,000 times more liable to "catch" an accident than a fatal case of liver disease.—"Amer. Medicine."