point. Von Graefe¹² says "if we imagine the substance of the optic nerve loosened (*geloockert*) through some internal process the papilla will give way to the normal pressure in an anomalous way." Later it was Schweigger who in 1891 (cf. Horstmann) insisted that cupping of the disc to the margin may exist without the least demonstrable increase of tension, and that "such a condition might arise from any atrophic disease of the optic nerve when there was previously a physiologic cup." Neither Schweigger nor Horstmann¹³ refer to the normal intraocular pressure as the direct cause for such "glaucomatous" cupping in apparently non-glaucomatous eyes.

Schnabel's¹⁴ views regarding true glaucomatous cupping as due to disintegration (*Zerfall*) of the optic fibers with formation of caverns, and "the glaucomatous excavation is one of these caverns"—are beside the point raised and are merely quoted for completeness' sake,¹⁵ so are also Mueller's¹¹ conclusions that excavation—in general—might sometimes also be brought about by traction from without through shrinking exudate, etc.

Clearly the point is not of merely anatomic interest; if we could discriminate in glaucoma simplex without increase of tension between cases in which the "glaucomatous" cupping is due solely to simple atrophy, i. e., of fibrillae and finer septa, and those in which it is the result directly of normal pressure on diseased or congenitally less resistant tissue, it would seem that the vexed question concerning the therapeutic effect of operative measures in glaucoma simplex would become clearer.

DISCUSSION.

DR. E. V. L. BROWN, Chicago, said that the pathogenesis of glaucomatous cupping in optic atrophy was much discussed at last year's meeting of the Vienna Ophthalmologic Society. The position was taken that this does occur in a great many cases without any tension whatever, and is due to the atrophy in the nerve head. Just back of it are found extensive areas of atrophic nerve fibers. Schmidt-Rimpler has also reported some observations on this point.

THE PHYSIOLOGIC ACTION OF DIONIN.*

WALTER HAMILTON SNYDER, M.D.

TOLEDO, OHIO.

Dionin is ethyl-morphin hydrochlorate, and was discovered by Grimaux ($C_{19}H_{23}NO_3 \cdot HCl + H_2O$). It is a white, odorless, bitter powder, soluble in water, 7 parts; in alcohol, 1.5; in syrup, 20. It is insoluble in ether and chloroform. The aqueous solution is neutral in reaction and keeps a long time. The general physiologic action is similar to that of morphin, but it is claimed that it is less toxic and does not produce nausea, constipation or habit. The respiratory and cardiac centers are unaffected. Applied to the eye, either as a solution or powder, it produces marked sneezing, vasodilation, edema, glandular excitation, burning and lachrymation and supposedly lymphatic extravasation, followed by analgesia, lasting from 2 to 48 hours. Wolffberg¹ of Breslau first used it in ophthalmic prac-

tice, but it was Darier's² report that brought it prominently before the profession.

The general literature on the drug is enormous, but I have been unable to find any experiments bearing on its actions on tissue and cells. I here cite my own:

Albino rabbit (medium size). Dionin in powder was placed in each eye in larger quantities than would be necessary if the eye was abnormal. The reaction was moderate, giving the usual appearance, only less in degree. The rabbit was killed. The globe and tissue were enucleated and placed in formalin 4 per cent. for 48 hours. Later sections from the cornea, through the ciliary region to the nerve were cut and then put through the usual technic of imbedding, sectioning, staining, etc. Control specimens were made from a normal rabbit's eye. I will here cite the findings:

Slide 1.—Magnification 210 diameters, section through cornea and ciliary region. (Fig. 1.) The lamellar spaces in substantia propria are dilated and edematous. The serous transudate involves the spaces at the junction of the cornea and ciliary region most. Is this not due to the dionin being placed directly over this region?

Slide 2.—Magnification 35 diameters. (Fig. 2.) This is same section under lower power and on close examination it is seen that the effect is greatest on the external side of the cornea and that the muscle is not affected, not even showing edema. Toward the sclera there is slight evidence of hyperemia.

Slide 3.—Magnification 210 diameters. (Fig. 3.) This is a control slide showing normal substantia propria. Compare this slide with No. 1 and see the effect dionin has had in elongating and widening the lymph spaces. In this section the lymph spaces are round, not widened or elongated as in the previous slide.

Slide 4.—Magnification 35 diameters. (Fig. 4.) This is the conjunctiva of a rabbit on whose eye dionin has been applied, suggesting the contention that the tissue changes are found only where there is tension; no edema or round cell infiltration can be found in this slide.

It might be well to introduce here an hypothesis to explain the clinical and microscopic action of dionin and to explain the clinical reports which, on comparison, at times are contradictory. Many slides were examined to arrive at certain conclusions and only the marked features photographed. From these I deduce that the action of dionin is purely local—greatest where the drug has actually rested. Its most marked action is in eyeballs in which the tension is increased, and I believe its entire action can be explained by saying that it has some disassociating action on the intercellular cement substance allowing a transudation of serum from a globe under pressure. That its analgesic action is to be explained by its lessening of tension and thus relieving pain, and by the well-known action of the derivatives of opium. It has been claimed for this drug that it is a lymphagogue, but opium lessens secretions, and the increased flow of apparent lymph, is not lymph, but serum which does not contain cells of any description. Lymph contains colorless blood cells and if the action of dionin was that of a lymphagogue, examination of the tissues would reveal the characteristic components of this fluid. I believe, therefore, that it is only a lymph stimulant secondarily; after the edema the fluid is absorbed as lymph, as it would be in edema from any cause. Careful examination of the slides reveals that locally where the drug is applied there is some change that permits fluid under pressure in the globe to pass through and into the tissue, causing edema and consequent lowering of tension. From these laboratory facts we should expect its action to be most marked in pathologic eyes, and, in these, greatest when the tension is

*Read in the Section on Ophthalmology of the American Medical Association, at the Fifty-sixth Annual Session, July, 1905.

12. Arch. f. Ophth., III, 2, p. 547.

13. "Glaucoma Simplex," Arch. of Ophth., 33, No. 5.

14. Arch. f. Augenheilkd., 1892, vol. xxiv; Wiener klin. Wochenschr., 1901, S. 469.

15. For a critical review of Schnabel's views see "Ginsberg, Grundriss der pathologischen Histologie" des Auges, Berlin, 1903.

1. Wochft. f. Ther. u. Hyg. des. Auges. vol. III, No. 16.

2. Ophth. Klin., No. VII, 1900.

highest. That its reaction subsequently is lessened rapidly, unless the tension is renewed, is a fact familiar to all who have used dionin.

I wish to enter into detail in a few findings not so generally admitted.

Darier³ and Reber⁴ have each reported cases of complete absorption of the lens and capsule during the use of dionin, and in a case of severe contusion of the eye

tension plus 2. Dionin and atropin were used daily. Each application would lessen the tension and it was used daily as a powder until December 28, when the fundus could be seen.

Condition.—Fluid was vitreous, the iris was dilated and maximum and drawn up in a mass in the inner upper quadrant.

Jan. 29, 1905; this iris had disappeared. Dr. Samuel D Risley has suggested as a possible explanation of the disappearance of the iris in this and similar cases that it inverts or

Fig. 1.—Magnified 210 diameters. Cornea and ciliary region showing the lamellar spaces in the substantia propria are dilated and edematous, especially at the corneal junction.

Fig. 3.—Magnified 210 diameters.—Control slides showing normal substantia propria. Compare with Fig. 1.

Fig. 2.—Magnified 35 diameters. Same section as in Fig. 1, through lower power. Note that effect on cornea is greatest on the external side. Muscle is not affected. Sclera is slightly hyperemic, showing how purely local is the action.

Fig. 4.—Magnified 35 diameters. Conjunctiva in fornix after dionin. No round cell infiltration or evidence of edema.

without penetration this has occurred in my own practice.

History.—Sept. 20, 1904, I first saw this patient and found hemorrhage in the anterior chamber, partially opaque lens—

3. Darier, A.: *Clin. ophth.*, 1899, No. 23; 1900, No. 6. *La Clin. Ophthal.*, Jan. 25, 1904.

4. Reber: *Ther. Gaz.*, Feb., 1904.

"umbrellas" over the ciliary processes. Tension was normal, and vitreous fluid.

Later History.—April 2, 1905, a careful examination showed no iris, lens or capsule. With plus 8.00, combined with plus 4.00 cyl. axis 180, vision is 20/40 and a spherical plus 4 added gives Jaeger No. 1 at 17". The reaction to dionin was marked every day, something I have never seen before.

In iritis with adhesions and plus tension the use of dionin lessens the tension and permits absorption of the

mydriatic, and prompt relief of pain and dilation of the pupil results. In corneal ulcers, especially the peripheral type, the repair process begins as soon as the ulcer is cleared. I have observed that the more recent the inflammation and the higher the tension the better the results obtained from its use.

Königstein agrees with this, as he states: "Dionin is almost completely abortive of parenchymatous keratitis if used in the earliest stages of the disease, but valueless in the late." Curiously enough, Graefe⁵ gets "good results in vitreous opacities only when they are of long standing." Bourdeaux⁶ and Batalow⁷ think it is useful in the later stages of corneal opacity and that they clear up more quickly than without its use. In old cases I have not been so successful, but in recent cases my experience coincides with theirs. Central opacities of the cornea, especially if of long standing and with tension minus or normal are but slightly, if at all benefited in my experience. Von Arlt⁸ has treated these patients "as long as 21 months with good results, especially in children." I have not persevered so long. Wolffberg⁹ uses this drug in hemorrhagic glaucoma, which is seemingly logical. Bloch¹⁰ states the greatest reaction occurs in pathologic condition. I should qualify this by adding "with increased tension." Sanz Blanco¹¹ "injects a 5 per cent. solution in intraocular hemorrhages with surprisingly good effects." "In one case the fundus could be seen in 5 days and in 30 days he (the patient) could read."

I have tried it in conjunctival hemorrhage with no reaction to speak of and no beneficial results over the ordinary treatment, evidently lacking the pressure factor. In beginning pannus I have cleared up the cornea and resisted permanent opacity more satisfactorily than with any previous treatment, the lid, of course, being treated for the cause.

In glaucoma, I prefer dionin to eserine, relief from pain being very marked, due, I think, to the mechanical relief from pressure. In old vitreous opacities I have had poor success, possibly because I have not used it long enough. I early abandoned the use of solutions and placed specks of the powder on two or three places on the cornea, especially finding the results better than solutions or applications in one place. This was when I thought the drug a lymph stimulant and is further proof that its action is as suggested in my hypothesis, because the action being purely local the larger number of places acted on the greater the relief from the tension. I wish here to express to W. J. Stone, B.S., M.D., my appreciation of his kindness in assisting me in the preparation of this paper.

DISCUSSION.

DR. E. V. L. BROWN, Chicago, referred to an article recently published on some work done in Breslau in which the author found that dionin had effect only on certain animals; not on all. The animals used were dogs, rabbits and cats, and the cats did not react in any way.

DR. W. H. SNYDER said that in rabbits the effect is very slight. It requires 100 times as much to get a reaction as in the human eye.

5. Graefe, A.: *Deut. med. Wochft.*, xxvi, No. 12.

6. Bourdeaux, B.: *Klin. Ther. Wochft.*, ix, No. 29.

7. Batalow, A. E.: *Vratch*, 1901, No. 19.

8. Von Arlt, F. R.: *Heilkunde*, vi, No. 11. *Wochft. für Ther. und Hyg. des Auges*, No. 11, 1902.

9. Wolffberg, *Wochft. f. Ther. u. Hyg. d. Auges*, iii, No. 4, Jan. 8, 1903.

10. Bloch, R.: *Therap. Monats.*, viii, p. 418; *Heilkunde*, vi, No. 11; *Deutsche Praxis*, 1903, No. 25.

11. Sanz Blanco: *Arch. de Ophthal.*, August, 1904.

Special Articles

MEDICAL AFFAIRS IN THE HEART OF THE ARCTICS.

N. SENN, M.D.

Professor of Surgery, The University of Chicago; Professor of Surgery and Head of the Surgical Department Rush Medical College.

CHICAGO.

The geographic distribution of disease, the study of primitive races, of their climatic conditions, customs, habits and manner of living as etiologic influences, their methods employed in combatting disease and their resources in the treatment of accidents are subjects which can not fail to interest physicians who are concerned in the progress of their profession and the welfare of humanity the world over.

My vacation trip this summer brought me in touch

Fig. 1.—Baffin Land Eskimo.

with, and afforded me an excellent opportunity to study, one of the most interesting peoples in the world, the Eskimos of the far north, in the very heart of the Arctic region. These people are scattered in small settlements along the northern part of the west coast of Greenland, extending from Upernavik to Etah, a distance of about 600 miles. Etah is the most northern habitation of the Eskimos. It is in latitude 78° 40' N. and is less than 700 miles from the North Pole. It is in the heart of the Arctic region, where so many fearless explorers have spent their long, dark winter night preliminary to the dash for the North Pole on foot and sledges over ice and hummocks of ice. It is here where so many hopes were shattered, where so much suffering was endured and where not a few brave sailors have found a premature grave.