to stand it, and the earlier it is done the better; but he would never hesitate to operate on an eye even if the fields were contracted to fixation, provided that there was any vision worth saving; even in the premonitory stage he advocated operation.

Gonorrhœa and Diseases of the Eye.—LAWFORD, London, in a discussion on the relation of the gonorrhœa to diseases of the eye (*British Medical Journal*, November 2, 1901), after excluding purulent ophthalmia, considers the following ocular affection as known to occur in association with gonorrhœa; a form of mild conjunctival inflammation (not due to direct inoculation), scleritis and episcleritis, iritis and iridocyclitis, neuroretinitis. Suppurative keratitis has been described in severe pyæmic case; of these iritis is the best known.

The Diagnosis of Malignant Intra-ocular Tumors in Infancy.—SOURDILLE, Nantes (*Gaz. Med. de Nantes*, November 16, 1901), calls attention to the great difficulties frequently existing in making the diagnosis of retinal glioma. Transillumination should be practised in every case before surgical intervention. If a tumor is shown, and any doubt as to its nature be present, it becomes necessary to make a puncture and to examine histologically and experimentally (by inoculation) the elements of the neoplasm.

Malignant tumors of the retina are all gliomata, properly so called. Mistakes of diagnosis are frequent. These mistakes are masked under the specious name of "pseudoglioma," when enucleation shows that an entirely different disease is present; and "cryptoglioma" when the disease is overlooked. For a long time the amaurotic cat's eye was believed to be adequate for the diagnosis of glioma. This has been the source of numerous errors. The writer concludes from his own histological examinations, that of ten eyes enucleated for glioma three are found to be cases of pseudoglioma.

The amaurotic cat's eye is by no means pathognomonic of glioma; this is found in suppurating retinal hyalitis, purulent choroiditis, affections following infectious maladies of infancy, particularly the eruptive fevers, and, above all, tuberculosis of the intra-ocular membranes, the choroid and retina.

It has therefore been proposed to narrow the diagnosis of these tumors and to lay down an absolute rule following Lagrange; in glioma the eye is glaucomatous or tends to glaucoma; in pseudoglioma the tension is below normal and tends to phthisis. This rule holds in the majority of cases, but there are many exceptions. There are cases in which glioma evolves under the form of iridocyclitis instead of glaucoma, where there is no increased tension whatever. Other cases directly opposite occur presenting the syndromes of glioma, amaurotic cat's eye and increased tension without the existence of any tumor. Transillumination and puncture are alone adequate to assure the diagnosis.

Benign Gangrene of the Eyelids.—ROGER and WEIL (*La Presse Médicale*, No. 76, 1901) report the case of a healthy male adult who was attacked with idiopathic gangrene of the left eyelids. There was no preceding inflammation of the conjunctiva nor any lesion of the lids. The disease began insidiously in a spot of inflammatory œdema. At the expiration of four or five

days general symptoms made their appearance, fever at first high (39.5°C .), but lasting only four days. It was accompanied by feeling of weight in the head and tension in the face, but without actual pain. There was a saburral condition of the digestive tract. The urine was albuminous. There was no delirium nor marked depression. Gangrenous plaques made their appearance upon the reddish erysipelatous lids forty-eight hours after the outbreak of the disease. The mortifying plaques were blackish in color and of very irregular contours. Line of demarcation quickly formed, and repair proceeded very rapidly. Eschars separated in fragments. The upper lid was free thereof on the nineteenth day; the lower on the twenty-first day. Healing was complete less than a month after the outbreak. There was no atresia of the palpebral orifice, and the only permanent trace was a slight ectropion at the external angle of the eye.

This is a very rare affection, indeed. The authors were able to find but a single analogous case in literature. It seems to resemble the spontaneous gangrene of the external male genital organs described by Fournier, of which more than twenty cases have been published since attention was first called to it.

Bacteriological study of the fluid obtained by puncturing the œdematous region showed the presence of an anaërobic microbe having a decided pathogenic action upon certain animals (rabbits, guinea-pigs), inoffensive for others (rats and mice). In the former, sometimes lethal septicæmia was produced, in other instances local lesions, œdemas, or indurations, or cold abscesses. True gangrene never occurred; but this is generally the case with all gangrene-producing microbes which do not retain or rapidly lose the virulence acquired for man.

DERMATOLOGY.

UNDER THE CHARGE OF

LOUIS A. DUHRING, M.D.,

PROFESSOR OF DERMATOLOGY IN THE UNIVERSITY OF PENNSYLVANIA,

AND

MILTON B. HARTZELL, M.D.,

INSTRUCTOR IN DERMATOLOGY IN THE UNIVERSITY OF PENNSYLVANIA.

Soft Pigmented Nævi.—FICK (*Archiv für Dermatologie und Syphilis*, Bd. lxx., Heft 111), who has studied the microscopical structure of eight pigmented nævi taken from various parts of the body, sums up the results of his investigation as follows: The pathological increase of pigment in a nævus is not the cause of the tumor formation. Nævus-cell formation and proliferation on the one side and increase of pigment on the other are two pathological processes running parallel courses, but also occurring separately. The nævus cells only contain pigment when they are found in close proximity to